

Effective Removal of Endocrine-Disrupting Compounds by Lignin Peroxidase from the White-Rot Fungus *Phanerochaete sordida* YK-624

著者	Wang Jianqiao, Majima Nayumi, Hirai Hirofumi, Kawagishi Hirokazu
journal or publication title	Current Microbiology
volume	64
number	3
page range	300-303
year	2011-12-28
出版者	Springer
権利	(C) Springer, Part of Springer Science+Business Media. The original publication is available at www.springerlink.com
URL	http://hdl.handle.net/10297/6730

doi: 10.1007/s00284-011-0067-2

Original Paper

**Effective removal of endocrine disrupting compounds by lignin peroxidase from
the white-rot fungus *Phanerochaete sordida* YK-624**

Jianqiao Wang,¹ Nayumi Majima,² Hirofumi Hirai,^{2*} Hirokazu Kawagishi^{1, 2}

¹ Graduate School of Science and Technology, Shizuoka University, 836 Ohya, Suruga-ku, Shizuoka
422-8529, Japan

² Department of Applied Biological Chemistry, Faculty of Agriculture, Shizuoka University, 836 Ohya,
Suruga-ku, Shizuoka 422-8529, Japan

Running title: Removal of ECDs by LiP

* Corresponding author: Hirofumi Hirai

Mailing address: Department of Applied Biological Chemistry, Faculty of Agriculture, Shizuoka
University, 836 Ohya, Suruga-ku, Shizuoka 422-8529, Japan.

Tel. & Fax: +81 54 238 4853

E-mail address: ahhirai@ipc.shizuoka.ac.jp

This work was partially supported by a Grant-in-Aid for Young Scientists (B) (No. 21780296) from the
Ministry of Education, Culture, Sports, Science and Technology of Japan.

Effective removal of endocrine disrupting compounds by lignin peroxidase from the white-rot fungus *Phanerochaete sordida* YK-624

The removal of endocrine disrupting compounds (EDCs) by lignin peroxidase from white-rot fungus *Phanerochaete sordida* YK-624 (YK-LiP1) was investigated. Five endocrine disruptors, *p*-t-octylphenol (OP), bisphenol A (BPA), estrone (E₁), 17 β -estradiol (E₂) and ethinylestradiol (EE₂) were eliminated by YK-LiP1 more effectively than lignin peroxidase from *P. chrysosporium* (Pc-LiP), and OP and BPA were disappeared almost completely in the reaction mixture containing YK-LiP1 after a 24-h treatment. Particularly, the removal of estrogenic activities of E₂ and EE₂, which show much higher estrogenic activities than other EDCs such as BPA and OP, were removed following 24-h treatment with YK-LiP1. Moreover, 5,5'-bis(1,1,3,3-tetramethylbutyl)-[1,1'-biphenyl]-2,2'-diol and 5,5'-bis-[1-(4-hydroxy-phenyl)-1-methyl-ethyl]-biphenyl-2,2'-diol were identified as the main metabolite from OP or BPA, respectively. These results suggest that YK-LiP1 is highly effective in removing of EDCs by the oxidative polymerization of these compounds.

1 **Introduction**

2 The occurrence of endocrine disrupting compounds (EDC) in the aquatic environment has
3 generated worldwide interest because these chemicals can cause feminization of fish as well as interfere
4 with the reproduction and development of other aquatic organisms [5, 12, 17]. Various natural and
5 synthetic chemical compounds have been identified as EDCs; including pharmaceuticals, pesticides,
6 industrial chemicals, and heavy metals [4]. Typical EDCs of anthropogenic origin with estrogen-like
7 action include *p*-*t*-octylphenol (OP) and bisphenol A (BPA). Natural estrogens, i.e., 17 β -estradiol (E₂)
8 and estrone (E₁), and synthetic estrogen, i.e., ethynylestradiol (EE₂), are excreted into wastewater by
9 humans and mammals mainly through their urine. The effluent concentrations of estrogens typically
10 range from a few ng/L to a few tens of ng/L [9, 11], but even these amounts are often high enough to
11 cause endocrine-disrupting effects in some aquatic species such as trouts [24] and minnows [16].
12 Estrogenic activities of estrogens are two or three orders of magnitude higher than those of EDCs such as
13 BPA [18, 23].

14 In recent years, white-rot fungi which can degrade lignin effectively are focused since white-rot
15 fungi can degrade various environmental pollutions such as polychlorinated dioxin [10], lindane [3],
16 heptachlor [27], trichlorobenzene [14] polycyclic aromatic hydrocarbons [1]. Moreover, ligninolytic
17 enzymes such as manganese peroxidase (MnP) and laccase were shown to be effective in removing the
18 estrogenic activities of bisphenol A, nonylphenol [25], 4-*tert*-octylphenol [21], and steroidal hormones
19 [20, 22]. However, the detail mechanisms on the detoxification of these compounds are still unknown.

20 White-rot fungus *Phanerochaete sordida* YK-624, which has been isolated from rotted wood,
21 showed much higher ligninolytic activity and selectivity than either *P. chrysosporium* or *Trametes*
22 *versicolor* [6]. The major extracellular ligninolytic enzymes of this strain are MnP [6] and LiP [13].
23 Particularly, this strain produces 2 novel LiP (YK-LiP1 and YK-LiP2), and these enzymes degrade lignin
24 model compounds more effectively than LiP from *P. chrysosporium* (Pc-LiP H8) [7, 19] . In the present
25 study, YK-LiP1 was applied to the removal of EDCs, and the removal properties were compared with
26 Pc-LiP H8. Moreover, the structures of metabolites from BPA and OP were determined to clarify the
27 removal mechanism of EDCs by YK-LiP1.

29 **Materials and methods**

30 **Fungi**

31 *P. sordida* YK-624 (ATCC 90872) and *P. chrysosporium* ME446 were used in this study. These
32 fungi were maintained on potato dextrose agar slants at 4°C.

Chemicals

BPA, E₁, E₂ and EE₂ were purchased from Tokyo Chemical Industry, Tokyo, Japan. OP was obtained from Wako Pure Chemical Industries, Osaka, Japan. All other chemicals were extra-pure grade and were used without further purification.

LiP preparation and determination of LiP activity

YK-LiP1 from *P. sordida* YK-624 and Pc-LiP (isozyme H8) from *P. chrysosporium* were prepared and purified as described by Sugiura et al. [19] and Wariishi and Gold [26], respectively. LiP activity was measured by monitoring the oxidation of veratryl alcohol (VA) to veratraldehyde ($\epsilon_{310} = 9.3 \text{ mM}^{-1}\text{cm}^{-1}$). The reaction mixture (1 ml) contained VA (1 mM) and H₂O₂ (0.2 mM) in 20 mM succinate buffer (pH 3.0). One katal (kat) was defined as the amount of enzyme producing 1 mol of product per second.

LiP treatment of EDCs

LiP reactions were performed in 1 ml of reaction mixture containing 2 nkat each LiP, 100 μM EDCs, and 100 μM H₂O₂ in 20 mM succinate, pH 3.0. Reactions were performed in triplicate for 24 h at 30°C and mixing at 150 rpm. The amount of EDCs was determined by high-performance liquid chromatography (HPLC) under the following conditions: column, Wakosil-II 5C18HG (4.6 mm x 150 mm, Wako Pure Chemical Industries, Japan); mobile phase, 50% aqueous acetonitrile containing 0.1% acetic acid (E₁, E₂, EE₂, and BPA) or 80% aqueous methanol containing 0.1% acetic acid (OP); flow rate, 1.0 ml/min; detection wavelength, 275 nm (BPA), 277 nm (OP), or 285 nm (E₁, E₂, and EE₂).

Estrogenic activity of E₁, E₂, or EE₂ treated with LiP

The estrogenic activities of E₁, E₂, or EE₂ before and after LiP treatment were evaluated by an *in vitro* screening test for chemicals with hormonal activities that used the yeast two-hybrid estrogenic assay system, developed by Nishikawa et al. [15]. The concentrations of E₁, E₂, and EE₂ before enzymatic treatment were 1 μM in the assay system (2.5 μl of reaction mixture containing 100 μM E₁, E₂, or EE₂ added to 50 μl of yeast culture and 200 μl of SD medium). Relative estrogenic activity (%) was defined as the percentage of β -galactosidase activity of enzyme-treated E₁, E₂, or EE₂ compared to that of untreated E₁, E₂, or EE₂.

Metabolism experiments

OP or BPA (final concentration 100 μM) were incubated at 30°C for 24 h in a 100-mL reaction mixture containing 200 nkat YK-LiP1 and 200 μM H₂O₂ in 20 mM succinic acid buffer, pH 3.0. The reaction mixtures were extracted 3 times with equal volume of ethyl acetate (EtOAc). The EtOAc extract were dried over anhydrous sodium sulfate and then evaporated to dryness. The concentrates were analyzed by thin-layer chromatography (TLC), HPLC, HR-ESI-MS and NMR. Silica gel plates (Merck

F254, Merck, Darmstadt, Germany) was used for analytical TLC. The metabolite of OP or BPA was further separated by HPLC (column: Wakosil-II 5C18HG) by 80% aqueous methanol containing 0.1% acetic acid or 70% aqueous methanol containing 0.1% acetic acid, respectively. The purified metabolites were then analyzed by HR-ESI-MS and NMR including COSY, HMQC, and HMBC experiments. The HR-ESI-MS data were measured by a JMS-T100LC mass spectrometer. ¹H-NMR spectra were recorded by a Jeol lambda-500 spectrometer at 500 MHz, while ¹³C-NMR spectra were recorded on the same instrument at 125 MHz.

Results and Discussion

We previously showed that YK-LiP1 from *P. sordida* YK-624 degrade dimeric lignin model compounds more effectively than Pc-LiP [7, 19]. Therefore, YK-LiP1 was applied to the removal of EDCs in the present study.

After a 24-h reaction using 2 nkat each LiP, the elimination of EDCs was determined (Fig. 1). Although YK-LiP1 belongs to LiP group, the substrate specificity of YK-LiP was different from that of Pc-LiP [7, 19]. Although same amounts (mole) of enzymes are popularly applied to these experiments, same activities of LiPs were used in this degradation experiment. Approximately 10% of LiP activities remained in each reaction mixture after 24-h. YK-LiP1 effectively removed OP, BPA, E₁, E₂, and EE₂. Particularly, OP, BPA, E₂, and EE₂ were disappeared almost completely in the reaction mixture containing YK-LiP1 whereas Pc-LiP removed OP, BPA, E₁, E₂, and EE₂ by 46.1%, 52.5%, 23.9%, 38.2%, and 45.0%, respectively. These results indicate that YK-LiP1 have a higher affinity than Pc-LiP for these phenolic compounds which were relatively high molecular weight (M.W. 206-296) since Huang et al. has reported that the activity of Pc-LiP toward various phenols is very low [8].

Because removal of toxicity is essential for the biodegradation of environmental pollutants, we examined estrogenic activities of E₁, E₂, and EE₂ treated with each LiP since these EDCs shows much higher estrogenic activities than OP or BPA [20, 21, 25]. Although the estrogenic activity of E₁ treated with YK-LiP1 was hardly decreased, treatments of E₂ and EE₂ by YK-LiP1 reduced the estrogenic activities by 72.6% and 82.6%, respectively (Fig. 2). On the other hands, estrogenic activities of E₁, E₂, and EE₂ treated with Pc-LiP were not reduced.

The removal of EDCs' estrogenic activities by ligninolytic enzymes from white rot fungi (MnP and laccase) has been reported [20, 21, 25]. However, there have been no studies focusing on the metabolic product of these EDCs by ligninolytic enzymes. Therefore, we attempted to identify the metabolites of OP and BPA, relatively simple structures compared with E₁, E₂, and EE₂. The metabolites were detected in

the analysis of TLC and HPLC (data not shown). The HR-ESI-MS data for the metabolite of OP, which yielded a molecular ion at m/z 409.3084 $[M-H]^-$ (calculated for $C_{28}H_{41}O_2$, 409.3061), indicated that the molecular formula of this compound was $C_{28}H_{42}O_2$. This formula suggested that the metabolite might be a dimer of OP. The structure of the purified metabolite was further characterized by NMR analyses. The ^{13}C -NMR and 1H -NMR spectra indicated that the metabolite of OP had four carbon atoms, one methylene, two methyls and 1,2,4 substituted benzene (data not shown). The 1,1',3,3'-tetramethylbutyl moiety was indicated by HMBC correlations (Fig. 3a) ($H-4,4'$, $H-2,2'/C-3,3'$, $H-1'-CH_3, H-2,2'/C-1,1'$). In addition, HMBC correlations ($H-1'-CH_3$, $H-2,2'$, $H-3,3'/C-5,5'$, $H-4,4', H-6,6'/C-1,1'$) confirmed the metabolite was 5,5'-bis(1,1',3,3'-tetramethylbutyl)-[1,1'-biphenyl]-2,2'-diol. The HR-ESI-MS data for the metabolite of BPA, which yielded a molecular ion at m/z 453.2062 $[M-H]^-$ (calculated for $C_{30}H_{30}O_4$, 453.2058), indicated that the molecular formula of this compound was $C_{30}H_{31}O_4$. This formula suggested that the metabolite might be a dimer of BPA. The structure of the purified metabolite was further characterized by NMR analyses. The ^{13}C -NMR and 1H -NMR spectra indicated that the metabolite of BPA had six carbon atoms, two methyls, five methylenes, 1,2,4 substituted benzene and 1,4 substituted benzene (data not shown). HMBC correlations (Fig. 3b) ($H-2',3'$ -methyl/ $C-1$, $C-2$, $C-5$, $H-3/C-1$, $C-2$, $C-5$) confirmed the metabolite was 5,5'-bis-[1-(4-hydroxy-phenyl)-1-methyl-ethyl]-biphenyl-2,2'-diol. Our current results suggest that each dimer was generated as a metabolite from OP or BPA by YK-LiP1. We propose that the formation of phenoxy radical followed by phenolic hydroxyl of OP was one-electron oxidized, and the radical was transferred to the *o*-position, the radical polymerization thus a dimer of OP was generated. BPA occurs a similar reaction was demonstrated, we proposed that the removal of the ECDs' estrogenic activities might be due to polymerization brought about by enzymatic oxidation.

Acknowledgement

This work was partially supported by a Grant-in-Aid for Young Scientists (B) (No. 21780296) from the Ministry of Education, Culture, Sports, Science and Technology of Japan.

References

- 1 Acevedo F, Pizzul L, Castillo Mdel P, Cuevas R & Diez MC (2011) Degradation of polycyclic aromatic hydrocarbons by the Chilean white-rot fungus *Anthracophyllum discolor*. *J Hazard Mater* **185**: 212-219.
- 2 Asgher M, Bhatti HN, Ashraf M & Legge RL (2008) Recent developments in biodegradation of

- industrial pollutants by white rot fungi and their enzyme system. *Biodegradation* **19**: 771–783.
- 3 Bumpus JA, Tien M, Wright D & Aust SD (1985) Oxidation of persistent environmental pollutants by a white rot fungus. *Science* **228**: 1434–1436.
- 4 Giesy JP, Hilscherova K, Jones PD, Kannan K & Machala M (2002) Cell bioassays for detections of aryl hydrocarbon (AhR) and estrogen receptor (ER) mediated activity in environmental samples. *Mar Pollut Bull* **45**: 3–16.
- 5 Harries JE, Sheahan DA, Jobling S, Matthiessen P, Neall P, Routledge EJ, Rycroft R, Sumpter JP & Tylor T (1996) A survey of estrogenic activity in United Kingdom inland waters. *Environ Toxicol Chem* **15**: 1993–2002.
- 6 Hirai H, Kondo R & Sakai K (1994) Screening of lignin-degrading fungi and their ligninolytic enzyme activities during biological bleaching of kraft pulp. *Mokuzai Gakkaishi* **40**: 980–986.
- 7 Hirai H, Sugiura M, Kawai S & Nishida T (2005) Characteristics of novel lignin peroxidases produced by white-rot fungus *Phanerochaete sordida* YK-624. *FEMS Microbiol Lett* **246**: 19–24.
- 8 Huang X, Wang D, Liu C, Hu M, Qu Y & Gao P (2003) The roles of veratryl alcohol and nonionic surfactant in the oxidation of phenolic compounds by lignin peroxidase. *Biochem Biophys Res Commun* **311**: 491–494.
- 9 Johnson AC & Sumpter JP (2001) Removal of endocrine-disrupting chemicals in activated sludge treatment works. *Environ Sci Technol* **35**: 4697–4703.
- 10 Kamei I, Suhara H & Kondo R (2005) Phylogenetical approach to isolation of white-rot fungi capable of degrading polychlorinated dibenzo-p-dioxin. *Appl Microbiol Biotechnol* **69**: 358–366.
- 11 Khanal SM, Xie B, Thompson ML, Sung S, Ong SK & Van Leeuwen J (2006) Fate, transport, and biodegradation of natural estrogens in the environment and engineered systems. *Environ Sci Technol* **40**: 6537–6546.
- 12 Larsson DGJ, Adolfsson-Erici M, Parkkonen J, Pettersson M, Berg AH, Olsson P & Forlin L (1999) Ethinylestradiol - an undesired fish contraceptive? *Aquat Toxicol* **45**: 91–97.
- 13 Machii Y, Hirai H & Nishida T (2004) Lignin peroxidase is involved in the biobleaching of manganese-less oxygen-delignified hardwood kraft pulp by white-rot fungi in the solid-fermentation system. *FEMS Microbiol Lett* **233**: 283–287.
- 14 Marco-Urrea E, Pérez-Trujillo M, Caminal G & Vicent T (2009) Dechlorination of 1,2,3- and 1,2,4-trichlorobenzene by the white-rot fungus *Trametes versicolor*. *J Hazard Mater* **166**: 1141–1147.
- 15 Nishikawa J, Saito K, Goto J, Dakeyama F, Matsuo M & Nishihara T (1999) New screening methods for chemicals with hormonal activities using interaction of nuclear hormone receptor with

- coactivator. *Toxicol Appl Pharmacol* **154**: 76-83.
- 16 Panter GH, Thompson RS & Sumpter JP (2000) Intermittent exposure of fish to estradiol. *Environ Sci Technol* **34**: 2756–2760.
- 17 Purdom CE, Hardiman PA, Bye VJ, Eno NC, Tyler CR & Sumpter J (1994) Estrogenic effects of effluents from sewage treatment works. *Chem Ecol* **8**: 275-285.
- 18 Routledge EJ & Sumpter JP (1996) Estrogenic activity of surfactants and some of their degradation products assessed using a recombinant yeast screen. *Environ Toxicol Chem* **15**: 241–248.
- 19 Sugiura M, Hirai H & Nishida T (2003) Purification and characterization of a novel lignin peroxidase from white-rot fungus *Phanerochaete sordida* YK-624. *FEMS Microbiol Lett* **224**: 285-290.
- 20 Suzuki K, Hirai H, Murata H & Nishida T (2003) Removal of estrogenic activities of 17 β -estradiol and ethinylestradiol by ligninolytic enzymes from white rot fungi. *Water Res* **37**: 1972-1975.
- 21 Tamagawa Y, Hirai H, Kawai S & Nishida T (2007) Removal of estrogenic activity of 4-*tert*-octylphenol by ligninolytic enzymes from white rot fungi. *Environ Toxicol* **22**: 281-286.
- 22 Tamagawa Y, Yamaki R, Hirai H, Kawai S & Nishida T (2006) Removal of estrogenic activity of natural steroidal hormone estrone by ligninolytic enzymes from white rot fungi. *Chemosphere* **65**: 97-101.
- 23 Tanaka H, Yakou Y, Takahashi A, Higashitani T & Komori K (2001) Comparison between estrogenicities estimated from DNA recombinant yeast assay and from chemical analyses of endocrine disruptors during sewage treatment. *Water Sci Technol* **43**: 125–132.
- 24 Thorpe KL, Hutchinson TH, Hetheridge MJ, Scholze M, Sumpter JP & Tyler CR (2001) Assessing the biological potency of environmental estrogens using vitellogenin induction in juvenile rainbow trout (*Oncorhynchus mykiss*). *Environ Sci Technol* **35**: 2476–2481.
- 25 Tsutsumi Y, Haneda T & Nishida T (2001) Removal of estrogenic activities of bisphenol A and nonylphenol by oxidative enzymes from lignin-degrading basidiomycetes. *Chemosphere* **42**: 271-276.
- 26 Wariishi H & Gold MH (1990) Lignin peroxidase compound III: mechanism of formation and decomposition. *J Biol Chem* **265**: 2070-2077.
- 27 Xiao P, Mori T, Kamei I & Kondo R (2011) Metabolism of organochlorine pesticide heptachlor and its metabolite heptachlor epoxide by white rot fungi, belonging to genus *Phlebia*. *FEMS Microbiol Lett* **314**: 140-146.

Figure Legends

Fig. 1. Decrease of BPA, OP, E₁, E₂ and EE₂ by YK-LiP1 and Pc-LiP. Reactions contained 2 nkat each LiP, 100 μM EDCs, and 100 μM H₂O₂ in 20 mM succinate, pH 3.0. Reactions were performed for 24 h at 30 °C and mixing at 150 r.p.m. Values are means ± SD of triplicate samples.

Fig. 2. Removal of estrogenic activities of E₁, E₂ and EE₂ by YK-LiP1 and Pc-LiP. Values are means ± SD of triplicate samples.

Fig. 3. HMBC correlations of the identified OP (a) and BPA (b) metabolites.




Fig. 1




Fig. 2





Fig. 3