Expression, purification and antigenicity of Neospora caninum-antigens using silkworm larvae targeting for subunit vaccines

著者	Otsuki Takahiro, Dong Jinhua, Kato Tatsuya, Park Enoch Y.
journal or publication title	Veterinary Parasitology
volume	192
number	1-3
page range	284-287
year	2013-02-18
出版者	Elsevier
権利	Copyright (C) 2012 Elsevier B.V. All rights reserved.
URL	http://hdl.handle.net/10297/6936

doi: 10.1016/j.vetpar.2012.09.038

**Expression, purification and antigenicity of *Neospora*
caninum-antigens using silkworm larvae targeting for
subunit vaccines**

Takahiro Otsuki ^a, Jinhua Dong ^a, Tatsuya Kato ^a and Enoch Y. Park ^{a,b,*}

^a *Laboratory of Biotechnology, Department of Applied Biological Chemistry, Faculty of
Agriculture, Shizuoka University, Shizuoka 422-8529, Japan*

^b *Laboratory of Biotechnology, Integrated Bioscience Section, Graduate School of
Science and Technology, Shizuoka University, 836 Ohya, Suruga-Ku, Shizuoka
422-8259, Japan*

* Corresponding author at: Integrated Bioscience Section, Graduate School of Science and Technology, Shizuoka University, 836 Ohya, Suruga-Ku, Shizuoka 422-8259, Japan. Tel. & Fax: +81-54-238-4887.

E-mail address: acypark@ipc.shizuoka.ac.jp (E.Y. Park).

ABSTRACT

Infection of *Neospora caninum* causes abortion in cattle, which has a serious worldwide impact on the economic performance of the dairy and beef industries. Now, inexpensive and efficacious vaccines are required to protect cattle from neosporosis in livestock industry. In this study, *N. caninum* surface antigen 1 (SAG1) and SAG1-related sequence 2 (SRS2) were expressed in hemolymph of silkworm larvae as a soluble form. Expressed SAG1 and SRS2 clearly showed antigenicity against *N. caninum*-positive sera of cow. SAG1 and SRS2 were purified to near homogeneity from hemolymph of silkworm larvae using anti-FLAG M2 antibody agarose: approximately 1.7 mg of SAG1 from 10 silkworm larvae and 370 µg of SRS2 from 17 silkworm larvae. Mice that were injected by antigens induced antibodies against SAG1 and SRS2. This study indicates that it is possible that this silkworm expression system leads to a large-scale production of *N. caninum*-antigens with biological function and low production cost. *Bombyx mori* nucleopolyhedrovirus (BmNPV) bacmid expression system paves the way to produce largely and rapidly these recombinant antigens for its application to subunit vaccines against neosporosis in cattle.

Keywords: *Neospora caninum*, neosporosis, silkworm, BmNPV bacmid

1. Introduction

Neospora caninum is a protozoan parasite of animals, which causes reproductive failure in cattle (Dubey et al., 2007). Neosporosis is a major cause of abortion in cattle and has serious impacts on the economic performance of dairy and beef industries (Reichel and Ellis, 2006). Vaccination is the most cost-effective way to control neosporosis, according to an economic analysis (Reichel and Ellis, 2006).

Subunit vaccines have been the most focused in this field. Various *N. caninum*-antigens have been reported and evaluated as a vaccine. *N. caninum* surface antigen 1 (SAG1), anchored on the surface of tachyzoites by glycosylphosphatidylinositol (GPI) anchor, was expressed in *E. coli* and purified recombinant SAG1 protected cerebral infection of *N. caninum* when immunized into mice (Cannas et al., 2003). SAG-related sequence protein 2 (SRS2) has also GPI anchor (Nishikawa et al., 2002), being localized on the surface of *N. caninum*, and is expressed in both tachyzoites and bradyzoites (Fuchs et al., 1998; Hemphill and Gottstein, 1996). Antibodies against SRS2 inhibit tachyzoite from attaching and invading to host cells and induce cellular and humoral immunity, supposing that SRS2 is a strong vaccine candidate (Baszler et al., 2008; Haldorson et al., 2006; Nishikawa et al., 2000).

Silkworms have been used for recombinant protein production because of its high capacity of producing proteins and cost-effectiveness for large-scale production (Kato et al., 2010; Li et al., 2010; Tsuji et al., 2011; Usami et al., 2010; Zhou et al., 2011).

Veterinary vaccines produced by recombinant baculoviruses are now on the market and some are under the development for licensing (Kamen et al., 2011). In addition, subunit vaccines produced in silkworms are immunogenic and efficacious in cattle when used as prophylactic ones (Li et al., 2008; Li et al., 2011). Especially, in these cases, a hemolymph of silkworm larvae containing expressed the recombinant protein was used as a subunit vaccine against infectious disease in cattle.

In this study, several antigens of *N. caninum* were expressed as FLAG-tagged proteins in silkworm larvae using BmNPV bacmid system. Purified antigens with an adjuvant induced antigen-specific antibodies in mice. This study demonstrates that BmNPV bacmid system can be applied to cost-effective large-scale production of subunit vaccines against cattle.

2. Materials and methods

2.1. Gene amplification, cloning, and construction of recombinant BmNPV bacmids

SAG1 or SRS2 was expressed in hemolymph of silkworm larvae using the signal peptide sequence of bombyxin from *Bombyx mori* (bx signal). bx signal peptide allows expressed proteins to be secreted into hemolymph of silkworm larvae efficiently (Park et al., 2007). Genomic DNA of *N. caninum* Nc-1 was purchased from American Type Culture Collection (ATCC No. 50843D) and used as a PCR template. Antigen genes, *sag1* and *srs2* have no intron in genomic DNA. Then *sag1* gene was amplified by PCR using SAG1-F (TATGGTACCGATCAGAAAAATCACCTCTA) and SAG1-R

(ATAGAGCTCTCACGCGACGCCAGCCGCTAT). *srs2* gene was amplified by PCR using SRS2-FL-F (TGCGGTACCGATTTCCTCGGGCAGTGAGAC) and SRS2-FL-R (ATAGAGCTCTCACGCGACGCCAGCCGCTAT). Each gene was cloned into pET52b vector at *Kpn* I-*Sac* I site. Next, each gene was amplified by PCR using CACC-bx-FLAG-HRV3C primer (CACCATGAAGATACTCCTTGCTATTGCATTAATGTTGTCAACAGTAATGTGG GTGTCAACAGACTACAAGGATGACGATGACAAGGGTGCACCTTGAAGTCCTC TTTCAG) and each reverse primer (SAG1-R or SRS2-FL-R). Each amplified gene was composed of bx signal peptide sequence, FLAG peptide sequence, human rhinovirus 3C protease cleavage site sequence, and each protein coding sequence. Each amplified gene was inserted into pENTR/D/TOPO (Life Technol. Japan Ltd, Tokyo, Japan) vector by TOPO reaction. A recombinant CP⁻ Chi⁻ BmNPV bacmid harboring each gene (BmNPV CP⁻ Chi⁻-bx-FLAGHRV3C-SAG1s or BmNPV CP⁻ Chi⁻-bx-FLAGHRV3C-SRS2FL) was constructed to express each *N. caninum*-antigen according to the previous report (Park et al., 2008).

2.2. Injection of BmNPV bacmid DNA into silkworm larvae, harvesting hemolymph of silkworm larvae, and purification of expressed recombinant *N. caninum*-antigens

BmNPV bacmid DNA injection into silkworm larvae and breeding silkworm larvae were performed according to the previous report (Park et al., 2008). Hemolymph was collected from silkworm larvae by cutting prolegs, and 1-phenyl-2-thiourea was put

into collected hemolymph at 0.1 mM to prevent melanization. Hemolymph was centrifuged at $10000 \times g$ for 15 min to remove hemocytes and debris, and its supernatant was used as a hemolymph sample.

To purify expressed recombinant *N. caninum*-antigens, 2 ml of anti-FLAG M2 antibody agarose (Sigma Aldrich Japan, Tokyo, Japan) was packed in an empty column and equilibrated with TBS (pH 7.5). Hemolymph diluted 5-fold with TBS was loaded onto the anti-FLAG M2 antibody agarose column. The column was washed with 40 ml of TBS after loading hemolymph, and proteins were eluted with 8 ml of glycine-HCl buffer (pH 3.5). Every fraction of 1 ml elute was collected.

2.3. SDS-PAGE, western blot and protein concentration determination

SDS-PAGE and western blot were carried out according to the previous report (Tsuji et al., 2011). Mouse anti-FLAG M2 antibody (Sigma Aldrich Japan) and goat anti-mouse labeled with horseradish peroxidase (HRP) (GE Healthcare Japan, Tokyo, Japan) were used as primary and secondary antibodies. Protein concentration was determined by BCA protein assay kit (Thermo SCIENTIFIC, Rockford, IL, USA).

2.4. Indirect enzyme-linked immunosorbent assay (Indirect ELISA)

One hundred microliters of hemolymph or purified antigens were immobilized on an ELISA plate at 4°C overnight. Solution in each well was removed, and 2% skim milk in PBST (PBS containing 0.05% Tween 20) as a blocking buffer was added into each well at room temperature for 2 hours for a blocking step, followed by collection

of their serum samples and washing each well using a plate washer (MODEL1575 ImmunoWash, Bio-Rad) with PBST. Then, 100 µl of *Neospora*-positive bovine serum (gifted by Mr. Junichi Noda of Shizuoka Prefecture Tobu Livestock Disease Diagnostic Center) diluted with 1000-fold with PBST was added to each well, and then, the ELISA plate was incubated at room temperature for an hour. Each well was washed using a plate washer with PBST, and 100 µl of goat anti-bovine IgG antibody-conjugated HRP (Jackson ImmunoResearch Lab. Inc.) was added into each well. After incubating the ELISA plate at room temperature for an hour, each well was washed using a plate washer with PBST. One hundred microliters of substrate [0.2-mg/ml 3,3',5,5'-tetramethylbenzidine (TMBZ) in 100-mM sodium acetate (pH 6.0), with 0.2% (v/v) of 30% hydrogen peroxide] were added to each well and left at room temperature for blue-color development. The reaction was stopped by the addition of 50 µl of 10% (w/v) H₂SO₄ solution. The developed color was measured at absorbance of 450 nm using a microplate reader (MODEL680, Bio-Rad).

2.5. Immunization of purified antigens to mice and collection of their serum samples

Purified recombinant *N. caninum*-antigens were dialyzed against PBS at 4°C overnight and mixed the same volume of Freund's complete adjuvant (Rockland Immunochemicals Inc., Gilbertsville, PA, USA). One hundred microliters of this mixture were immunized subcutaneously into a BALB/cCrSlc mouse (Japan SLC Inc., Hamamatsu, Japan). Immunization was performed every two weeks four times, and

200 µg of purified antigen was used for this immunization to a mouse.

After fourth immunization, blood was collected from the heart of the mouse and centrifuged at 3000 × g for 5 min. Sodium azide was added into this supernatant (serum) to 0.1%, and this mixture was kept at -30°C for further analysis.

3. Results and discussion

SAG1 and SRS2 have a signal peptide at its N-terminus and SAG1 also has a transmembrane domain at its C-terminus (Howe et al., 1998; Nishikawa et al., 2002). To express these antigens, *sag1* gene lacking the sequences of its native signal peptide and transmembrane domain region was amplified by PCR. Regarding SRS2, full-length *srs2* gene fused with bx signal peptide sequence was amplified by PCR. Each PCR product was connected with bx signal peptide sequence and FLAG tag sequence at its N-terminus to secrete each protein into hemolymph of silkworm larvae and make the purification of expressed protein easy.

SAG1 and SRS2 were expressed in hemolymph of silkworm larvae (Fig. 1A). Hemolymph containing expressed SAG1 and SRS2 had strong reactivity against serum from a *N. caninum*-infected cow than that from a *N. caninum*-negative cow (Fig. 1B), suggesting that SAG1, SRS2 are the major immunodominant antigens of *N. caninum* and candidates for effective vaccines against *N. caninum*-infection. SAG1 was also detected by Western blot using a *N. caninum*-positive serum (data not shown).

SAG1 and SRS2, which have antigenicity against a *N. caninum*-positive serum,

were purified from hemolymph of silkworm larvae using anti-FLAG M2 antibody agarose. Both SAG1 and SRS2 were purified (Fig. 2A) and its purity was higher than 90% when purified proteins were analyzed using Experion system (Bio-Rad). Approximately 1.7 mg of SAG1 and 370 µg of SRS2 were purified from 10 and 17 silkworm larvae, respectively. Protein yield in silkworm larvae is dependent on protein properties, for example, pI, hydrophobicity and structure etc. (Kato et al., 2010), likely to other protein expression system.

Purified SAG1 or SRS2 was immunized with Freund's complete adjuvant to mice 4 times every two weeks. Their sera were collected, and the production of antibodies against SAG1 or SRS2 was confirmed using indirect ELISA. Absorbance value higher than that in BSA was detected in both proteins, although both sera were diluted by more than 50000-fold (Fig. 2B). It indicates that both proteins induced antibodies against SAG1 and SRS2, respectively, and purified proteins from silkworm larvae have a potential to be used for a vaccine against neosporosis. Up to now, several *N. caninum*-antigens were expressed heterologously. SAG1 and SRS2 that are major immunodominant surface proteins in tachyzoites of *N. caninum* were expressed in *E. coli* as fusion proteins with poly-histidine tag. Purified antigens immunized into mice protected against cerebral *N. caninum* infection to some extent (Cannas et al., 2003).

Several antigens have been expressed in *E. coli* and cultured insect cells using recombinant baculoviruses and purified antigens have been evaluated as subunit vaccines to cattle. In this study, SAG1 and SRS2 were successfully expressed in

171 silkworm larvae using recombinant BmNPV bacmids. SAG1 and SRS2, expressed in
172 hemolymph of silkworm larvae and showed antigenicity to a *N. caninum*-positive
173 serum from cow. Purified SAG1 and SRS2 amounts were 1.7 and 0.37 mg from only
174 10-20 silkworm larvae, respectively. To use recombinant antigen for a vaccine to cattle,
175 large-scale antigen production system is needed inevitably. In the point of the
176 large-scale antigen production, silkworm system is advantageous. This point has a
177 tremendous impact on the production cost of subunit vaccines because any purification
178 steps of recombinant subunit vaccines from hemolymph of silkworm larvae are not
179 needed. BmNPV bacmid silkworm expression system for the production of *N.*
180 *caninum*-antigens provides its practical application as a recombinant antigen vaccine in
181 the field of livestock industry.

182 **Acknowledgements**

183 This work was supported by a Grant-in-Aid for Scientific Research (A) Grant
184 No.22248009 from the Ministry of Education, Culture, Sports, Science and Technology.
185 We thank Mr. Junichi Noda of Shizuoka Prefecture Tobu Livestock Disease Diagnostic
186 Center (101 Nitta Kannamicho Tagata-gun, Shizuoka Prefecture, Japan) for providing
187 cattle serum samples.

188 **References**

189 Baszler, T.V., Shkap, V., Mwangi, W., Davies, C.J., Mathison, B.A., Mazuz, M.,

190 Resnikov, D., Fish, L., Leibovitch, B., Staska, L.M., Savitsky, I., 2008. Bovine
 191 immune response to inoculation with *Neospora caninum* surface antigen SRS2
 192 lipopeptides mimics immune response to infection with live parasites. Clin
 193 Vaccine Immunol 15, 659-667.

194 Cannas, A., Naguleswaran, A., Muller, N., Eperon, S., Gottstein, B., Hemphill, A.,
 195 2003. Vaccination of mice against experimental *Neospora caninum* infection
 196 using NcSAG1- and NcSRS2-based recombinant antigens and DNA vaccines.
 197 Parasitology 126, 303-312.

198 Dubey, J.P., Schares, G., Ortega-Mora, L.M., 2007. Epidemiology and control of
 199 neosporosis and *Neospora caninum*. Clinical microbiology reviews 20,
 200 323-367.

201 Fuchs, N., Sonda, S., Gottstein, B., Hemphill, A., 1998. Differential expression of cell
 202 surface- and dense granule-associated *Neospora caninum* proteins in
 203 tachyzoites and bradyzoites. The Journal of parasitology 84, 753-758.

204 Haldorson, G.J., Stanton, J.B., Mathison, B.A., Suarez, C.E., Baszler, T.V., 2006.
 205 Neospora caninum: antibodies directed against tachyzoite surface protein
 206 NcSRS2 inhibit parasite attachment and invasion of placental trophoblasts in
 207 vitro. Experimental parasitology 112, 172-178.

208 Hemphill, A., Gottstein, B., 1996. Identification of a major surface protein on
 209 Neospora caninum tachyzoites. Parasitology research 82, 497-504.

210 Howe, D.K., Crawford, A.C., Lindsay, D., Sibley, L.D., 1998. The p29 and p35
 211 immunodominant antigens of Neospora caninum tachyzoites are homologous to
 212 the family of surface antigens of Toxoplasma gondii. Infection and immunity
 213 66, 5322-5328.

214 Kamen, A.A., Aucoin, M.G., Merten, O.W., Alves, P., Hashimoto, Y., Airene, K., Hu,
 215 Y.C., Mezzina, M., van Oers, M.M., 2011. An initiative to manufacture and
 216 characterize baculovirus reference material. Journal of invertebrate pathology
 217 107 Suppl, S113-117.

218 Li, X.H., Wang, D., Zhou, F., Yang, H.J., Bhaskar, R., Hu, J.B., Sun, C.G., Miao, Y.G.,
 219 2010. Cloning and expression of a cellulase gene in the silkworm, *Bombyx*
 220 *mori* by improved Bac-to-Bac/BmNPV baculovirus expression system.
 221 *Molecular biology reports* 37, 3721-3728.

222 Li, Z., Yi, Y., Yin, X., Zhang, Z., Liu, J., 2008. Expression of foot-and-mouth disease
 223 virus capsid proteins in silkworm-baculovirus expression system and its
 224 utilization as a subunit vaccine. *PloS one* 3, e2273.

225 Li, Z., Yin, X., Yi, Y., Li, X., Li, B., Lan, X., Zhang, Z., Liu, J., 2011. FMD subunit
 226 vaccine produced using a silkworm-baculovirus expression system: protective
 227 efficacy against two type Asia1 isolates in cattle. *Veterinary microbiology* 149,
 228 99-103.

229 Nishikawa, Y., Tragoolpua, K., Makala, L., Xuan, X., Nagasawa, H., 2002. *Neospora*
 230 *caninum* NcSRS2 is a transmembrane protein that contains a
 231 glycosylphosphatidylinositol anchor in insect cells. *Veterinary parasitology* 109,
 232 191-201.

233 Nishikawa, Y., Xuan, X., Nagasawa, H., Igarashi, I., Fujisaki, K., Otsuka, H., Mikami,
 234 T., 2000. Monoclonal antibody inhibition of *Neospora caninum* tachyzoite
 235 invasion into host cells. *International journal for parasitology* 30, 51-58.

236 Park, E.Y., Kageshima, A., Kwon, M.S., Kato, T., 2007. Enhanced production of
 237 secretory beta1,3-N-acetylglucosaminyltransferase 2 fusion protein into
 238 hemolymph of *Bombyx mori* larvae using recombinant BmNPV bacmid
 239 integrated signal sequence. *Journal of biotechnology* 129, 681-688.

240 Reichel, M.P., Ellis, J.T., 2006. If control of *Neospora caninum* infection is technically
 241 feasible does it make economic sense? *Veterinary parasitology* 142, 23-34.

242 Tsuji, Y., Deo, V.K., Kato, T., Park, E.Y., 2011. Production of Rous sarcoma virus-like
 243 particles displaying human transmembrane protein in silkworm larvae and its
 244 application to ligand-receptor binding assay. *Journal of biotechnology* 155,
 245 185-192.

246 Usami, A., Suzuki, T., Nagaya, H., Kaki, H., Ishiyama, S., 2010. Silkworm as a host of

247 baculovirus expression. Current pharmaceutical biotechnology 11, 246-250.

248 Zhou, Y., Chen, H., Li, X., Wang, Y., Chen, K., Zhang, S., Meng, X., Lee, E.Y., Lee,

249 M.Y., 2011. Production of recombinant human DNA polymerase delta in a

250 Bombyx mori bioreactor. PloS one 6, e22224.

251

252

Figure legends

Fig. 1. (A) Expression of SAG1 and SRS2 in hemolymph of silkworm larvae. Protein expression was analyzed by Western blot using mouse anti-FLAG M2 antibody. (B) Antigenicity of SAG1 and SRS2 expressed in hemolymph of silkworm larvae against a *N. caninum*-positive serum. Antigenicity of each antigen was analyzed using indirect ELISA as described in Materials and methods. Closed and open bars indicate *Neospora*-positive and negative sera, respectively.

Fig. 2. (A) Purification of SAG1 and SRS2 from hemolymph of silkworm larvae using anti-FLAG M2 antibody agarose. Lanes of CBB and WB show SDS-PAGE with Coomassie Brilliant Blue (CBB) staining and Western blot analysis of each purified antigen using mouse anti-FLAG M2 antibody, respectively. (B) Indirect ELISA analysis of serum from mice immunized with purified SAG1 (I) or SRS2 (II). Indirect ELISA method was described in Materials and methods in detail.

(A)

(B)

(A) **(B)**