Hydroxylation of bisphenol A by hyper lignin-degrading fungus *Phanerochaete sordida* YK-624 under non-ligninolytic condition

著者	Wang Jianqiao, Yamamoto Ryoko, Yamamoto Yotaro, Tokumoto Toshinobu, Dong Jing, Thomas Peter, Hirai Hirofumi, Kawagishi Hirokazu
journal or publication title	Chemosphere
volume	93
number	7
page range	1419-1423
year	2013-10
出版者	Elsevier
権利	Copyright (C) 2013 Elsevier Ltd. Published by Elsevier Ltd. All rights reserved.
URL	http://hdl.handle.net/10297/7430

doi: 10.1016/j.chemosphere.2013.07.026

**Hydroxylation of bisphenol A by hyper lignin-degrading fungus *Phanerochaete sordida*
YK-624 under non-ligninolytic condition**

Jianqiao Wang^a, Ryoko Yamamoto^b, Yotaro Yamamoto^c, Toshinobu Tokumoto^a, Jing Dong^d,
Peter Thomas^d, Hirofumi Hirai^{c, e*}, Hirokazu Kawagishi^{a, c, e}

^a Graduate School of Science and Technology, Shizuoka University, 836 Ohya, Suruga-ku, Shizuoka 422-8529, Japan

^b Department of Applied Biological Chemistry, Faculty of Agriculture, Shizuoka University, 836 Ohya, Suruga-ku, Shizuoka 422-8529, Japan

^c Graduate School of Agriculture, Shizuoka University, 836 Ohya, Suruga-ku, Shizuoka 422-8529, Japan

^d Marine Science Institute, University of Texas at Austin, 750 Channel View Drive, Port Aransas, TX 78373, USA

^e Research Institute of Green Science and Technology, Shizuoka University, 836 Ohya, Suruga-ku, Shizuoka 422-8529, Japan

Key words: *Phanerochaete sordida* YK-624; Bisphenol A; Hydroxylation; Cytochrome P450

* Corresponding author: Hirofumi Hirai

Mailing address: Graduate School of Agriculture, Shizuoka University, 836 Ohya, Suruga-ku, Shizuoka 422-8529, Japan.

Tel. & Fax: +81 54 238 4853

E-mail address: ahhirai@ipc.shizuoka.ac.jp

ABSTRACT

1 Bisphenol A (BPA) is one of the representative compounds of the endocrine disrupting
2 compounds group and the highest volume chemicals produced worldwide. As a result, BPA is
3 often detected in many soil and water environments. In this study, we demonstrated the
4 transformation of BPA from liquid cultures inoculated with hyper lignin-degrading fungus
5 *Phanerochaete sordida* YK-624. Under non-ligninolytic conditions, approximately 80% of
6 BPA was eliminated after 7 d of incubation. High-resolution electrospray ionization mass
7 spectra and nuclear magnetic resonance analyses of a metabolite isolated from the culture
8 supernatant suggested that BPA was metabolized to hydroxy-BPA,
9 4-(2-(4-hydroxyphenyl)propan-2-yl)benzene-1,2-diol, which has a much lower estrogenic
10 activity than BPA. In addition, we investigated the effect of the cytochrome P450 inhibitor
11 piperonyl butoxide (PB) on the hydroxylation of BPA, markedly lower transformation activity
12 of BPA was observed in cultures containing PB. These results suggest that cytochrome P450
13 plays an important role in the hydroxylation of BPA by *P. sordida* YK-624 under
14 non-ligninolytic conditions.

1. Introduction

The occurrence of endocrine disrupting compounds (EDCs) in the aquatic environment has generated worldwide interest because these chemicals can cause feminization of fish as well as interfere with the reproduction and development of other aquatic organisms (Purdum et al., 1994; Harries et al., 1996; Larsson et al., 1999). One of the representative compounds of the EDCs group is bisphenol A (2,2-bis(4-hydroxyphenyl)propane; BPA). BPA, a monomer component of polycarbonate plastics, is used in many consumer products, including lacquers applied as food-can linings and dental composite fillings and sealants (Brotons et al., 1995; Olea et al., 1996). Many countries throughout the world have large production capacities for BPA, especially Germany, the Netherlands, the USA and Japan. BPA is one of the highest volume chemicals produced worldwide, with an estimated production of 5 Mt in 2010 (Ballesteros-Gómez et al., 2009). As a result, BPA has been detected often in many soil and water environments (Kitada et al., 2008; Matsumura et al., 2009).

Lignin-degrading white-rot fungi, which have the unique ability to degrade lignin to the level of CO₂ (Kirk and Farrell, 1987), and their ligninolytic enzymes have also attracted interest for the biotransformation of contaminants because of their industrial potential for degrading recalcitrant environmental pollutants, such as polychlorinated dibenzodioxin (Kamei et al., 2005), lindene (Bumpus et al., 1985), chlorophenols (Joshi and Gold, 1993), mycotoxin aflatoxin B₁ (Wang et al., 2011), and acetamiprid as a neonicotinoid insecticide (Wang et al., 2012a).

The degradation of EDCs by ligninolytic enzymes has attracted the attention of many researchers (Hirano et al., 2000; Suzuki et al., 2003; Saito et al., 2004; Tamagawa et al., 2006). The degradation of BPA using fungi is also reported (Chai et al., 2005; Shin et al., 2007) but the detail mechanisms are still unknown. More recently, we have reported that the removal of BPA by lignin peroxidase from hyper lignin-degrading fungus *Phanerochaete sordida* YK-624 (Wang et al., 2012b) and *P. sordida* YK-624 under ligninolytic condition (Wang et al., 2013). In our previous reports, BPA dimers have been formed as metabolites by radical couplings of BPA phenoxy radical generated by ligninolytic enzymes. Unfortunately, these BPA dimers are possible to be decomposed to toxic BPA by various reactions. Therefore, the transformation techniques of BPA without polymerization are necessary. In the present study, we examined the removal of BPA by *P. sordida* YK-624 under non-ligninolytic condition which hardly oxidize BPA to BPA phenoxy radical. We also detected the metabolite from BPA and proposed the metabolic pathway of BPA by *P. sordida* YK-624.

2. Materials and methods

2.1. Fungi and chemicals

P. sordida YK-624 (ATCC 90872), which has been isolated from rotted wood, was used in the present study (Hirai et al., 1994). The fungus was maintained on potato dextrose agar (PDA) slants at 4 °C.

BPA was purchased from Alfa Aesar, A Johnson Matthey (Lancashire, United Kingdom). All other chemicals were analytical pure grade and were used without further purification.

2.2. Transformation of BPA

Potato dextrose broth (PDB) medium which is a non-lignolytic (no production of lignolytic enzymes) condition was used for BPA transformation experiments. The fungus was incubated on a PDA plate at 30 °C for 3 d, and then 10 mm-diameter disks were punched out from the growing edge of mycelium. Two disks were each placed into a 100-mL Erlenmeyer flask containing 10 mL of PDB medium. After statically incubating the flasks at 30 °C for 3 d, 100 µL of 100 mM BPA/dimethyl sulfoxide were added to the cultures (final concentration at 1 mM), and then further incubated for 0-7 d (each in triplicate). The internal standard material was added to the culture, and then the culture was homogenized with 20 mL of acetone by using a Polytron PT1200E (Kinematica, Canada). The homogenate was filtrated and then evaporated to dryness. The residue was analyzed by HPLC for the quantification of BPA under the following conditions: column, Wakosil-II 5C18HG (4.6×150 mm; Wako Pure Chemical Industries, Japan); mobile phase, 10 to 100% MeOH gradient for 60 min; flow rate, 1 mL min⁻¹; and UV wavelength, 277 nm.

2.3. Metabolism experiment of BPA

Inoculated cultures of *P. sordida* YK-624 were prepared under the PDB medium. After static incubation at 30 °C for 3 d, BPA (final concentration, 1 mM) was added to these cultures. These cultures were further incubated for 7 d and then filtrated with membrane filters (pore size; 0.2 µm). The resulting solution was extracted 3 times with equal amount of ethyl acetate (EtOAc). The EtOAc extract was dried over anhydrous sodium sulfate and then evaporated to dryness. The residue was cleaned on a thin-layer chromatography (TLC, glass plates) to obtain 3 fractions. Each fraction was analyzed by TLC (aluminum sheets). Silica gel plates (Merck F₂₅₄; Merck, Darmstadt, Germany) were used for TLC. The metabolite was further separated by HPLC (column: CAPCELL PAK C18 AQ; 20×250 mm, Shiseido,

Japan) using 10 to 100% MeOH gradient. The purified metabolite was analyzed by HR-ESI-MS and NMR, including distortionless enhancement by polarization transfer, correlated spectroscopy, heteronuclear multi quantum correlation, and heteronuclear multiple bond correlation (HMBC) experiments, respectively. The HR-ESI-MS data were measured using a JMS-T100LC mass spectrometer. ¹H-NMR spectra were recorded using a Jeol Lambda-500 spectrometer at 500 MHz, while ¹³C-NMR spectra were recorded on the same instrument at 125 MHz.

2.4. Cytochrome P450 inhibitor experiment

After preculturing *P. sordida* YK-624 for 3 d, BPA (final concentration at 1 mM) and the cytochrome P450 inhibitor piperonyl butoxide (PB, final concentration, 0, 0.1 and 1 mM) were added to cultures. The cultures were further incubated for 1-4 d, and each culture was then subjected to HPLC for the quantification of BPA, as described above.

2.5. Competition assay using membrane estrogen receptor

Culture of HEK231 cells were stably transfected with membrane estrogen receptor (GPR30) and estrogen receptor competition studies were carried out as described before (Thomas and Dong, 2006). Briefly, human HEK293 cells stably transfected with human GPR30 were cultured in dulbecco's modification of eagle's medium/Ham's F-12 medium with 5% fetal bovine serum and 100 µg mL⁻¹ of G418 in 150 mm diameter plates. The medium was replaced every 1-2 d and the cells which were reached 80% confluence after 3 d were used in the experiments.

Procedures for the preparation of plasma membrane fractions and assay of steroid membrane receptors were followed with few modifications for measurement of [³H] estradiol (E₂) binding to plasma membranes of HEK293 cells transfected with goldfish mPRα (Tokumoto et al., 2007) except that presoaking with Tween 80 was omitted. As a radio-labeled steroid [2, 4, 6, 7-³H]E₂ ([³H]E₂, ~89 Ci mmol⁻¹ was purchased from PerkinElmer Life & Analytical Sciences (Waltham, MA). The competitive binding assay tubes contained 4 nM [³H]E₂ and the competitors (concentration range: 1 nM to 10 µM). After a 30 min incubation at 4 °C with the membrane fractions, the reaction was stopped by filtration (Whatman GF/B filters), the filters were washed and bound radioactivity measured by scintillation counting. The displacement of [³H]E₂ binding by the competitors was expressed as a percentage of the maximum specific binding of E₂.

3. Results

3.1. Transformation of BPA by *P. sordida* YK-624 under non-ligninolytic condition

In the present study, PDB medium was used for the transformation experiment of BPA. When *P. sordida* YK-624 was cultured in PDB medium, BPA was decreased by 51, 71, 83% after 1, 4, and 7 d of incubation, respectively (Fig. 1a). These results suggested that *P. sordida* YK-624 had the ability for the transformation of BPA under the non-ligninolytic conditions such as PDB medium.

3.2. Identification of the metabolite from BPA

To determine the metabolite(s) produced during the degradation of BPA, 7-d cultures of *P. sordida* YK-624 in PDB medium was supplemented with 0.5 mM BPA were subjected to TLC and HPLC. From this analysis, one potential BPA metabolite was detected (data not shown). The purified metabolite was then subjected to high-resolution electrospray ionization mass spectrometry analysis, which yielded a molecular ion at m/z 243.1015 $[M-H]^-$ (calculated for $C_{15}H_{15}O_3$, 243.1021), indicating that the molecular formula of this compound was $C_{15}H_{16}O_3$. This formula suggested that the metabolite might be a hydroxylation form of BPA. The structure of the purified metabolite was further characterized by NMR analyses. Table 1 lists the chemical-shift assignment data for the metabolite of BPA. HMBC (Fig. 2) (H-3/C-2', H-5/C-2') confirmed that the metabolite was 4-(2-(4-hydroxyphenyl)propan-2-yl)benzene-1,2-diol, which hydroxylated at the *ortho*-position of BPA.

3.3. Effect of cytochrome P450 inhibitor

The effect of the cytochrome P450 inhibitor on the transformation of BPA by *P. sordida* YK-624 was investigated using PB. The transformation rate of BPA after the addition of 0.1 and 1 mM PB into cultures of *P. sordida* YK-624 is shown in Fig. 1b. In contrast to the transformation rate of BPA in cultures without PB, markedly lower transformation activity of BPA was observed in cultures containing PB. In the presence of 0.1 and 1 mM PB, 63 and 38% of BPA were eliminated after 4 d of incubation, respectively, compared to 70% in cultures lacking PB.

3.4. Steroid binding assay of BPA and the metabolite

To evaluate the estrogenic activity of BPA metabolite, we used the steroid binding assay in the present study. BPA showed significant competition with IC_{50} of 1.8 μ M. Just a few affinities for binding to the membrane estrogen receptor were detected in hydroxy-BPA. These

results demonstrated that hydroxylation of BPA by *P. sordida* YK-624 causes the 67% loss of estrogenic activity of BPA (Fig. 3). In other words hydroxylation of BPA by *P. sordida* YK-624 is sufficient for diminish possible environmental estrogenic activity of BPA.

4. Discussion

Biotransformation of BPA has also been reported by many researchers. More recently, we reported the removal of BPA by *P. sordida* YK-624 under ligninolytic condition (Wang et al., 2013). BPA dimers are formed by radical couplings at extracellular region without the incorporation into the cell of *P. sordida* YK-624. However, BPA dimers may be returned to BPA by various reactions. BPA glucuronide is the major metabolite of BPA in rat hepatocytes (Nakagawa and Tayama, 2000; Pritchett et al., 2002). In plants, cells of *Eucalyptus perriniana* and *Portulaca oleracea* cv. metabolize BPA to its hydroxyl products (Hamada et al., 2002; Watanabe et al., 2012). In fungi, BPA is metabolized to BPA-*O*- β -D-glucopyranoside by *Aspergillus fumigatus* (Yim et al., 2003). In the present study, we detected a hydroxy-BPA as a metabolite of BPA.

Watanabe et al. (2012) suggested polyphenol oxidases were likely to contribute to BPA hydroxylation by portulaca. However, *P. sordida* YK-624 produces lignin peroxidases and manganese peroxidase as ligninolytic enzymes but not laccase. The functional diversity of cytochrome P450s in white-rot fungi has been studied because cytochrome P450s play an important role on degrading a wide variety of recalcitrant aromatic compounds (Hiratsuka et al. 2001; Xiao et al., 2011; Wang et al., 2012a). Recently, Hata et al. (2010) suggested that hydroxylation catalyzed by cytochrome P450 in *P. sordida* YK-624 might be involved in the elimination and detoxification of diclofenac and mefenamic acid. Our present study showed that the transformation of BPA was efficiently inhibited by the addition of PB (Fig. 1b), which is a common inhibitor of cytochrome P450 and is often used for demonstrating whether a reaction is catalyzed by cytochrome P450 enzymes (Mori and Kondo, 2002; Mori et al., 2003; Kamei et al., 2005). Since hydroxylation of BPA was inhibited by PB, we propose that cytochrome P450s are involved in the hydroxylation of BPA by *P. sordida* YK-624 under non-ligninolytic condition. We have described for the first time the hydroxylation of BPA by a white-rot fungus.

Due to the high production volumes, the toxicity of BPA has been intensively studied since the 1970s. Estrogenic activity of BPA was first reported in 1993 (Krishnan et al., 1993). We used the steroid binding assay to evaluate the estrogenic activity of BPA and BPA metabolite in the present study. BPA showed significant competition with IC₅₀ of 1.8 μ M. Just

a few affinities for binding to the membrane estrogen receptor were detected in hydroxy-BPA (Fig. 3). Since the metabolite hydroxy-BPA has lower estrogenic toxicity than BPA, diminishing environmental estrogenic activity of BPA is possible using *P. sordida* YK-624.

The findings of the present study confirm that *P. sordida* YK-624 can remove the estrogenic activity of BPA by converting to hydroxy-BPA. In the next study, we will investigate whether hydroxy-BPA could be further metabolized by *P. sordida* YK-624 for complete removal of estrogenic activity.

5. Acknowledgement

This work was partially supported by Grants-in-Aid for Scientific Research (No. 21780296) from the Ministry of Education, Culture, Sports, Science and Technology of Japan.

6. References

- Ballesteros-Gómez, A., Rubio, S., Pérez-Bendito, D., 2009. Analytical methods for the determination of bisphenol A in food. *J. Chromatogr. A* 1216, 449-469.
- Brotons, J.A., Olea-Serrano, M.F., Villalobos, M., Pedraza, V., Olea, N., 1995. Xenoestrogens released from lacquer coatings in food cans. *Environ. Health Persp.* 103, 608-612.
- Bumpus, J.A., Tien, M., Wright, D., Aust, S.D., 1985. Oxidation of persistent environmental pollutants by a white rot fungus. *Science* 228, 1434-1436.
- Chai, W., Handa, Y., Suzuki, M., Saito, M., Kato, N., Horiuchi, C.A., 2005. Biodegradation of bisphenol A by fungi. *Appl. Biochem. Biotechnol.* 120, 175-182.
- Hamada, H., Tomi, R., Asada, Y., Furuya, T., 2002. Phytoremediation of bisphenol A by cultured suspension cells of *Eucalyptus perriniana*-regioselective hydroxylation and glycosylation. *Tetrahedron Lett.* 43, 4087-4089.
- Harries, J.E., Sheahan, D.A., Jobling, S., Matthiessen, P., Neall, P., Routledge, E.J., Rycroft, R., Sumpter, J.P., Tylor, T., 1996. A survey of estrogenic activity in United Kingdom inland waters. *Environ. Toxicol. Chem.* 15, 1993-2002.
- Hata, T., Kawai, S., Okamura, H., Nishida, T., 2010. Removal of diclofenac and mefenamic acid by the white rot fungus *Phanerochaete sordida* YK-624 and identification of their metabolites after fungal transformation. *Biodegradation* 21, 681-689.
- Hirai, H., Kondo, R., Sakai, K., 1994. Screening of lignin-degrading fungi and their ligninolytic enzyme activities during biological bleaching of kraft pulp. *Mokuzai Gakkaishi* 40, 980-986.

205 Hirano, T., Honda, Y., Watanabe, T., Kuwahara, M., 2000. Degradation of bisphenol A by
 206 the lignin-degrading enzyme, manganese peroxidase, produced by the white-rot
 207 basidiomycete, *Pleurotus ostreatus*. Biosci. Biotechnol. Biochem. 64, 1958-1962.

208 Hiratsuka, N., Wariishi, H., Tanaka, H., 2001. Degradation of diphenyl ether herbicides
 209 by the lignin-degrading basidiomycete *Coriolus versicolor*. Appl. Microbiol. Biotechnol. 57,
 210 563-571.

211 Joshi, D.K., Gold, M.H., 1993. Degradation of 2,4,5-trichlorophenol by the
 212 lignin-degrading basidiomycete *Phanerochaete chrysosporium*. Appl. Environ. Microbiol. 59,
 213 1779-1785.

214 Kamei, I., Suhara, H., Kondo, R., 2005. Phylogenetical approach to isolation of white-rot
 215 fungi capable of degrading polychlorinated dibenzo-*p*-dioxin. Appl. Microbiol. Biotechnol.
 216 69, 358-366.

217 Kirk, T.K., Farrell, R.L., 1987. Enzymatic combustion: the microbial degradation of
 218 lignin. Annu. Rev. Microbiol. 41, 465-505.

219 Krishnan, A.V., Stathis, P., Permuth, S.F., Tokes, L., Feldman, D., 1993. Bisphenol-A: an
 220 estrogenic substance is released from polycarbonate flasks during autoclaving. Endocrinology
 221 132, 2279-2286.

222 Kitada, Y., Kawahata, H., Suzuki, A., Oomori, T., 2008. Distribution of pesticides and
 223 bisphenol A in sediments collected from rivers adjacent to coral reefs. Chemosphere 71,
 224 2082-2090.

225 Larsson, D.G.J., Adolfsson-Erici, M., Parkkonen, J., Pettersson, M., Berg, A.H., Olsson,
 226 P., Forlin, L., 1999. Ethinyloestradiol - an undesired fish contraceptive? Aquat. Toxicol. 45,
 227 91-97.

228 Matsumura, Y., Hosokawa, C., Sasaki-Mori, M., Akahira, A., Fukunaga, K., Ikeuchi, T.,
 229 Oshiman, K., Tsuchido, T., 2009. Isolation and characterization of novel
 230 bisphenol-A--degrading bacteria from soils. Biocontrol Sci. 14, 161-169.

231 Mori, T., Kondo, R., 2002. Oxidation of chlorinated dibenzo-*p*-dioxin and dibenzofuran
 232 by white-rot fungus, *Phlebia lindtneri*. FEMS Microbiol. Lett. 216, 223-227.

233 Mori, T., Kitano, S., Kondo, R., 2003. Biodegradation of chloronaphthalenes and
 234 polycyclic aromatic hydrocarbons by the white-rot fungus *Phlebia lindtneri*. Appl. Microbiol.
 235 Biotechnol. 61, 380-383.

236 Nakagawa, Y., Tayama, S., 2000. Metabolism and cytotoxicity of bisphenol A and other
 237 bisphenols in isolated rat hepatocytes. Arch. Toxicol. 74, 99-105.

238 Olea, N., Pulgar, R., Pérez, P., Olea-Serrano, F., Rivas, A., Novillo-Fertrell, A., Pedraza,

239 V., Soto, A.M., Sonnenschein, C., 1996. Estrogenicity of resin-based composites and sealants
240 used in dentistry. *Environ. Health Persp.* 104, 298-305.

241 Pritchett, J.J., Kuester, R.K., Sipes, I.G., 2002. Metabolism of bisphenol a in primary
242 cultured hepatocytes from mice, rats, and humans. *Drug Metab. Dispos.* 30, 1180-1185.

243 Purdom, C.E., Hardiman, P.A., Bye, V.J., Eno, N.C., Tyler, C.R., Sumpter, J., 1994.
244 Estrogenic effects of effluents from sewage treatment works. *Chem. Ecol.* 8, 275-285.

245 Saito, T., Kato, K., Yokogawa, Y., Nishida, M., Yamashita, N., 2004. Detoxification of
246 bisphenol A and nonylphenol by purified extracellular laccase from a fungus isolated from
247 soil. *J. Biosci. Bioeng.* 98, 64-66.

248 Shin, E.H., Choi, H.T. Song, H.G., 2007. Biodegradation of endocrine-disrupting
249 bisphenol A by white rot fungus *Irpex lacteus*. *J. Microbiol. Biotechnol.* 17, 1147-1151.

250 Suzuki, K., Hirai, H., Murata, H., Nishida, T., 2003. Removal of estrogenic activities of
251 17 β -estradiol and ethinylestradiol by ligninolytic enzymes from white rot fungi. *Water Res.*
252 37, 1972-1975.

253 Tamagawa, Y., Yamaki, R., Hirai, H., Kawai, S., Nishida, T., 2006. Removal of
254 estrogenic activity of natural steroidal hormone estrone by ligninolytic enzymes from white
255 rot fungi. *Chemosphere* 65, 97-101.

256 Thomas, P., Dong, J., 2006. Binding and activation of the seven-transmembrane estrogen
257 receptor GPR30 by environmental estrogens: a potential novel mechanism of endocrine
258 disruption. *J. Steroid. Biochem.* 102, 175-179.

259 Tokumoto, T., Tokumoto, M., Thomas P., 2007. Interactions of diethylstilbestrol (DES)
260 and DES analogs with membrane progesterin receptor-alpha and the correlation with their
261 nongenomic progesterin activities. *Endocrinology* 148, 3459-3467.

262 Wang, J., Hirai, H., Kawagishi, H., 2012a. Biotransformation of acetamiprid by the
263 white-rot fungus *Phanerochaete sordida* YK-624. *Appl. Microbiol. Biotechnol.* 93, 831-835.

264 Wang, J., Majima, N., Hirai, H., Kawagishi, H., 2012b. Effective removal of
265 endocrine-disrupting compounds by lignin peroxidase from the white-rot fungus
266 *Phanerochaete sordida* YK-624. *Curr. Microbiol.* 64, 300-303.

267 Wang, J., Ogata, M., Hirai, H., Kawagishi, H., 2011. Detoxification of aflatoxin B₁ by
268 manganese peroxidase from the white-rot fungus *Phanerochaete sordida* YK-624. *FEMS*
269 *Microbiol. Lett.* 314, 164-169.

270 Wang, J., Yamamoto, Y., Hirai, H., Kawagishi, H., 2013. Dimerization of bisphenol A by
271 hyper lignin-degrading fungus *Phanerochaete sordida* YK-624 under ligninolytic condition.
272 *Curr. Microbiol.* 66, 544-547.

273 Watanabe, I., Harada, K., Matsui, T., Miyasaka, H., Okuhata, H., Tanaka, S., Nakayama,
274 H., Kato, K., Bamba, T., Hirata, K., 2012. Characterization of bisphenol A metabolites
275 produced by *Portulaca oleracea* cv. by liquid chromatography coupled with tandem mass
276 spectrometry. Biosci. Biotechnol. Biochem. 76, 1015-1017.

277 Xiao, P., Mori, T., Kamei, I., Kondo, R., 2011. A novel metabolic pathway for
278 biodegradation of DDT by the white rot fungi, *Phlebia lindtneri* and *Phlebia brevispora*.
279 Biodegradation 22, 859-867.

280 Yim, S.H., Kim, H.J., Lee, I.S., 2003. Microbial metabolism of the environmental
281 estrogen bisphenol A. Arch. Pharm. Res. 26, 805-808.
282

Fig. 1 Time course for BPA transformation by *P. sordida* YK-624 under non-ligninolytic condition (a) and effect of cytochrome P450 inhibitor PB on the transformation of BPA by *P. sordida* YK-624 (b). ● without PB, ▲ 0.1 mM PB, ■ 1 mM PB. Values are the means $\pm$ SD of triplicate samples.

Fig. 2 COSY and HMBC correlations for the identified BPA metabolite.

Fig. 3 Competition by BPA and hydroxy-BPA for binding to the membrane estrogen receptor. Samples were incubated with 4 nM [³H]E2 and 10 μ M competitor. Competition rate (%) is represented as relative value of the displacement of [³H]E2 binding by BPA set as 100 %.

Fig. 1

4-(2-(4-hydroxyphenyl)propan-2-yl)benzene-1,2-diol

Fig. 2

Fig. 3

Table 1 ^1H - and ^{13}C -NMR data for BPA metabolite (in CD_3OD)

Position	^1H	^{13}C
	δ_{H} (mult, J in Hz)	d
1	-	142.3
2	-	142.9
3	6.61 (d, 2.2)	114.1
4	-	141.9
5	6.56 (dd, 7.9, 2.2)	117.3
6	6.64 (d, 7.9)	113.9
1''	-	143.9
2''	7.02 (d, 9.0)	127.2
3''	6.65 (d, 8.5)	113.9
4''	-	154.3
5''	6.65 (d, 8.5)	113.9
6''	7.02 (d, 9.0)	127.2
1'-CH ₃	1.54 (s)	30.1
2'	-	41.0
3'-CH ₃	1.54 (s)	30.0