Improving xylitol production through recombinant expression of xylose reductase in the white-rot fungus *Phanerochaete sordida* YK-624

著者	Hirabayashi Sho, Wang Jianqiao, Kawagishi Hirokazu, Hirai Hirofumi
journal or publication title	Journal of Bioscience and Bioengineering
volume	120
number	1
page range	6-8
year	2015-07
出版者	Elsevier
権利	Copyright (C) 2014 Elsevier B.V. except certain content provided by third parties. ScienceDirect(R) is a registered trademark of Elsevier B.V.
注記	NOTICE: this is the author's version of a work that was accepted for publication in Journal of Bioscience and Bioengineering. Changes resulting from the publishing process, such as peer review, editing, corrections, structural formatting, and other quality control mechanisms may not be reflected in this document. Changes may have been made to this work since it was submitted for publication. A definitive version was subsequently published in Journal of Bioscience and Bioengineering, Volume 120, Issue 1, July 2015, 10.1016/j.jbiosc.2014.11.024
URL	http://hdl.handle.net/10297/8000

doi: 10.1016/j.jbiosc.2014.11.024

Note

**Improving xylitol production through recombinant expression of xylose reductase
in the white-rot fungus *Phanerochaete sordida* YK-624**

Sho Hirabayashi ^{1,†}, Jianqiao Wang ^{1,†}, Hirokazu Kawagishi ^{1,2,3}, Hirofumi Hirai ^{1,2*}

¹ Graduate School of Agriculture, Shizuoka University, 836 Ohya, Suruga-ku, Shizuoka
422-8529, Japan

² Research Institute of Green Science and Technology, Shizuoka University, 836 Ohya,
Suruga-ku, Shizuoka 422-8529, Japan

³ Graduate School of Science and Technology, Shizuoka University, 836 Ohya,
Suruga-ku, Shizuoka 422-8529, Japan

* Corresponding author. Department of Applied Biological Chemistry, Graduate School
of Agriculture, Shizuoka University, 836 Ohya, Suruga-ku, Shizuoka 422-8529, Japan.
Tel. & fax: +81 54 238 4853
E-mail address: ahhirai@ipc.shizuoka.ac.jp (H. Hirai).

[†] These authors contributed equally to this work.

Keywords: white-rot fungi, *Phanerochaete sordida* YK-624, xylose reductase,
recombinant expression, xylitol

Running title: Improving xylitol production by XR gene expression

Abstract

20 We generated an expression construct consisting of the xylose reductase (XR) gene
21 (*xr*) from *P. chrysosporium*. Transformant X7 exhibited increased xylitol production and
22 markedly higher XR activities than the wild-type strain. RT-PCR analysis demonstrated
23 that the increased XR activity was associated with constant expression of the
24 recombinant *xr* gene.

Lignocellulosic biomass is the most abundant organic compound and has therefore attracted worldwide interest as a feedstock for the production of bioethanol (1, 2). For the cost-effective production of value-added products from renewable lignocellulosic resources, microbial bioconversion processes must effectively utilize the pentose sugar xylose, as it is a major component of lignocellulose hydrolysates (3-5). However, the yeast *Saccharomyces cerevisiae*, which is typically used in ethanol production from hexoses, is unable to use pentose sugars such as a xylose (6, 7). Therefore, the development of a microbial-based system that can directly utilize pentose is necessary for the economic conversion of lignocellulose in biorefinery processes.

In yeasts and fungi, typically convert xylose to xylulose in a two-step reaction mediated by two kinds of oxidoreductases. An NADPH xylose reductase (XR) first reduces xylose to xylitol, which is then oxidized to xylulose by an NAD-linked xylitol dehydrogenase. And, improvement of xylitol production by recombinant microorganisms was received much attention (4, 8). Thus, the simultaneous production of ethanol and xylitol would significantly increase the efficiency of bioethanol production from lignocellulosic biomass.

The white-rot fungus *Phanerochaete sordida* YK-624, which was originally isolated from rotten wood, exhibits greater ligninolytic activity and selectivity than

either *P. chrysosporium* or *Trametes versicolor* (9). In our previous studies, we successfully developed a superior lignin-degrading strain of *P. sordida* YK-624 using a molecular breeding approach with a homologous expression system (10, 11). Here, we used a similar approach to generate high XR gene (*xr*)-expressing transformants of *P. sordida* YK-624 and investigated whether xylitol production was improved in these transformants.

Genomic DNA of *P. chrysosporium* ME-446 was extracted from mycelia using ISOPLANT II (Nippon Gene, Tokyo) and was then used as template for the PCR amplification of the full-length genomic *xr* gene (1144 bp) using the specific primers PcXRF1 and PcXRR1. The obtained PCR product was ligated into the cloning vector pMD20-T (Takara Bio, Shiga, Japan) and introduced into *Escherichia coli* DH5 α for sequencing. Primers XRF1 and XRR1 were designed to amplify the *xr* gene and introduce an *Xba* I site for cloning into the expression plasmid p*PsGPD-pro*, which was generated in our previous study (12). The amplified DNA fragment was digested with *Xba* I and cloned into *Xba* I-digested p*PsGPD-pro*, yielding plasmid p*PsGPD-XR*.

After p*PsGPD-XR* was sequenced to verify the absence of PCR errors, it was co-transformed with p*PsURA5* using the PEG method into UV-64 protoplasts that were prepared by a standard technique using cellulases (13). Co-transformed clones (X

strains) were identified by PCR with the primers *gpdF1* and *XRR2*, which were designed to amplify the recombinant *xr* gene. A total of 13 strains (X1-13) that were co-transformed with *pPsGPD-XR* and *pPsURA5* were obtained.

The effect of recombinant *xr* expression on xylitol production by the X strains was next investigated. *P. sordida* YK-624 (WT) and the X strains were incubated on PDA plates at 30 °C for 3 days, and 10-mm diameter disks were then punched out from the growing edge of the mycelia using a sterile cork borer. Two mycelial disks for each strain were placed into a 100-mL Erlenmeyer flask containing 10 mL nitrogen-limited Kirk medium (14) supplemented with 1.5% xylose as a carbon source, and the flask was then statically incubated at 30 °C for 7 days. Xylitol and xylose concentrations were determined using a high-performance liquid chromatograph equipped with a Shodex SH1821 column (8.0 mm × 300 mm, Showa Denko K.K., Tokyo, Japan) at 75 °C with 0.5 mM H₂SO₄ as the mobile phase at a flow rate of 0.6 mL min⁻¹. Xylitol and xylose were detected using an online refractive index detector. Transformant strain X7 exhibited the highest xylitol production among the 13 transformant strains (data not shown) and was therefore selected for subsequent analyses.

We next analyzed the transcription levels of *xr* by RT-PCR and measured the XR activity of strain X7 during 8 days of culture in nitrogen-limited Kirk medium. Mycelial

79 samples were collected and stored at -80 °C. Total RNA was isolated from the thawed
80 samples using an RNeasy Plant Mini kit (Qiagen, Valencia, CA). RT-PCR was
81 performed using 200 ng of total RNA with a PrimeScript RT-PCR Kit (Takara Bio) and
82 the gene-specific primer sets *gpdF1-XRR2* and *ActinF-ActinR*.

83 For the measurement of XR activity, each mycelial sample was added to 2 ml of
84 250 mM phosphate buffer (pH 7.0) and was then homogenized using a Polytron
85 PT1200E (Kinematica, Canada) at 4 °C. The homogenate was centrifuged (4 °C, 10,000
86 x g, 10 min) and the obtained supernatant was used as a cell-free extract for the
87 measurement of XR activity, which was determined by monitoring the oxidation of
88 NADPH to NADP⁺ ($\epsilon_{340} = 6.22 \text{ mM}^{-1} \text{ cm}^{-1}$) (15). The reaction mixture (1 ml) consisted
89 of 10 mM 2-mercaptoethanol, 50 mM xylose, and 0.17 mM NADPH in 250 mM
90 phosphate buffer (pH 7.0).

91 The time course of xylitol production by strains X7 and WT over 2, 4, 6, and 8
92 days of incubation in nitrogen-limited Kirk medium was monitored. After 4 days of
93 incubation, xylitol production by strain X7 reached 1.39 g L⁻¹, whereas the WT strain
94 had only produced 0.61 g L⁻¹ (Fig. 1). In contrast to the WT strain, the production of
95 xylitol by strain X7 continued to increase, reaching 3.61 g L⁻¹ of xylitol after 8 days of
96 incubation (Fig. 1). We next evaluated the consumption rates by the xylose

consumption, and the xylose consumption rate of the strain X7 (31.8%) was higher than that of WT strain (11.2%).

XR activity of strain X7 increased gradually for 8 days and was higher than that of WT throughout the incubation period (Fig. 2a). The highest XR activity for strain X7 was 9.14 nkat flask⁻¹ after 8 days of incubation, whereas that for WT was 6.54 nkat flask⁻¹ (Fig. 2a). The difference of fungal growth in WT strain and strain X7 was not observed under glucose- or xylose-containing medium. Consistent with these findings, RT-PCR indicated that strain X7 had constant *xr* expression between days 2 to 8 (Fig. 2b). Taken together, these results suggest that the increase of XR activity in strain X7 was due to constant expression of the recombinant *xr* gene, and that the recombinant expression of *xr* in the white-rot fungus *P. sordida* YK-624 improves xylitol production.

To conclude, the recombinant expression of *xr* is effective for the improvement of the xylitol production in white-rot fungi. Recently, direct ethanol production by the white-rot fungus *Phlebia* sp. MG-60 in pure culture without the addition of exogenous chemicals or enzymes was reported (16). In future studies, we intend to develop a molecularly bred strain of white-rot fungus that can simultaneously produce ethanol and xylitol from woody biomass with high yields.

This work was partially supported by Grants-in-Aid for Scientific Research (A) (Nos. 21248023 and 24248030) from the Ministry of Education, Culture, Sports, Science and Technology of Japan.

Reference

1. FitzPatrick, M., Champagne, P., Cunningham, M.F., and Whitney, R.A.: A biorefinery processing perspective: treatment of lignocellulosic materials for the production of value-added products, *Bioresour. Technol.*, **101**, 8915–8922 (2010).
2. Lin, Y. and Tanaka, S.: Ethanol fermentation from biomass resources: current state and prospects, *Appl. Microbiol. Biotechnol.*, **69**, 627–642 (2006).
3. Sasaki, M., Jojima, T., Inui, M., and Yukawa, H.: Xylitol production by recombinant *Corynebacterium glutamicum* under oxygen deprivation, *Appl. Microbiol. Biotechnol.*, **86**, 1057–1066 (2010).
4. Kötter, P., and Ciriacy, M.: Xylose fermentation by *Saccharomyces cerevisiae*, *Appl. Microbiol. Biotechnol.*, **38**, 776–783 (1993).
5. Eliasson, A., Christensson, C., Wahlbom, C.F., and Hahn-Hägerdal, B.: Anaerobic xylose fermentation by recombinant *Saccharomyces cerevisiae* carrying *XYL1*, *XYL2*, and *XKS1* in mineral medium chemostat cultures, *Appl. Environ. Microbiol.*, **66**, 3381–3386 (2000).

6. Vallet, C., Said, R., Rabiller, C., and Martin, M.L.: Natural abundance isotopic fractionation in the fermentation reaction: influence of the nature of the yeast, *Bioorg. Chem.*, **24**, 319–330 (1996).
7. Leticia, P., Miguel, C., Humberto, G., and Jaime, A.J.: Fermentation parameters influencing higher alcohol production in the tequila process, *Biotechnol. Lett.*, **19**, 45–47 (1997).
8. Marcus, D., Dignard, D., Lépine, G., Askew, C., Raymond, M., Whiteway, M., and Wu, C.: Comparative xylose metabolism among the Ascomycetes *C. albicans*, *S. stipitis* and *S. cerevisiae*, *PLoS One*, **8**, DOI: 10.1371/journal.pone.0080733 (2013).
9. Hirai, H., Kondo, R., and Sakai, K.: Screening of lignin degrading fungi and their ligninolytic enzyme activities during biological bleaching of kraft pulp. *Mokuzai Gakkaishi*, **40**, 980–986 (1994).
10. Sugiura, T., Mori, T., Kamei, I., Hirai, H., Kawagishi, H., and Kondo, R.: Improvement of ligninolytic properties in the hyper lignin-degrading fungus *Phanerochaete sordida* YK-624 using a novel gene promoter, *FEMS Microbiol. Lett.*, **331**, 81–88 (2012).
11. Sugiura, T., Yamagishi, K., Kimura, T., Nishida, T., Kawagishi, H., and Hirai, H.: Cloning and homologous expression of novel lignin peroxidase genes in the

white-rot fungus *Phanerochaete sordida* YK-624, Biosci. Biotechnol. Biochem., **73**,
1793–1798 (2009).

12. Suzuki, T., Dohra, H., Omae, S., Takeshima, Y., Choi, J.H., Hirai, H., and
Kawagishi, H.: Heterologous expression of a lectin from *Pleurocybella porrigens*
(PPL) in *Phanerochaete sordida* YK-624, J. Microbiol. Methods, **100**, 70–76
(2014).

13. Yamagishi, K., Kimura, T., Oita, S., Sugiura, T., and Hirai, H.: Transformation by
complementation of a uracil auxotroph of the hyper lignin-degrading basidiomycete
Phanerochaete sordida YK-624, Appl. Microbiol. Biotechnol., **76**, 1079–1091
(2007).

14. Tien, M., and Kirk, T.K.: Lignin peroxidase of *Phanerochaete chrysosporium*,
Methods Enzymol., **161**, 238–249 (1988).

15. Yokoyama, I., Suzuki, T., Kawai, K., Horitsu, H., and Takamizawa, K.: Purification,
characterization and structure analysis of NADPH-dependent d-xylose reductases
from *Candida tropicalis*, J. Ferm. Bioeng., **79**, 217–223 (1995).

16. Kamei, I., Hirota, Y., and Meguro, S.: Integrated delignification and simultaneous
saccharification and fermentation of hard wood by a white-rot fungus, *Phlebia* sp.
MG-60, Bioresour. Technol., **126**, 137–141 (2012).

Table 1 Oligonucleotides used as primers in this study.

Primer name	Nucleotide sequence (5'-sequence-3')
PcXRF1	ATGCTTTCTTCGCCAACCTC
PcXRR1	TCAGCATTACGGAGAAAGTACG
XRF1	AATCTAGAATGCTTTCTTCGCCAACCTC
XRR1	AATCTAGATCAGCATTACGGAGAAAGTACG
gpdF1	AAGCAGCGAGGATTGTACC
XRR2	GATGGAAGGTGTTCCACAG
ActinF	AGCACGGTATCGTCACCAAC
ActinR	AGCGAAACCCTCGTAGATGG

Fig. 1 Time course of xylitol production by the wild-type (WT) strain (closed diamonds) and strain X7 (closed squares). Values are the means $\pm$ SD of triplicate samples. Asterisks indicate values that were determined by the Student's t-test to be significantly different from WT (* $P < 0.05$, ** $P < 0.01$).

Fig. 2 XR activity and expression of the *xr* gene by strain X7. a) Time course of XR activities in the wild-type strain (closed diamonds) and strain X7 (closed squares). Values are the means $\pm$ SD of triplicate samples. Asterisks indicate values that were determined by the Student's t-test to be significantly different from WT (* $P < 0.05$, ** $P < 0.01$). b) RT-PCR analysis of *xr* gene transcription in strain X7. The expression of *actin* was used as an internal control.