Improvement of ligninolytic properties by recombinant expression of glyoxal oxidase gene in hyper lignin-degrading fungus *Phanerochaete sordida* YK-624

著者	Yamada Yuto, Wang Jianqiao, Kawagishi Hirokazu, Hirai Hirofumi
journal or publication title	Bioscience, Biotechnology, and Biochemistry
volume	78
number	12
page range	2128-2133
year	2014-08-13
出版者	Taylor & Francis
権利	This is an electronic version of an article published in Bioscience, Biotechnology, and Biochemistry Volume 78, Issue 12, pages 2128-2133, 2014. Bioscience, Biotechnology, and Biochemistry is available online at: www.tandfonline.com/Article DOI; 10.1080/09168451.2014.946398
URL	http://hdl.handle.net/10297/9090

doi: 10.1080/09168451.2014.946398

Regular paper

**Improvement of ligninolytic properties by recombinant expression of glyoxal
oxidase gene in hyper lignin-degrading fungus *Phanerochaete sordida* YK-624**

Yuto Yamada ^{1, †}, Jianqiao Wang ^{2, †}, Hirokazu Kawagishi ^{1, 2, 3}, Hirofumi Hirai ^{1, 3 *}

¹ Graduate School of Agriculture, Shizuoka University, 836 Ohya, Suruga-ku, Shizuoka
422-8529, Japan

² Graduate School of Science and Technology, Shizuoka University, 836 Ohya,
Suruga-ku, Shizuoka 422-8529, Japan

³ Research Institute of Green Science and Technology, Shizuoka University, 836 Ohya,
Suruga-ku, Shizuoka 422-8529, Japan

Received June 5, 2014; Accepted July 2, 2014

* Corresponding author. Department of Applied Biological Chemistry, Graduate School
of Agriculture, Shizuoka University, 836 Ohya, Suruga-ku, Shizuoka 422-8529, Japan.

Tel. & fax: +81 54 238 4853

E-mail address: ahhirai@ipc.shizuoka.ac.jp (H. Hirai).

[†] These authors contributed equally to this work.

Keywords: *Phanerochaete sordida* YK-624, lignin degradation, glyoxal oxidase,
molecular breeding

Running title: Ligninolytic improvement by expression of GLOX gene

Abstract

Glyoxal oxidase (GLOX) is a source of the extracellular H₂O₂ required for the oxidation reactions catalyzed by the ligninolytic peroxidases. In the present study, the GLOX-encoding gene (*glx*) of *Phanerochaete chrysosporium* was cloned, and *bee2* promoter of *P. sordida* YK-624 was used to drive the expression of *glx*. The expression plasmid was transformed into a *P. sordida* YK-624 uracil auxotrophic mutant (strain UV-64), and 16 clones were obtained as GLOX-introducing transformants. These transformants showed higher GLOX activities than wild-type *P. sordida* YK-624 and control transformants harboring marker plasmid. RT-PCR analysis indicated that the increased GLOX activity was associated with elevated recombinant *glx* expression. Moreover, these transformants showed higher ligninolytic activity than control transformants. These results suggest that the ligninolytic properties of white-rot fungi can be improved by recombinant expression of *glx*.

Introduction

Lignin is one of the most abundant and recalcitrant natural polymers, consisting of various nonphenolic phenylpropanoid units complexed into a three-dimensional network. Therefore, lignin degradation is a key step for carbon recycling in forest ecosystems, and the process is a central issue in lignocellulose-based industries (*e.g.*, in paper pulp manufacture and production of bioethanol and other renewable chemicals).¹⁾ In nature, white-rot fungi (members of the basidiomycete group) are able to degrade recalcitrant lignin efficiently ²⁾. The biodegradation process is initiated via a one-electron oxidation of lignin units, a step that is mediated by extracellular heme peroxidases (including lignin peroxidase (LiP), manganese peroxidase (MnP), and versatile peroxidase), enzymes that are of keen interest as industrial biocatalysts.^{3, 4)} Activation of molecular oxygen to hydrogen peroxide by extracellular oxidases represents a common step in the fungal decay strategy, as shown by biochemical ^{2, 5)} and genomic ^{6, 7)} evidence. White-rot decay is based on peroxide activation of high-redox-potential fungal heme peroxidases that depolymerize lignin, leaving a cellulose-rich residue.⁸⁾ The mechanism of enzymatic attack on lignin by these fungi has been extensively investigated because of its biotechnological interest.^{1, 9)}

The white-rot fungus *Phanerochaete chrysosporium* secretes three known classes of extracellular enzymes under ligninolytic (secondary metabolic) conditions in defined glucose medium: glyoxal oxidases (GLOXs), LiPs, and MnPs.³⁾ GLOX represented a family of radical copper oxidases produced by filamentous fungi, mainly organisms belonging to the group of white-rot fungi; the GLOX from *P. chrysosporium* is the most extensively studied of this enzyme family. GLOX acts as a source of the extracellular hydrogen peroxide that is required for the oxidations catalyzed by the LiPs and MnPs involved in lignin degradation.^{10, 11)} The enzyme catalyzes the oxidation of aldehydes to the corresponding carboxylic acids by coupling the reaction to the reduction of dioxygen to H₂O₂. Although GLOX activity was detected under ligninolytic conditions,^{10, 11)} there is no evidence for GLOX involved in lignin degradation.

The white-rot fungus *Phanerochaete sordida* YK-624, which was isolated from rotted wood, exhibits greater ligninolytic activity and selectivity than either *P. chrysosporium* or *Trametes versicolor*.¹²⁾ In our previous study, we demonstrated that *P. sordida* YK-624 produces MnP^{12, 13)} and LiP^{14, 15)} as ligninolytic enzymes. Moreover, we previously identified a protein (BUNA2) that accumulated to high levels in cultures of *P. sordida* YK-624 under wood-rotting conditions. The promoter region of the BUNA2-encoding gene, designated *bee2*, was successfully cloned and demonstrated to

be a useful regulator for the high expression of genes under ligninolytic conditions.¹⁶⁾

More recently, we have reported that the production of MnP was increased by the recombinant expression of a 5-aminolevulinic acid synthase-encoding gene.¹⁷⁾

In the present study, we report the transformation of *P. sordida* YK-624 with a recombinant *glx* gene under regulation of the *bee2* promoter, yielding elevated transcript accumulation and GLOX activity under wood-rotting conditions. Moreover, we demonstrate that these transformants exhibit improved ligninolytic properties.

Materials and Methods

Strains

P. sordida strain YK-624 (ATCC 90872), uracil auxotrophic *P. sordida* strain UV-64,¹⁸⁾ and *P. chrysosporium* ME-446 (ATCC 34541) were used in this study. All strains and transformants were maintained on potato dextrose agar (PDA) slants at 4 °C.

Construction of a GLOX-encoding gene expression vector, co-transformation of UV-64, and screening of regenerated clones

Genomic DNA was isolated from *P. chrysosporium* ME-446 mycelium using

ISOPLANT II (Nippon Gene, Tokyo). Gene sequences for the GLOX-encoding gene were obtained from the DOE Joint Genome Institute and used to design PCR primers. The full-length *glx* (1907 bp) gene was amplified using the primer pair PcGLOX F1-PcGLOX R1. The procedure for constructing the *glx* expression plasmid, pBUNA2pro-GLOX, is described in Fig. 1. The plasmid pBUNA2pro, which was generated in our previous study,¹⁶⁾ was used for the construction of the *glx* expression plasmid. Primers GLOX F1 and Pc GLOX R1 were designed to amplify the *glx* gene and to introduce a *Bgl* II site just before the first methionine codon and to add a *Bgl* II site just after the stop codon, respectively (step 1). The amplified DNA fragment was digested with *Bgl* II (New England Biolabs) and cloned into *Bgl* II digested pBUNA2pro (step 2), yielding plasmid pBUNA2pro-GLOX (step 3). pBUNA2pro-GLOX was sequenced to verify the absence of PCR errors.

Fig. 1

UV-64 protoplasts were prepared by a standard technique using cellulases and then co-transformed with p*PsURA5* and pBUNA2pro-GLOX using the PEG method.¹⁸⁾ The co-transformed clones (G strains) were screened by PCR, as described previously,¹⁹⁾ with the following modifications: primers PcGLOX F2 and PsGPD R1 were designed to amplify the *glx* gene fused with the *bee2* promoter.

Enzyme assay

P. sordida YK-624, G strains, and U strains (UV-64 protoplasts were singly-transformed with p*PsURA5*, yielding transformants designated as U strains) were incubated on PDA plates at 30 °C for 3 days, and 10-mm diameter disks then were punched out from the growing edge of the mycelia using a sterile cork borer. For determining the activities in the extracellular culture fluids, two mycelial disks from each source plate were placed into a 100-mL Erlenmeyer flask containing 10 mL nitrogen-limited Kirk medium,²⁰⁾ which then was statically incubated at 30 °C for 2, 4, 6, and 8 days. Meanwhile, for detecting the activities in fungal-treated beech wood meals, these strains were cultured in 50-mL Erlenmeyer flasks containing 0.5 g of extractive-free beech wood meal (80-100 mesh) and 1.25 ml of distilled water, and incubated at 30 °C for 5, 10, 15, and 20 days. Following the culture period, the method described by Hirai *et al.* (1994) was modified for enzyme extraction. Briefly, fungal-treated wood meal was homogenized with 25 mL of 50 mM malonate (pH 4.5) or phosphate (pH 6.0) buffer containing 0.05% Tween 20 (Wako) using a Polytron PT1200E (Kinematica, Canada) homogenizer for a total of 5 min (20 s blending with 10-min intervals) at 4 °C.

MnP activity was measured by monitoring the oxidation of 2,6-dimethoxyphenol

to coerulignone (extinction coefficient at 470 nm (ϵ_{470}) = 49.6 mM⁻¹ cm⁻¹).²¹⁾ The reaction mixture (1 mL) contained 2,6-dimethoxyphenol (1 mM), MnSO₄ (1 mM), and H₂O₂ (0.2 mM) in 50 mM malonate buffer (pH 4.5). GLOX activity was measured by monitoring the oxidation of guaiacol to the tetramer (ϵ_{470} = 26.6 mM⁻¹ cm⁻¹). The reaction mixture (1 mL) contained guaiacol (1 mM), 0.08 nkat of horseradish peroxidase, and 2-methylglyoxal (1 mM) in 50 mM phosphate buffer (pH 6.0). A standard curve was constructed for horseradish peroxidase (Y axis: production of the tetramer; X axis: H₂O₂ concentration) and used to calculate the production of H₂O₂ in the reaction system. One katal (kat) was defined as the amount of enzyme producing 1 mol of product per second.

Determination of ligninolytic properties

G strains, U strains, and *P. sordida* YK-624 were cultured in 50-mL Erlenmeyer flasks containing 0.5 g of extractive-free beech wood meal (80-100 mesh) and 1.25 ml of distilled water, which then were incubated at 30 °C for 28 days. After incubation, weight loss, Klason lignin content, and acid-soluble lignin content of the fungal-treated wood meal were determined, as previously described.¹²⁾ The selection factor (SF), which is an indicator of ligninolytic selectivity, was calculated as follows: SF = lignin

loss/holocellulose loss. Holocellulose loss was calculated as follows: total weight loss – lignin loss.

Transcriptional analysis of strain G-22

Strain G-22 was inoculated in liquid medium as described above and then statically incubated at 30 °C for 2, 4, 6, and 8 days. The mycelia were then collected and stored at -80 °C pending transcriptional analysis. One hundred milligrams of mycelial mat was ground to a fine powder using a mortar and pestle under liquid nitrogen, and then total RNA was isolated from the powder using an RNeasy Plant Mini kit (Invitrogen). RT-PCR was performed using 200 ng of total RNA with a PrimeScript RT-PCR Kit (TaKaRa Bio, Otsu) and the gene-specific primer sets PcGLOX F3-PsGPD R1 (to amplify recombinant *glx*) and ActinF-ActinR (to amplify the actin-encoding gene, *act*). PCR was performed for 28 cycles of template denaturation at 95 °C for 30 s, primer annealing at 56 °C for 1 min, and DNA extension at 72 °C for 1 min using Ex Taq DNA Polymerase (TaKaRa Bio).

Nucleotide sequence accession number

The nucleotide sequences of *glx* derived from *P. chrysosporium* have been

deposited in the DDBJ database (<http://www.ddbj.nig.ac.jp/>) under accession no.

AB910536.

Results and Discussion

Isolation of recombinant *glx*-expressing transformants

P. chrysosporium was cultured in Kirk liquid medium for 5 days, and then genomic DNA was extracted from the mycelia. Based on the genome database of *P. chrysosporium*, two primers were designed to amplify the genomic DNA encoding GLOX. Genomic DNA of *glx* obtained by genome PCR was inserted between the *bee2* promoter and *gpd* terminator by restriction enzyme digestion and ligation (Fig. 1). This expression plasmid, pBUNA2pro-GLOX, was introduced into strain UV-64 together with p*PsURA5* as a marker plasmid. The presence of the *bee2* promoter-*glx* fusion gene in each uracil prototrophic clone was confirmed by PCR using genomic DNA as the template (data not shown). We obtained 16 of these “G” strains that were co-transformed with pBUNA2pro-GLOX and p*PsURA5*.

GLOX production and ligninolytic properties of recombinant *glx*-expressing

transformants

In our previous reports, *P. sordida* YK-624 produces only LiP and MnP as ligninolytic enzymes,¹²⁻¹⁵⁾ and LiP activity has been hardly detected in the culture containing Mn ion.¹²⁾ Therefore, we focused MnP among ligninolytic enzymes, and the production of GLOX and MnP by G strains was compared with those by singly-transformed (p*PsURA5* alone) “U” strains. The mean GLOX activity in the culture medium inoculated with G strains for 5 days was 0.66 nkat/flask, whereas that in the culture medium of U strains was 0.19 nkat/flask (Fig. 2a). The mean MnP activity in the culture medium inoculated with G strains was 8.51 nkat/flask, whereas that in the culture media of U strains was 7.78 nkat/flask (Fig. 2b). These results indicate that G strains showed higher GLOX activities than U strains, although MnP activities of G strains did not apparently differ from those of U strains.

Fig. 2

The effect of recombinant *glx* expression on the ligninolytic properties of G strains was investigated. G strains showed 1.08-fold higher ligninolytic activity (Fig. 3a) and 1.18-fold higher SF values than U strains (Fig. 3b). In particular, strain G-22 showed 1.22-fold higher ligninolytic activity and 1.25-fold higher SF values than U strains (data not shown). These results indicated that the ligninolytic properties were improved by the recombinant expression of *glx* using the *bee2* promoter, with strain G-22 exhibiting

Fig. 3

the highest lignin degradation activity among the 16 screened co-transformants. In our previous study, the promoter region of *bee2* was demonstrated to be a useful regulator for the high expression of genes under ligninolytic conditions.¹⁶⁾ The results of the present study are consistent with that previous work. The proposed role of GLOX in lignin biodegradation is to generate H₂O₂ for LiP- and MnP-mediated reactions.²²⁾ Moreover, the identification of *glx* transcripts in decayed wood is consistent with a close physiological connection between extracellular peroxidases and GLOX.²³⁾ These reports, together with our present results, suggest that GLOX plays an important role in lignin degradation by white-rot fungi.

To confirm whether the improvement of the ligninolytic properties resulted from an increase in GLOX production, GLOX and MnP activities produced in beech wood meals inoculated with G-22 and wild type (WT) were determined. The MnP activity of G-22 was similar to that of WT, and no drastic fluctuations were observed (Fig. 4b). In contrast, higher GLOX activities were detected at 10 and 15 days in G-22, although similar GLOX activities for each strain were detected at 5 days (Fig. 4a). These results indicate that the improvement of the ligninolytic properties correlates with the higher production of GLOX.

Fig. 4

GLOX production and transcriptional analysis of G-22

Transcriptional analysis was performed by RT-PCR to confirm whether the increment of GLOX activity reflected increased accumulation of *glx* transcript. Specifically, we analyzed a time course of GLOX activity in G-22 and WT. GLOX activity in G-22 increased gradually after 4 days, remaining relatively stable through 8 days (Fig. 5a). GLOX activity in G-22 was higher than that in WT during the incubation. The highest GLOX activity in G-22 was observed at 8 days, when activity was 7.2-fold higher than that in WT.

Fig. 5

We next analyzed the transcription of recombinant *glx* in G-22 by RT-PCR (Fig. 5b). The analysis indicated that high levels of recombinant *glx* transcript were detected after 4 to 8 days. These results suggest that the increase of GLOX activity in G-22 was caused by *bee2*-driven expression of the recombinant *glx*.

To conclude, we found that the recombinant expression of *glx* under the control of the *bee2* promoter is effective for the improvement of the ligninolytic properties in white-rot fungi. We also provide the evidence that physiologic productivity of GLOX in white-rot fungi is not enough under ligninolytic condition. In our next study, we propose to investigate the molecular breeding of superior lignin-degrading fungi by the simultaneous expression of *glx* and a MnP-encoding gene (*mnp4*). To our knowledge,

this is the first report to demonstrate that GLOX is involved in lignin degradation.

Acknowledgements

This work was partially supported by Grants-in-Aid for Scientific Research (A) (Nos. 21248023 and 24248030) from the Ministry of Education, Culture, Sports, Science and Technology of Japan.

Reference

1. Martínez AT, Ruiz-Dueñas FJ, Martínez MJ, del Río JC, Gutiérrez A. Enzymatic delignification of plant cell wall: from nature to mill. *Curr. Opin. Biotechnol.* 2009;20:348–357.
2. Martínez AT, Speranza M, Ruiz-Dueñas FJ, Ferreira P, Camarero S, Guillén F, Martínez MJ, Gutiérrez A, del Río JC. Biodegradation of lignocellulosics: microbial, chemical, and enzymatic aspects of the fungal attack of lignin. *Int. Microbiol.* 2005;8:195–204.
3. Kirk TK, Farrell R. Enzymatic "combustion": the microbial degradation of lignin. *Annu. Rev. Microbiol.* 1987;41:465–505.

- 231 4. Ruiz-Dueñas FJ, Martínez AT. Microbial degradation of lignin: how a bulky
232 recalcitrant polymer is efficiently recycled in nature and how we can take
233 advantage of this. *Microb. Biotechnol.* 2009;2:164–177.
- 234 5. Baldrian P, Valášková V. Degradation of cellulose by basidiomycetous fungi.
235 *FEMS Microbiol. Rev.* 2008;32:501–521.
- 236 6. Martinez D, Challacombe J, Morgenstern I *et al.* Genome, transcriptome, and
237 secretome analysis of wood decay fungus *Postia placenta* supports unique
238 mechanisms of lignocellulose conversion. *Proc. Natl. Acad. Sci. USA.*
239 2009;106:1954–1959.
- 240 7. Martinez D, Larrondo LF, Putnam N *et al.* Genome sequence of the lignocellulose
241 degrading fungus *Phanerochaete chrysosporium* strain RP78. *Nat. Biotechnol.*
242 2004;22:695–700.
- 243 8. Martínez AT, Rencoret J, Nieto L, Jiménez-Barbero J, Gutiérrez A, del Río JC.
244 Selective lignin and polysaccharide removal in natural fungal decay of wood as
245 evidenced by in situ structural analyses. *Environ. Microbiol.* 2011;13:96–107.
- 246 9. Hammel KE, Cullen D. Role of fungal peroxidases in biological ligninolysis. *Curr.*
247 *Opin. Plant Biol.* 2008;11:349–355.
- 248 10. Kersten PJ. Glyoxal oxidase of *Phanerochaete chrysosporium*: its characterization

249 and activation by lignin peroxidase. Proc. Natl. Acad. Sci. USA.
250 1990;87:2936–2940.

251 11. Kersten PJ, Kirk TK. Involvement of a new enzyme, glyoxal oxidase, in
252 extracellular hydrogen peroxide production by *Phanerochaete chrysosporium*. J.
253 Bacteriol. 1987;169:2195–201.

254 12. Hirai H, Kondo R, Sakai K. Screening of lignin degrading fungi and their
255 ligninolytic enzyme activities during biological bleaching of kraft pulp. Mokuzai
256 Gakkaishi. 1994;40:980–986.

257 13. Hirai H, Kondo R, Sakai K. Effect of metal ions on biological bleaching of kraft
258 pulp with *Phanerochaete sordid* YK-624. Mokuzai Gakkaishi. 1995;41:69–75.

259 14. Sugiura M, Hirai H, Nishida T. Purification and characterization of a novel lignin
260 peroxidase from white-rot fungus *Phanerochaete sordida* YK-624. FEMS
261 Microbiol. Lett. 2003;224:285–290.

262 15. Hirai H, Sugiura M, Kawai S, Nishida T. Characteristics of novel lignin
263 peroxidases produced by white-rot fungus *Phanerochaete sordida* YK-624. FEMS
264 Microbiol. Lett. 2005;246:19–24.

265 16. Sugiura T, Mori T, Kamei I, Hirai H, Kawagishi H, Kondo R. Improvement of
266 ligninolytic properties in the hyper lignin-degrading fungus *Phanerochaete sordida*

YK-624 using a novel gene promoter. FEMS Microbiol. Lett. 2012;331:81–88.

17. Hirai H, Misumi K, Suzuki T, Kawagishi H. Improvement of manganese peroxidase production by the hyper lignin-degrading fungus *Phanerochaete sordida* YK-624 by recombinant expression of the 5-aminolevulinic acid synthase gene. Curr. Microbiol. 2013;67:708–711.

18. Yamagishi K, Kimura T, Oita S, Sugiura T, Hirai H, Transformation by complementation of a uracil auxotroph of the hyper lignin-degrading basidiomycete *Phanerochaete sordida* YK-624. Appl. Microbiol. Biotechnol. 2007;76:1079–1091.

19. Sugiura T, Yamagishi K, Kimura T, Nishida T, Kawagishi H, Hirai H. Cloning and homologous expression of novel lignin peroxidase genes in the white-rot fungus *Phanerochaete sordida* YK-624. Biosci. Biotechnol. Biochem. 2009;73:1793–1798.

20. Tien M, Kirk TK. Lignin peroxidase of *Phanerochaete chrysosporium*. Methods Enzymol. 1988;161:238–249.

21. Périé FH, Gold MH. Manganese regulation of manganese peroxidase expression and lignin degradation by the white rot fungus *Dichomitus squalens*. Appl. Environ. Microbiol. 1988;57:2240–2245.

- 285 22. Vanden Wymelenberg A, Sabat G, Mozuch M, Kersten PJ, Cullen D, Blanchette
286 RA. Structure, organization, and transcriptional regulation of a family of copper
287 radical oxidase genes in the lignin-degrading basidiomycete *Phanerochaete*
288 *chrysosporium*. Appl. Environ. Microbiol. 2006;72:4871–4877.
- 289 23. Janse BJH, Gaskell J, Akhtar M, Cullen D. Expression of *Phanerochaete*
290 *chrysosporium* genes encoding lignin peroxidases, manganese peroxidases, and
291 glyoxal oxidase in wood. Appl. Environ. Microbiol. 1998;64:3536–3538.

Fig. 1 Plasmid map of the *glx* expression plasmid pBUNA2pro-GLOX.

Fig. 2 GLOX (a) and MnP (b) activities in the extracellular culture fluids inoculated with U strains and G strains for 5 days.

Fig. 3 Ligninolytic properties of U strains and G strains. Lignin degradation (a) and selection factor (b) by each strain for 28 days.

Fig. 4 GLOX (a) and MnP (b) activities detected in fungal-treated beech wood meals. G-22 (triangles) and WT (squares).

Fig. 5 Production of GLOX and expression of *glx* by strain G-22. (a) Time course of GLOX activity in strain G-22 (triangles) and WT (squares). (b) RT-PCR analysis of *glx* in strain G-22. The expression of actin was used as an internal control.

Fig. 1

Fig. 2

Fig. 3

Fig. 4

Fig. 5