

Volatiles from wasabi inhibit entomopathogenic fungi: implications for tritrophic interactions and biological control

メタデータ	言語: eng 出版者: 公開日: 2016-04-19 キーワード (Ja): キーワード (En): 作成者: Atsumi, Ayako, Saito, Tsutomu メールアドレス: 所属:
URL	http://hdl.handle.net/10297/9349

Title

Volatiles from wasabi inhibit entomopathogenic fungi: Implications for tritrophic interactions and biological control

Authors

Ayako Atsumi and Tsutomu Saito*
Faculty of Agriculture, Shizuoka University, 836 Ohya, Suruga, Shizuoka 422-8529,
Japan

E-mail addresses

Ayako Atsumi: japanesegirl-no-185@ezweb.ne.jp
Tsutomu Saito: atsaito@ipc.shizuoka.ac.jp

***Corresponding author**

Tsutomu Saito: Faculty of Agriculture, Shizuoka University, 836 Ohya, Suruga,
Shizuoka 422-8529, Japan
E-mail address: atsaito@ipc.shizuoka.ac.jp
Tel/Fax: +81 54 238 4790

ABSTRACT

When wasabi plants are damaged by plant pathogens or herbivores they produce volatile allyl isothiocyanate (AITC) as a defence mechanism. In the current study we conducted experiments to determine whether volatiles from damaged wasabi leaves or synthetic AITC also have negative effects on the beneficial entomopathogenic fungi, *Beauveria bassiana sensu lato* (Balsamo) Vuillemin and *Isaria fumosorosea* Wize, which contribute to regulation of a major pest of wasabi, the striated white butterfly, *Pieris melete* Ménétrières. Conidial germination of both fungi was inhibited when exposed to volatiles from macerated wasabi leaves; the concentration of AITC in the volatiles was also quantified. The inhibitory activity of macerated wasabi leaves was compared with that of synthetic AITC, demonstrating that the inhibitory effect of wasabi volatiles was due to the presence of AITC. These results indicate that AITC released from damaged wasabi leaves has the potential to inhibit fungal infection of insect pests in wasabi fields.

Keywords: wasabi, plant volatiles, allyl isothiocyanate, *Beauveria bassiana*, *Isaria fumosorosea*, *Pieris melete*

1. Introduction

Japanese horseradish, or wasabi (*Wasabia japonica* Matsumura), is a cruciferous plant widely used in Japanese cuisine, particularly as a pungent spice in sushi. Wasabi plants are grown for 2 to 3 years in rice paddy-like fields, filled throughout the year with shallow flowing water from mountain streams (Figure 1a). These conditions should be very suitable for biological control of herbivores using entomopathogenic fungi because high humidity is an absolute requirement for conidial germination and the establishment of infection (Hall 1981; Helyer et al. 1992; Luz & Fargues 1999; Milner & Lutton 1986). Furthermore, to avoid downstream water pollution, there are enforced restrictions on chemical applications in wasabi fields; this should also be beneficial for entomopathogenic fungi that can be susceptible to pesticides, particularly fungicides (Saito & Yabuta 1996; Shinohara et al. 2013).

The strong flavour of wasabi is due to the presence of allyl isothiocyanate (AITC), also called mustard oil, which is produced in plant tissues that have been damaged by biotic or abiotic stressors; the enzyme myrosinase hydrolyses allyl glucosinolate (sinigrin) producing the breakdown product, AITC (Rask et al. 2000; Wittstock & Halkier 2002; Wittstock et al. 2003). The compound is highly toxic to a wide range of organisms including various herbivores (Agrawal & Kurashige 2003; Eltayeb et al. 2010; Rask et al. 2000; Wittstock et al. 2003) and plant pathogens (Mayton et al. 1996; Oliver et al. 1999). As such, AITC contributes to the plant's defence strategy against herbivores and disease (Chew 1988; Louda & Mole 1991; Rask et al. 2000; Wittstock et al. 2003). Wasabi actually produces larger quantities of AITC in leaves, petioles, rhizomes and roots, than other cruciferous plants (Kumagai et al. 1994; Sultana et al. 2003).

As AITC is toxic to plant pathogens (including fungi) it is possible that it could also have toxic effects on beneficial entomopathogenic fungi in the field; this may decrease their potential as biological control agents of herbivorous pests on wasabi. Therefore, to ensure successful biological control, fungal isolates with potential for augmentation in the field, should be evaluated for their susceptibility to volatile AITC prior to use. To date, there have already been some studies describing interactions between other plant volatiles and entomopathogenic fungi (Hountondji et al. 2005; Baverstock et al. 2005; Brown et al. 1995); however little is known about the effects of volatile AITC released from cruciferous plants on entomopathogenic fungi, with the exception of one study describing effects of synthetic isothiocyanates, including AITC, produced by cruciferous plants, on conidial germination and mycelial growth of *Metarhizium anisopliae* (Metschn.) Sorokin when incorporated into the culture medium (Inyang et al. 1999).

The purpose of our study was to examine the effects of volatiles released from wasabi leaves on conidial germination of entomopathogenic fungi that had been isolated from populations of the small white butterfly, *Pieris melete* Ménétrières, a serious pest of wasabi plants in Japan (Nakata 1963; Saito et al. 2010). Based on these results, we discuss the importance of AITC in terms of biological control and plant volatile-mediated tritrophic interactions between plants, herbivores and their natural enemies.

2. Materials and methods

2.1. Collection of entomopathogenic fungi

In 2011 and 2012, *P. melete* larvae were collected from several wasabi fields in

Shizuoka, Japan, and then reared in groups of 20–30 on leaves of kale, *Brassica oleracea* L. var. *acephala*, in containers (20 × 15 × 5 cm) at 25 ± 1°C and 16L: 8D. Dead larvae and pupae were transferred to Petri dishes (9 cm diameter) with a wet filter paper and incubated for a week at 25 ± 1°C in darkness to induce fungal sporulation on cadavers. Entomopathogenic fungi were isolated from infected cadavers on to Sabouraud dextrose agar (SDA) (Difco BD Bioscience, MD, USA) and all isolates were stored on slopes of culture medium in glass test tubes at –60°C prior to use in each experiment. Each isolate was identified according to morphological characteristics using the taxonomic keys for the genus *Beauveria* (Rehner et al. 2011), genus *Isaria* (Liang et al. 2005; Luangsa-ard et al. 2005; Samson 1974; Shimazu 2001) and genus *Lecanicillium* (Zare & Gams 2001, 2008).

2.2. Effect of volatiles from macerated wasabi leaves on conidial germination

For use in experiments, two isolates were selected from those collected in the field (see section 3.1); these isolates were retrieved from storage, subcultured on SDA in Petri dishes (9 cm diameter) and incubated at 25 ± 1°C for 10 d in darkness. Conidial suspensions of each isolate were prepared by scraping the mycelium from plate cultures into sterile 0.1% Tween 80, agitating with a vortex mixer and then filtering through sterile muslin (0.2 mm mesh size). Conidial concentrations of the suspensions were determined using a Thoma haemocytometer and adjusted to 1×10^7 conidia mL⁻¹ by the addition of sterile 0.1% Tween 80. Fifty µL aliquots of suspension were placed, individually, on to 10 mL SDA in Petri dishes (9 cm diameter); the dishes, without their lids, were each suspended upside-down beneath the lids of airtight glass chambers (12 cm diameter, 7 cm height, 0.79 L volume) using double-sided adhesive tape. The

prepared chambers were used to expose conidia to volatiles from macerated wasabi leaves as described below.

Fresh wasabi leaves collected from the field were macerated for about 5 min in a food processor (SKH-A100, Tiger, Osaka, Japan) and then immediately 0, 1, 3, 5 or 10 g were placed in the base of each chamber; these quantities were calculated as 0, 1.3, 3.8, 6.3 and 13 g L⁻¹ chamber, respectively. The chambers were sealed using petrolatum and then incubated at 25 ± 1°C and 16L:8D. After 24 h, the conidia were fixed and stained with drops of lactophenol-cotton blue solution (Sigma-Aldrich, MO, USA); the germination rate of approximately 100 conidia in each replicate of each treatment was determined under a microscope, for both isolates. Conidia were considered to have germinated when the germ tube was at least half as long as the width of the conidium. Five replicate chambers were used for each treatment. The proportion of conidia that had germinated in each replicate was arcsine square-root transformed and submitted to factorial analysis for two factors, the quantity of wasabi and the fungal isolate, using a two-way ANOVA. If there was an interaction between the two factors, multiple comparisons using a Bonferroni correction were made. These analyses were all done in the software package SPSS (2009).

2.3. Effect of synthetic AITC on conidial germination

The toxicity of synthetic AITC (99%, Wako Pure Chem., Osaka, Japan) was examined using the same method as described above in section 2.2, except that 100 µL of acetone solution containing AITC was used instead of macerated wasabi leaves. A series of different concentrations of AITC that had been selected from preliminary experiments (data not shown) were evaluated: 0, 6.3, 63 and 630 µg L⁻¹ chamber. Five replicate

chambers were used for each treatment. Data were analyzed using the same methods as described above in section 2.2.

2.4. Quantifying the concentration of AITC in volatiles from macerated wasabi leaves

AITC in volatiles from macerated wasabi leaves was quantified using a headspace sampling method in conjunction with Solid Phase Microextraction (SPME). Three g of macerated wasabi leaves (3.8 g L^{-1} chamber) were placed in the bases of each of five of the glass chambers described in section 2.2. An SPME fibre (DVB/CAR/PDMS, Sigma-Aldrich, Dorset, UK) was inserted through a 4 mm diameter hole (sealed with a rubber washer) into the side of each chamber for 30 min at $25 \pm 1^\circ\text{C}$. Each SPME fibre was then introduced into the injector of a gas chromatograph (3800GC, VARIAN, CA, USA) with a DB-WAX capillary column ($60 \text{ m} \times 0.32 \text{ mm}$, $0.25 \mu\text{m}$). Helium was used as the carrier gas (2 mL min^{-1}). Thermal conditions for the column were 35°C for 5 min, followed by a temperature increase from $35\text{--}120^\circ\text{C}$ at a rate of 5°C min^{-1} and then from $120\text{--}320^\circ\text{C}$ at a rate of $15^\circ\text{C min}^{-1}$.

The concentration of AITC produced was quantified based on the area beneath the peak, which was estimated using a standard curve for AITC. Data for the standard curve were collected using $100 \mu\text{L}$ acetone solutions containing known concentrations of AITC (0, 6.3, 63, 630 and $6300 \mu\text{g L}^{-1}$ chamber); there were three replicate chambers for each concentration and these data were subjected to linear regression to determine the relationships between peak areas and log doses using the software package SPSS (2009).

3. Results

3.1. Entomopathogenic fungi isolated from *P. melete*

A total of 1557 larvae of *P. melete* were collected from wasabi fields (Figure 1a) and five species of fungi were isolated from 67 cadavers (mainly pupae): *Beauveria bassiana sensu lato* (Balsamo) Vuillemin (six isolates) (Figure 1b); *Isaria fumosorosea* Wize (three isolates) (Figure 1c); *Isaria tenuipes* Peck (52 isolates) (Figure 1d); *Isaria cateniannulata* (Z. Q. Liang) Samson & Hywel-Jones (eight isolates) (Figure 1e); and *Lecanicillium* sp. (one isolate) (Figure 1f). Three mycosed cadavers had mixed infections of two fungal species: these were *I. tenuipes* and *I. cateniannulata* (one cadaver) and *I. fumosorosea* and *I. cateniannulata* (two cadavers). Of the isolates collected, *B. bassiana s.l.* and *I. fumosorosea* had also been recorded in previous preliminary field surveys (Saito et al. 2010); for this reason one isolate of each of these species (BB006 and IF002, respectively), both with reliable spore production capacities, were selected for evaluation as they were representative of the most common species affecting *P. melete* in wasabi fields.

3.2. Effect of volatiles from macerated wasabi leaves on conidial germination

There was no significant difference in germination between the *B. bassiana s.l.* and the *I. fumosorosea* isolates in response to wasabi leaves ($F_{1,40} = 3.14, P = 0.0839$) (Table 1). However, there was a significant negative effect of macerated wasabi leaves on germination of both isolates ($F_{4,40} = 3018.38, P < 0.0001$) (Table 1). Both *B. bassiana s.l.* and *I. fumosorosea* conidia had a germination rate of 100% in the control, and *B. bassiana s.l.* conidia exposed to volatiles from 1.3 g L⁻¹, 3.8 g L⁻¹, and 6.3 g L⁻¹ of macerated wasabi leaves had germination rates of 100%, 2.7% and 1.8%, respectively, while *I. fumosorosea* conidia exposed to the same levels of wasabi leaves had

germination rates of 100%, 4.5%, and 0%, respectively. Neither isolates germinated when exposed to volatiles from 13 g L⁻¹ of macerated wasabi leaves. There was a significant interaction between the quantity of wasabi leaves and isolate ($F_{4,40} = 3.14$, $P = 0.0245$) but only in one treatment (6.3 g L⁻¹ of wasabi leaves) indicating a significant difference in germination rate between the two isolates at this quantity of wasabi leaves ($F_{1,40} = 15.71$, $P = 0.0003$).

2.3. Effect of synthetic AITC on conidial germination

There was no significant difference in germination of *B. bassiana* s.l. and *I. fumosorosea* isolates in response to synthetic AITC ($F_{1,32} = 1.82$, $P = 0.1864$) (Table 2). However, there was a significant negative effect of synthetic AITC on germination of both isolates ($F_{3,32} = 1048.74$, $P < 0.0001$) (Table 2). Both *B. bassiana* s.l. and *I. fumosorosea* conidia had a germination rate of 100% in the control. *Beauveria bassiana* s.l. conidia exposed to 6.3 µg L⁻¹ and 63 µg L⁻¹ of AITC had germination rates of 99.0% and 12.3%, respectively, and *I. fumosorosea* conidia exposed to the same levels of AITC had germination rates of 90.5% and 15.1%, respectively. Neither isolate germinated when exposed to volatiles from 630 µg L⁻¹ of AITC. There was a significant interaction between the concentration of AITC and the isolate ($F_{3,32} = 6.67$, $P = 0.0013$) but only in one treatment (6.3 µg L⁻¹ of AITC) indicating a significant difference in germination rate between the two isolates at this concentration of AITC ($F_{1,32} = 19.07$, $P = 0.0001$).

3.4. Concentration of AITC in volatiles from macerated wasabi leaves

The quantity of AITC contained in volatiles released from macerated wasabi leaves (3.8

g L⁻¹) was quantified by gas chromatography. The standard curve for AITC was determined as $Y = 86.5X - 104.3$ ($F_{1,10} = 36.73$, $r^2 = 0.786$) for peak areas (Y) and log doses (X). Based on linear regression analysis, the concentration of AITC released from the sample leaves was $45 \pm 5 \mu\text{g L}^{-1}$ (mean \pm SE).

4. Discussion

There have been some studies describing interactions between plant volatiles and entomopathogenic fungi. For example, *Neozygites tanajoae* Delalibera Jr., Humber & Hajek produced more conidia when exposed to volatiles from cassava leaves damaged by mites (Hountondji et al. 2005). *Pandora neoaphidis* (Remaudière & Hennebert) Humber increased the number of conidia germinating on broad bean plants damaged by aphid feeding (Baverstock et al. 2005), and, in another study demonstrated that conidial germination of *P. neoaphidis* was inhibited by the volatiles released from tobacco plants in response to either aphid attack or artificial damage (Brown et al. 1995). These reports indicate that plant volatiles influence interactions between herbivores and entomopathogenic fungi in different ways, and that understanding these interactions is essential for the development of successful biological control strategies using fungal agents.

In the present study, we have demonstrated that volatiles from artificially damaged wasabi leaves, inhibited germination of *B. bassiana* s.l. and *I. fumosorosea* conidia (Table 1). This is likely to be as a result of the plant volatile AITC, because AITC levels in 3.8 g L⁻¹ of macerated leaves, which caused low germination rates (2.7% and 4.5% for *B. bassiana* s.l. and *I. fumosorosea*, respectively) (Table 1), was estimated at 45 $\mu\text{g L}^{-1}$ and a similar concentration (63 $\mu\text{g L}^{-1}$) of synthetic AITC also led to low

germination rates (12.3% and 15.1% for *B. bassiana s.l.* and *I. fumosorosea*, respectively) (Table 2).

For both fungi, their susceptibility to volatiles from macerated wasabi leaves or AITC was similar, with significant differences between them only occurring at one concentration in each experiment (Table 1 and 2). This negative effect of plant volatiles on entomopathogenic fungi is more severe than previously observed for fungal plant pathogens of cruciferous plants; for example, the fungal plant pathogens, *Helminthosporium solani* Dur. & Mont. and *Verticillium dahliae* Kleb., were able to grow a little at 7 mg L⁻¹ of AITC (Olivier et al., 1999), while in our study, germination of *B. bassiana s.l.* and *I. fumosorosea* was completely inhibited at just 630 µg L⁻¹ of AITC (Table 2). This suggests that entomopathogenic fungi may be less well adapted to plant volatiles than plant pathogens.

Inyang et al. (1999) examined toxicity of eleven isothiocyanates, potentially relating to cruciferous plants, on germination of *M. anisopliae* conidia and reported that two isothiocyanates (phenylethyl- and 3-butenyl isothiocyanate) decreased mortality of inoculated adults of the mustard beetle, *Phaedon cochleariae* (Fabricius), when exposed to vapours from 1 µL of each neat compound in a 55 mm-diameter glass chamber; its length was not described, but in the figure it appeared to be approximately 200 mm in length (about 500 mL in volume). The inhibitory concentration of these isothiocyanates (1 µL per 500 mL chamber) was apparently higher than the 63 µg L⁻¹ or 630 µg L⁻¹ of synthetic AITC that we observed resulted in little (12.3% and 15.1% for *B. bassiana s.l.* and *I. fumosorosea*, respectively) or no germination in both fungal species (Table 2). Therefore, such AITC levels may negatively affect the infection process of *B. bassiana s.l.* and *I. fumosorosea* and reduce herbivore mortality. While confirmation of this

requires further research beyond the scope of this study, it remains possible that wasabi volatiles may contribute to unsatisfactory biological control of herbivorous pests.

As mentioned previously, AITC plays an important role in the plant's defence strategy against herbivores and disease (Chew 1988; Louda & Mole 1991; Rask et al. 2000; Wittstock et al. 2003). However, in the evolutionary arms race between plants and their herbivores, specialist crucifer herbivores have undergone selection to overcome this defence. For example, in the small white butterfly, *Pieris rapae* L., the hydrolysis reaction is redirected toward the formation of nitriles such as allyl cyanide in the gut, rather than AITC, and this is then excreted in the faeces (Wittstock et al. 2004). The diamondback moth, *Plutella xylostella* L., has an enzyme, sulfatase, which desulfates the glucosinolates to metabolites that are not substrates for myrosinase and so do not produce AITC (Ratzka et al. 2002). In addition, these crucifer specialists use volatile AITC as an attractant to detect host plants (Hovanitz & Chang 1963; Hovanitz et al. 1963; Li et al. 2000; Rask et al. 2000), and their parasitoids also use the compound as a cue to detect their herbivore hosts (Fatouros et al. 2005; Geervliet et al. 1994, 1996; Mattiacci et al. 2001; Rask et al. 2000). Thus, AITC can enhance the fitness of both specialist herbivores and their natural enemies resulting in complex coevolving tritrophic interactions. Negative effects of AITC on entomopathogenic fungi may give an additional advantage to the host herbivores in such interactions.

Cruciferous plants emit AITC and many other isothiocyanates (Kumagai et al., 1994; Olivier et al., 1999); some of these other isothiocyanates have fungicidal activity against entomopathogenic fungi (Inyang et al., 1999) and the plant pathogens *H. solani* and *V. dahlia* (Olivier et al., 1999). In our study, other compounds in addition to AITC were detected in the volatiles from macerated wasabi leaves (data not shown) and it is

possible that these compounds may also have negative effects on conidial germination of *B. bassiana s.l.* and *I. fumosorosea*; further studies are required to examine the effects of these other compounds on entomopathogenic fungi.

In conclusion, we have provided evidence that volatiles (including AITC) from wasabi had negative effects on conidial germination of the entomopathogenic fungi, *B. bassiana s.l.* and *I. fumosorosea*. Successful development of biological control strategies for herbivores of wasabi, and other crucifers, must, therefore, be developed under a conceptual framework that accounts for plant volatile-mediated tritrophic interactions between plants, herbivores and their natural enemies including fungal pathogens, parasites and predators (Agrawal 2000; Cory & Hoover 2006; Dicke & Van Loon 2000; Elliot et al. 2000). AITC may play a key role in such interactions on crucifers such as wasabi, and its effects on biological control agents should be considered specifically, though this speculation needs to be confirmed by further laboratory and field studies.

Acknowledgments

We are grateful to Dr. H. Etoh for the AITC analysis of leaves, Dr. M. Shimazu for fungal identification and Professor I. Uwafuji for helpful advice on statistical analyses. We thank J. K. Pell Consulting for editing the manuscript.

References

- Agrawal A. 2000. Mechanisms, ecological consequences and agricultural implications of tri-trophic interactions. *Curr Opin Plant Biol.* 3:329–335.
- Agrawal A, Kurashige N. 2003. A role for isothiocyanates in plant resistance against the

305 specialist herbivore *Pieris rapae*. J Chem Ecol. 29:1403–1415.

306 Baverstock J, Elliot SL, Anderson PG, Pell JK. 2005. Response of the

307 entomopathogenic fungus *Pandora neoaphidis* to aphid-induced plant volatiles. J

308 Invertebr Pathol. 89:157–164.

309 Brown GC, Prochaska GL, Hildebrand DF, Nordin GL, Jackson DM. 1995. Green leaf

310 volatiles inhibit conidial germination of the entomopathogen *Pandora neoaphidis*

311 (Entomophthorales: Entomophthoraceae). *Environ Entomol* 24:1637–1647.

312 Chew FS. 1988. Biological effects of glucosinolates. In: Cutler HG, editor. Biologically

313 active natural products, Potential use in agriculture. Washington DC: American

314 Chemical Society: p. 155–181.

315 Cory JS, Hoover K. 2006. Plant-mediated effects in insect-pathogen interactions. Trends

316 Ecol Evol. 21:278–286.

317 Dicke M, Van Loon JA. 2000. Multitrophic effects of herbivore-induced plant volatiles

318 in an evolutionary context. Entomol Exp Appl. 97:237–249.

319 Elliot SL, Sabelis MW, Janssen A, Van der Geest LPS, Beerling EAM, Fransen J. 2000.

320 Can plants use entomopathogens as bodyguards? Ecol Lett. 3:228–235.

321 Eltayeb EM, Ahang G-A, Xie J-G, Mi F-Y. 2010. Study on the effect of

322 allyl isothiocyanate formulations on three Lepidopterous insect larvae, the diamond

323 back moth, *Plutella xylostella* (L.), the small cabbage white butterfly, *Pieris rapae*

324 (L.) and tobacco cut worm, *Spodoptera litula* Fabr. Am J Environ Sci. 6:168–176.

325 Fatouros NE, Van Loon JJA, Hordijk KA, Smid HM, Dicke M. 2005.

326 Herbivore-induced plant volatiles mediate in-flight host discrimination by

327 parasitoids. J Chem Ecol. 31:2033–2047.

328 Geervliet JBF, Vet LEM, Dicke M. 1994. Volatiles from damaged plants as major cues

329 in long-range host-searching by the specialist parasitoid *Cotesia rubecula*. Entomol
 330 Exp Appl. 73:289–297.

331 Geervliet JBF, Vet LEM, Dicke M. 1996. Innate responses of the parasitoids *Cotesia*
 332 *glomerata* and *C. rubecula* (Hymenoptera: Braconidae) to volatiles from different
 333 plant-herbivore complexes. J Insect Behav. 9:525–538.

334 Hall RA. 1981. The fungus *Verticillium lecanii* as a microbial insecticide against aphids
 335 and scales. In: Burgess HD, editor. Microbial control of pest and diseases 1970-1980.
 336 London: Academic Press: p. 483–498.

337 Helyer N, Gill G, Bywater A, Chambers R. 1992. Elevated humidities for control of
 338 chrysanthemum pests with *Verticillium lecanii*. Pestic Sci. 36:373–378.

339 Hountondji FCC, Sabelis MW, Hanna R, Janssen A. 2005. Herbivore-induced plant
 340 volatiles trigger sporulation in entomopathogenic fungi: the case of *Neozygites*
 341 *tanajoae* infecting the cassava green mite. J Chem Ecol. 31:1003–1021.

342 Hovanitz W, Chang VCS. 1963. Selection of allyl isothiocyanate by larvae of *Pieris*
 343 *rapae* and the inheritance of this trait. J Res Lepid. 1:169–182.

344 Hovanitz W, Chang VCS, Honch G. 1963. The effectiveness of different isothiocyanates
 345 on attracting larvae of *Pieris rapae*. J Res Lepid. 1:249–259.

346 Inyang EN, Butt TM, Doughty KJ, Todd AD, Archer S. 1999. The effects of
 347 isothiocyanates on the growth of the entomopathogenic fungus *Metarhizium*
 348 *anisopliae* and its infection of the mustard beetle. Mycol Res. 103:974–980.

349 Kumagai H, Kashima N, Seki T, Sakurai H, Ishii K, Ariga T. 1994. Analysis of volatile
 350 components in essential oil of upland *wasabi* and their inhibitory effects on platelet
 351 aggregation. Biosci Biotech Biochem. 58:2131–2135.

352 Li Q, Eigenbrode SD, Stringam GR, Thiagarajah MR. 2000. Feeding and growth of

353 *Plutella xylostella* and *Spodoptera eridania* on *Brassica juncea* with varying
354 glucosinolate concentrations and myrosinase activities. J Chem Ecol. 26:2401–2419.

355 Liang ZQ, Han YF, Chu HL, Liu AY. 2005. Studies on the genus *Paecilomyces* in China.
356 I. Fungal Divers. 20:83–101.

357 Louda S, Mole S. 1991. Glucosinolates: chemistry and ecology. In: Rosenthal GA,
358 Berenbaum MR, editors. Herbivores, their interactions with secondary plant
359 metabolites. 2nd edn. Vol. I. The chemical participants. London: Academic Press: p.
360 123–164.

361 Luangsa-ard JJ, Hywel-Jones NL, Manoch L, Samson RA. 2005. On the relationships of
362 *Paecilomyces* sect. *Isarioidea* species. Mycol Res. 109:581–589.

363 Luz C, Fargues J. 1999. Dependence of the entomopathogenic fungus, *Beauveria*
364 *bassiana*, on high humidity for infection of *Rhodnius prolixus*. Mycopathologia.
365 146:33–41.

366 Mattiacci L, Rudelli S, Rocca BA, Genini S, Dorn S. 2001. Systemically-induced
367 response of cabbage plants against a specialist herbivore, *Pieris brassicae*.
368 Chemoecology. 11:167–173.

369 Mayton HS, Oliver C, Vaughn SF, Loria R. 1996. Correlation of fungicidal activity of
370 *Brassica* species with allyl isothiocyanate production in macerated leaf tissue.
371 Phytopathology. 86:267–271.

372 Milner RJ, Lutton GG. 1986. Dependence of *Verticillium lecanii* (Fungi:
373 Hyphomycetes) on high humidities for infection and sporulation using *Myzus*
374 *persicae* (Homoptera: Aphididae) as hosts. Environ Entomol. 15:380–382.

375 Nakata M. 1963. Notes on the natural enemies of larvae and pupae of *Pieris melete*
376 Ménétriès injurious to the Japanese horseradish. Jpn J Appl Entomol Zool. 7:348–

377 349 (in Japanese).

378 Oliver C, Vaughn SF, Mizubuti CSG, Loria R. 1999. Variation in allyl isothiocyanate
 379 production within *Brassica* species and correlation with fungicidal activity. J Chem
 380 Ecol. 25:2687–2701.

381 Rask L, Andréasson E, Ekbom B, Eriksson S, Pontoppidan B, Meijer J. 2000.
 382 Myrosinase: gene family evolution and herbivore defense in Brassicaceae. Plant
 383 Mol Boil. 42:93–113.

384 Ratzka A, Vogel H, Kliebenstein DJ, Mitchell-Olds T, Kroymann J. 2002. Disarming the
 385 mustard oil bomb. Proc Natl Acad Sci USA. 99:11223–411228.

386 Rehner SA, Minnis AM, Sung G-H, Luangsa-ard JJ, Devotto L, Humber RA. 2011.
 387 Phylogeny and systematics of the anamorphic entomopathogenic genus *Beauveria*.
 388 Mycologia. 103:1055–1073.

389 Saito T, Yabuta M. 1996. Laboratory studies on effect of pesticides on
 390 entomopathogenic fungus, *Verticillium lecanii*. Jpn J Appl Entomol Zool. 40:71–76
 391 (in Japanese).

392 Saito T, Sugiyama Y, Haga H, Sugiyama K. 2010. Integrated pest management in
 393 wasabi. 4. Parasites of the striated white, *Pieris melete* Ménétriès, in Shizuoka
 394 Prefecture. Ann Rept Kansai Pl Prot. 52:135–137 (in Japanese).

395 Samson RA. 1974. *Paecilomyces* and some allied Hyphomycetes. Stud Mycol. 6:1–119.

396 Shimazu M. 2001. *Paecilomyces cateniannulatus* Liang, a commonly found, but an
 397 unrecorded entomopathogenic fungus in Japan. Appl Entomol Zool. 36:283–288.

398 Shinohara S, Fitriana Y, Satoh K, Narumi I, Saito T. 2013. Enhanced fungicide
 399 resistance in *Isaria fumosorosea* following ionizing radiation-induced mutagenesis.
 400 FEMS Microbiol Lett. 349:54–60.

401 SPSS. 2009. PASW Statistics 18. Chicago: SPSS Inc.

402 Sultana T, Savage GP, McNeil DL, Porter NG, Clark B. 2003. Comparison of flavor
 403 compounds in wasabi and horseradish. J Food Agr Environ. 1:117–121.

404 Wittstock U, Halkier A. 2002. Glucosinolate research in the *Arabidopsis* era. Trends
 405 Plant Sci. 7:263–270.

406 Wittstock U, Kliebenstein DJ, Lambrix V, Reichelt M, Gershenzon J. 2003.
 407 Glucosinolate hydrolysis and impact on generalist and specialist insect herbivores.
 408 Recent Adv Phytochem. 37:101–125.

409 Wittstock U, Agerbirk N, Stauber E, Oisen CE, Hippler M, Mitchell-Olds T,
 410 Gershenzon J, Vogel H. 2004. Successful herbivore attack due to metabolic
 411 diversion of a plant chemical defense. Proc Natl Acad Sci USA. 101:4859–4864.

412 Zare R, Gams W. 2001. A revision of *Verticillium* section *Prostrata*. IV. The genera
 413 *Lecanicillium* and *Simplicillium* gen. nov. Nova Hedwigia. 73:1–50.

414 Zare R, Gams W. 2008. A revision of the *Verticillium fungicola* species complex and its
 415 affinity with the genus *Lecanicillium*. Mycol Res. 112:811–824.

Table 1. Percent germination of *B. bassiana s.l.* and *I. fumosorosea* conidia exposed to different quantities of macerated wasabi leaves for 24 h in a chamber.

Quantity of macerated wasabi leaves (g L ⁻¹ chamber)	Percent germination (mean ± SE)	
	<i>B. bassiana s.l.</i>	<i>I. fumosorosea</i>
0	100	100
1.3	100	100
3.8	2.7 ± 1.2	4.5 ± 4.0
6.3	1.8 ± 0.5	0
13	0	0

Table 2. Percent germination of *B. bassiana s.l.* or *I. fumosorosea* conidia exposed to different quantities of allyl isothiocyanate (AITC) for 24 h in a chamber.

Dose of AITC ($\mu\text{g L}^{-1}$ chamber)	Percent germination (mean \pm SE)	
	<i>B. bassiana s.l.</i>	<i>I. fumosorosea</i>
0	100	100
6.3	99.0 \pm 0.3	90.5 \pm 2.8
63	12.3 \pm 3.6	15.1 \pm 0.6
630	0	0





Figure caption

Figure 1. (a) A wasabi field. (b)–(f) Mycosed pupae of *P. melete* infected by *B. bassiana* s.l., *I. fumosorosea*, *I. tenuipes*, *I. cateniannulata*, and *Lecanicillium* sp., respectively.