

Plant growth regulatory compounds from the mushroom *Russula vinosa*

メタデータ	言語: eng 出版者: 公開日: 2017-08-28 キーワード (Ja): キーワード (En): 作成者: Matsuzaki, Nobuo, Wu, Jing, Kawaide, Misaki, Choi, Jae-Hoon, Hirai, Hirofumi, Kawagishi, Hirokazu メールアドレス: 所属:
URL	http://hdl.handle.net/10297/10377

Note

Plant growth regulatory compounds from the mushroom *Russula vinosa*

Nobuo Matsuzaki ^a, Jing Wu ^b, Misaki Kawaide ^c, Jae-Hoon Choi ^{b,c}, Hirofumi Hirai ^{b, c},
Hirokazu Kawagishi ^{a, b, c, *}

^a Graduate School of Science and Technology, Shizuoka University, 836 Ohya, Suruga-ku,
Shizuoka 422-8529, Japan

^b Research Institute of Green Science and Technology, Shizuoka University, 836 Ohya,
Suruga-ku, Shizuoka 422-8529, Japan

^c Graduate School of Agriculture, Shizuoka University, 836 Ohya, Suruga-ku, Shizuoka 422-
8529, Japan

* Correspondence author:
H. Kawagishi
Tel: +81-54-238-4885
Fax: +81-54-238-4885
E-mail: kawagishi.hirokazu@shizuoka.ac.jp

Text: 7 pages; figures: 2

1 Abstract

2
3 Five compounds, (1*R*,2*S*)-1-phenylpropane-1,2-diol, isolactarorufin, lactarorufin A, 8 α ,13-
4 dihydroxy-marasm-5-oic acid γ -lactone, and 7 α ,8 α ,13-trihydroxy-marasm-5-oic acid γ -
5 lactone were isolated from the fruiting bodies of *Russula vinosa*. In the bioassay examining
6 plant-growth regulatory activity using lettuce, all the compounds regulated the growth.
7

8
9
10
11 *Keywords*

12
13 Plant growth regulator; Structural identification
14
15
16
17
18
19
20
21
22
23
24
25
26
27
28
29
30
31
32
33
34
35
36
37
38
39
40
41
42
43
44
45
46
47
48
49
50
51
52
53
54
55
56
57
58
59
60
61
62
63
64
65

We have reported the isolation of plant growth regulators, 2-azahypoxanthine and imidazole-4-carboxamide, from the fungus *Lepista sordida* that form “fairy rings” (Choi et al. 2010a, b). These compounds increased yields of rice and wheat in field experiments (Tobina et al. 2014; Asai et al. 2015). Based on these studies, we are screening for plant growth regulators from fungi that interact with plants. During the screening using lettuce, we found strong activity in the extracts of the fruiting bodies of *Russula vinosa*.

Russula vinosa is an edible wild mushroom with high medicinal value (Chen et al. 2007). The extracts of *R. vinosa* have an inhibition effect on bacteria, yeasts and molds (Li et al. 1998).

Here, we describe the isolation and identification of plant growth regulators from the fruiting bodies of this species.

Extraction and isolation

Fruiting bodies of *R. vinosa* were collected in Wuyi Mountains of Fujian Province in China in Aug 2010. There were *Cyclobalanopsis chungii* nearby the fruiting bodies. The collected fruiting bodies were air-dried soon after collection. The dried fruiting bodies were extracted and fractionated twice. In the first time, 1.5 kg of the fruiting bodies was crushed in a blender and the crushed fruiting bodies were extracted with hexane and then EtOAc. The EtOAc-soluble part (25.1 g) was fractionated by silica gel flash column chromatography (Silica gel 60 N, Kanto Chemical, Tokyo, Japan; CH₂Cl₂ (fraction 1); 90% (fractions 2 to 3), 80% (fractions 4 to 5), 70% (fractions 6 to 7), 60% (fraction 8), 40% (fractions 9 to 10), 30% (fraction 11), 10% (fractions 12 to 13) CH₂Cl₂/EtOAc; 90% (fraction 14), 70% (fraction 15), 50% (fractions 16 to 17), 40% (fractions 18 to 20), 30% (fraction 21), 20% (fractions 22 to 23), 10% (fraction 24) EtOAc/MeOH; MeOH; 80% (fraction 25), 70% (fraction 26) MeOH/acetic acid) to obtain 26 fractions (fractions 1 to 26). HPLC separations were performed with a Jasco Gulliver system. Fraction 14 (301 mg) was separated by reverse-phase HPLC (Develosil C30-UG-5, Nomura Chemical, Seto, Japan; 40% MeOH) to afford 1 (2.0 mg). Fraction 15 (173 mg) was fractionated by ODS gel flash column chromatography (Cosmosil 140 C18-OPN, Nacalai Tesque, Kyoto, Japan; 90%, 98%, 99% MeOH, MeOH) to obtain 10 fractions (fractions 15-1 to 15-10). Fraction 15-3 (59.0 mg) was further separated by reverse-phase HPLC (Develosil C30-UG-5, 55% MeOH) to obtain 9 fractions (fractions 15-3-1 to 15-3-9). Fraction 15-3-4 (5.9 mg) was fractionated by reverse-phase HPLC (Cosmosil πNAP, Nacalai Tesque, Kyoto, Japan; 50% MeOH) to afford 2 (2.7 mg). In the second time, 5.0 kg of the fruiting bodies was crushed in a blender and the crushed fruiting bodies were extracted with hexane and then EtOAc. The EtOAc-soluble part (41.5 g) was fractionated by silica gel flash column chromatography (Silica gel 60 N, Kanto Chemical, Tokyo, Japan; CH₂Cl₂ (fraction 1’); 90% (fraction 2’), 80% (fraction 3’), 60% (fraction 4’), 40% (fraction 5’), 30% (fraction 6’), 10% (fraction 7’) CH₂Cl₂/EtOAc; 90% (fraction 8’), 70% (fraction 9’), 60% (fraction 10’), 50% (fraction 11’), 40% (fraction 12’), 30% (fraction 13’), 20% (fraction 14’) EtOAc/MeOH; MeOH (fractions 15’ to 16’); 70% (fraction 17’) MeOH/acetic acid) to obtain 17 fractions (fractions 1’-17’), respectively. Fraction 8’ (333 mg) and fraction 9’ (1.19 g) were separated by ODS gel flash column chromatography (70%, 80%, 90% MeOH, MeOH) to give 13 and 19 fractions (fractions 8’-1 to 8’-13 and fractions

9'-1 to 9'-19), respectively. Fraction 8'-2 (16.7 mg) was separated by reverse-phase HPLC (Develosil C30-UG-5, 55% MeOH) to afford 4 (1.8 mg) and 5 (1.4 mg). Fraction 9'-2 (40.9 mg) was separated by reverse-phase HPLC (Develosil C30-UG-5, 55% MeOH) to obtain 23 fractions (fractions 9'-2-1 to 9'-2-23) and fraction 9'-2-13 (5.1 mg) was further separated by reverse-phase HPLC (Cosmosil π NAP, 50% MeOH), affording 3 (1.1 mg).

¹H NMR spectra (one- and two-dimensional) were recorded on a Jeol lambda-500 spectrometer at 500 MHz, and ¹³C NMR spectra were recorded on the same instrument at 125 MHz (Jeol, Tokyo, Japan). ESIMS spectra were measured on a JMS-T100LC mass spectrometer (Jeol, Tokyo, Japan). IR spectra were recorded on a FT/IR-4100 (Jasco, Tokyo, Japan).

We obtained five compounds from the fruiting bodies of *R. vinosa*.

Compound 1: white crystals; ESIMS m/z 175 [M+Na]⁺; IR (neat): 3351 cm⁻¹; ¹H-NMR (in CD₃OD) δ_H : 1.10 (d, J = 6.4), 2.03, 4.00 (dq, J = 4.0, 6.4), 2.64, 4.48 (d, J = 5.2), 7.30 (m, J = 7.6); ¹³C-NMR, δ_C : 18.1, 72.4, 79.1, 128.0, 128.3, 129.0, 143.4.

Compound 2: white crystals; ESIMS m/z 289 [M+Na]⁺; IR (neat): 3392 cm⁻¹; ¹H-NMR (in CD₃OD) δ_H : 0.95 (s), 1.0 (d), 1.08 (s), 1.18 (dd), 1.45 (ddd), 1.53 (s), 1.65 (dd), 1.70 (d), 1.81 (ddd), 1.93 (m), 2.12 (m), 4.16 (d), 4.25 (d), 4.28 (d); ¹³C-NMR δ_C : 18.6, 24.4, 27.3, 29.8, 34.6, 39.1, 39.4, 41.3, 44.3, 47.7, 48.8, 72.1, 73.4, 75.6, 178.1.

Compound 3: white amorphous; ESIMS m/z 289 [M+Na]⁺; IR (neat): 3221 cm⁻¹; ¹H-NMR (in CD₃OD) δ_H : 1.03 (s), 1.09 (s), 1.21 (s), 1.37 (d), 1.40 (m), 1.60 (d), 1.70 (m), 2.46 (d), 2.60 (m), 2.62 (dd), 2.69 (d), 4.32 (d), 4.74 (d), 4.88 (d); ¹³C-NMR δ_C : 28.3, 29.3, 30.4, 36.7, 37.4, 45.0, 46.3, 47.1, 52.2, 69.1, 72.6, 74.0, 124.1, 164.3, 177.4.

Compound 4: white amorphous; ESIMS m/z 273 [M+Na]⁺; IR (neat): 3392 cm⁻¹; ¹H-NMR (in CD₃OD) δ_H : 1.04 (s), 1.06 (d), 1.13 (s), 1.24 (d), 1.25 (s), 1.50 (m), 1.54 (m), 1.62 (d), 1.65 (m), 1.85 (m), 2.41 (d), 2.64 (m), 3.23 (t), 4.17 (m), 4.69 (m); ¹³C-NMR δ_C : 17.8, 30.2, 30.9, 32.5, 32.7, 36.7, 37.8, 42.7, 44.7, 45.6, 46.2, 46.7, 73.2, 73.8, 180.6.

Compound 5: white amorphous; ESIMS m/z 289 [M+Na]⁺; IR (neat): 3335 cm⁻¹; ¹H-NMR (in CD₃OD) δ_H : 0.74 (d), 0.93 (s), 1.03 (s), 1.41 (s), 1.47 (q), 1.60 (d), 1.80 (q), 1.82 (d), 2.17 (d), 2.25 (d), 2.76 (m), 3.65 (d), 4.12 (d), 4.51 (d); ¹³C-NMR δ_C : 17.8, 27.4, 28.9, 29.6, 30.0, 33.6, 38.4, 45.0, 45.2, 46.9, 50.6, 74.5, 76.3, 79.6, 180.5.

Their structures were identified by interpretation of their spectroscopic data and all the compounds have been reported previously (Fig. 1). Compound 1, (1*R*,2*S*)-1-phenylpropane-1,2-diol, has been isolated from *Trametes* sp. (Brambilla et al. 1995). Compound 2 was identified as isolactarorufin and has been isolated from *Lactarius piperatus* (Wang et al. 2003). Compound 3 was lactarorufin A, which has been isolated from *Lactarius vellereus* (Kamo et al. 2006), *Strobilurus stephanocystis* (Hiramatsu et al. 2008) and *Russula emetica* (Kobata et al. 1995). Lactarorufin A showed selective cytotoxicity against the A549, and HCT-15 cell lines (Kim et al. 2010) and anti-feedant activity of insects *Sitophilus granarius* (Daniewski et al. 1993). Compound 4, 8 α ,13-dihydroxy-marasm-5-oic acid γ -lactone, has been isolated from *L. vellereus* (Daniewski et al. 2003) and *Russula foetens* (Wang et al. 2006). Compound 5, 7 α ,8 α ,13-trihydroxy-marasm-5-oic acid γ -lactone, was isolated from *R. foetens* (Wang et al. 2006) and *L. piperatus* (Wang et al. 2003).

Plant growth regulatory activity

Lettuce seeds (*Lactuca sativa* L. cv. Cisko; Takii Co., Ltd., Tokyo, Japan) were put on filter paper (Advantec No. 2, ϕ 55 mm; Toyo Roshi Kaisha, Japan), soaked in distilled water in a Petri dish (ϕ 60 \times 20 mm), and incubated in a growth chamber in the dark at 25 °C for 1 d. Each sample was dissolved in 1 mL of MeOH (1, 10^{-1} , 10^{-2} and 10^{-3} μ mol/mL) and then poured on filter paper (ϕ 55 mm) in a Petri dish (ϕ 60 \times 20 mm). After the sample-loaded paper had been air-dried, 1 mL of distilled water was poured on the sample-loaded paper or intact filter paper (control). The preincubated lettuces (n = 9 in each Petri dish) were transferred onto the sample-loaded filter paper or control filter paper and incubated in a growth chamber in the dark at 25 °C for 3 d. The lengths of the hypocotyl and the root were measured using a ruler.

Effects of all the compounds on the growth of lettuce were examined (Fig. 2). Compound 4 weakly inhibited the root and hypocotyl growth of lettuce at 1 μ mol/paper, while compound 3 showed an inhibition as low as 100 nmol/paper. As for the root growth of lettuce, compound 2 showed promotion at 10 and 100 nmol/paper, but showed inhibition at 1 nmol/paper and 1 μ mol/paper. As for the hypocotyl growth of lettuce, compound 1 showed inhibition at 10 and 100 nmol/paper and promotion at 1 nmol/paper, compound 2 showed a promotion at 100 nmol/paper, compound 5 showed promotion at lower doses (1 and 10 nmol/paper).

We isolated five compounds from the mushroom *R. vinosa*. The structures were identified by spectroscopic analyses. Although these compounds have been already known as mentioned above, their biological activity against plant was first found in this study. In the previous report (Kamo et al. 2006), compound 3 at 3.6×10^2 μ M showed no growth activity against lettuce, while this compound at 100 nmol/paper inhibited lettuce growth in this study. The reason of the difference between the two results might be due to the difference of the procedures of the bioassay. However, our results will provide useful information for the development of plant-growth regulators.

Disclosure

The authors declare no conflict of interest. All the experiments undertaken in this study comply with the current laws of Japan.

Acknowledgments

We thank Dr. V. K. Deo (Shizuoka University) for valuable discussion, and Mr. H. Oka for collecting the mushroom.

REFERENCES

- Asai T, Choi JH, Ikka T, Fushimi K, Abe N, Tanaka H, Yamakawa Y, Kobori H, Kiriiwa Y, Motohashi R, Deo VP, Asakawa T, Kan T, Morita A, Kawagishi H, 2015. Effect of 2-azahypoxanthine (AHX) produced by the fairy-ring-forming fungus on the growth and

- the grain yield of rice. *Japan Agricultural Research Quarterly* 49: 45–49;
<http://dx.doi.org/10.6090/jarq.49.45>.
- Brambilla U, Nasini G, Vajna de Pava O, 1995. Secondary mold metabolites, part 49.
 Isolation, structural elucidation, and biomimetic synthesis of trametol, a new 1-
 arylpropane-1,2-diol produced by the fungus *Trametes* sp. *Journal of Natural Products*
 58: 1251–1253.
- Chen YH, Chen ZM, 2007. Effect of mycorrhizal biodiversity on fine root mass of
Cyclobalanopsis chungii and basidiocarp collection of *Russula vinosa*. *Chinese Journal*
of Eco-Agriculture 15: 171–173.
- Choi JH, Abe N, Tanaka H, Fushimi K, Nishina Y, Morita A, Kiriiwa Y, Motohashi R,
 Hashizume D, Koshino H, Kawagishi H, 2010b. Plant-growth regulator, imidazole-4-
 carboxamide produced by fairy-ring forming fungus *Lepista sordida*. *Journal of*
Agricultural and Food Chemistry 58: 9956–9959; <http://dx.doi.org/10.1021/jf101619a>.
- Choi JH, Fushimi K, Abe N, Tanaka H, Maeda S, Morita A, Hara M, Motohashi R,
 Matsunaga J, Eguchi Y, Ishigaki N, Hashizume D, Koshino H, Kawagishi H, 2010a.
 Disclosure of the “fairy” of fairy-ring forming fungus *Lepista sordida*. *ChemBioChem* 11:
 1373–1377; <http://dx.doi.org/10.1002/cbic.201000112>.
- Daniewski WM, Gumulka M, Ptaczynska K, Skibicki P, Bloszyk E, Drozd B, Stromberg S,
 Norin T, Holub M, 1993. Antifeedant activity of some sesquiterpenoids of the genus
Lactarius (Agaricales: Russulaceae). *European Journal of Entomology* 90: 65–70.
- Daniewski WM, Gumulka M, Ptaczynska K, Skibicki P, Krajewski J, Gluzinski P, 2003.
 Marasmane lactones from *Lactarius vellereus*. *Phytochemistry* 31: 913–915;
[http://dx.doi.org/10.1016/0031-9422\(92\)80038-G](http://dx.doi.org/10.1016/0031-9422(92)80038-G).
- Hiramatsu F, Murayama T, Koseki T, Shiono Y, 2008. A new secolactarane-type sesquiterpene
 from *Strobilurus stephanocystis*. *Natural Product Research* 22: 1001–1006.
- Kamo T, Matsue M, Kashiwabara M, Hirota M, 2006. 1,2-Dehydrolactarolide A, a new plant
 growth regulatory lactarane sesquiterpene from *Lactarius vellereus*. *Bioscience,*
Biotechnology, & Biochemistry 70: 2307–2309. <http://dx.doi.org/10.1271/bbb.60148>.
- Kim KH, Noh HJ, Choi SU, Park KM, Seok SJ, Lee KR, 2010. Lactarane sesquiterpenoids
 from *Lactarius subvellereus* and their cytotoxicity. *Bioorganic & Medicinal Chemistry*
Letters 20: 5385–5388; <http://dx.doi.org/10.1016/j.bmcl.2010.07.119>.
- Kobata K, Kano S, Shibata H, 1995. New lactarane sesquiterpenoid from the fungus *Russula*
emetica. *Bioscience, Biotechnology, & Biochemistry* 59: 316–318.
- Li HZ, Huang DX, Xu XP, Zhang DC, Shi QQ, Wu SG, 1998. Studies on the chemical
 compositions in *Russula vinosa*. *Mycosystema* 17: 68–74.
- Tobina H, Choi JH, Asai T, Kiriiwa Y, Asakawa T, Kan T, Morita A, Kawagishi H, 2014.
 2-Azahypoxanthine and imidazole-4-carboxamide produced by the fairy-ring-forming
 fungus increase yields of wheat. *Field Crops Research* 162: 6–11.
- Wang XN, Shen JH, Du JC, Liu JK, 2006. Marasmane sesquiterpenes isolated from
Russula foetens. *The Journal of Antibiotics* 59: 669–672.
- Wang Y, Yang SP, Yue JM, 2003. Novel sesquiterpenes from the fungus *Lactarius piperatus*.
Helvetica Chimica Acta 86: 2424–2433.

Figure legends

- 1
- 2 Fig. 1– Structures of compounds 1–5.
- 3 Fig. 2– Effect of 1 to 5 on the growth of lettuce. Lettuce seedlings were treated with 1 to 5.
- 4 Respective length of growth compared with the control \pm standard deviation (* $p < 0.05$, ** $p <$
- 5 0.01 vs control, n = 7)

Highlights

- Five compounds were isolated from the fruiting bodies of *Russula vinosa*.
- All the compounds regulated lettuce growth.
- The biological activity of the compounds against plant was first found in this study.




Figure 1. Matsuzaki *et al.*




Figure 2 Matsuzaki *et al.*