

Plant growth regulators from the edible mushroom *Leccinum extremiorientale*

著者	Ito Akinobu, Wu Jing, Ozawa Nobuhiko, Choi Jae-Hoon, Hirai Hirofumi, Kawagishi Hirokazu
journal or publication title	Mycoscience
volume	58
number	6
page range	383-386
year	2017-11
出版者	Elsevier
権利	(C) 2017. This manuscript version is made available under the CC-BY-NC-ND 4.0 license http://creativecommons.org/licenses/by-nc-nd/4.0/
URL	http://hdl.handle.net/10297/00025654

doi: 10.1016/j.myc.2017.04.008

Mycoscience

Elsevier Editorial System(tm) for

Manuscript Draft

Manuscript Number: MYC-D-17-00019R2

Title: Plant growth regulators from the edible mushroom *Leccinum extremiorientale*

Article Type: Notes

Corresponding Author: Professor Hirokazu Kawagishi,

Corresponding Author's Institution: Shizuoka University

First Author: Akinobu Ito

Order of Authors: Akinobu Ito ; Jing Wu, Ph.D.; Nobuhiko Ozawa; Jae-Hoon Choi, Ph.D.; Hirofumi Hirai Hirofumi Hirai , Ph.D.; Hirokazu Kawagishi

Abstract: Two compounds were isolated from the edible mushroom *Leccinum extremiorientale*. In the bioassay examining plant growth regulatory activity using lettuce, compound 1 promoted the root growth as low as 100 nmol/paper dose-dependently and inhibited the hypocotyl growth at 1000 nmol/paper. Compound 2 inhibited the root and hypocotyl growth at 1000 nmol/paper.

1 **Note**

2
3 **Plant growth regulators from the edible mushroom *Leccinum extremiorientale***

4
5 Akinobu Ito ^a, Jing Wu ^b, Nobuhiko Ozawa ^a, Jae-Hoon Choi ^{a,b}, Hirofumi Hirai ^{a,b}, Hirokazu
6 Kawagishi ^{a,b,c,*}

7
8 ^a Graduate School of Integrated Science and Technology, Shizuoka University, 836 Ohya,
9 Suruga-ku, Shizuoka 422-8529, Japan

10 ^b Research Institute of Green Science and Technology, Shizuoka University, 836 Ohya,
11 Suruga-ku, Shizuoka 422-8529, Japan

12 ^c Graduate School of Science and Technology, Shizuoka University, 836 Ohya, Suruga-ku,
13 Shizuoka 422-8529, Japan

14
15 * Corresponding author.

16 H. Kawagishi

17 Tel: +81 54 238 4885.

18 Fax: +81 54 238 4885.

19 E-mail address: kawagishi.hirokazu@shizuoka.ac.jp

20
21 Text: 6 pages; figures: 2

1 ABSTRACT

2
3 Two compounds were isolated from the edible mushroom *Leccinum extremiorientale*. In the
4 bioassay examining plant growth regulatory activity using lettuce, compound **1** promoted the
5 root growth as low as 100 nmol/paper dose-dependently and inhibited the hypocotyl growth at
6 1000 nmol/paper. Compound **2** inhibited the root and hypocotyl growth at 1000 nmol/paper.
7
8
9

10
11 *Keywords:*

12 Structural identification
13
14
15
16
17
18
19
20
21
22
23
24
25
26
27
28
29
30
31
32
33
34
35
36
37
38
39
40
41
42
43
44
45
46
47
48
49
50
51
52
53
54
55
56
57
58
59
60
61
62
63
64
65

1 Fungi are good sources of plant growth regulators, and we have been searching for the
2 regulators from various fungi. For example, we have reported the plant growth regulators, 2-
3 azahypoxanthine and imidazole-4-carboxamide, produced by a fairy ring-forming fungus
4 *Lepista sordida* (Choi et al. 2010a, b). Furthermore, these compounds increased yields of rice
5 and wheat in field experiments (Tobina et al. 2014; Asai et al. 2015). We have also reported
6 the isolations of agrocybynes A to E from *Agrocybe praecox*, erinaceolactones A to C from
7 *Hericium erinaceus*, (1*R*,2*S*)-1-phenylpropane-1,2-diol, isolactarorufin, lactarorufin A, 8 α ,13-
8 dihydroxy-marasm-5-oic acid γ -lactone, and 7 α ,8 α ,13-trihydroxy-marasm-5-oic acid γ -
9 lactone from *Russula vinosa* (Fushimi et al. 2012; Wu et al. 2015; Matsuzaki et al. 2016). All
10 these compounds showed the growth regulatory activity against lettuce. During further
11 screening for plant growth regulators using lettuce, we found the activity in the extracts of the
12 fruiting bodies of *Leccinum extremiorientale*.

13 The edible mushroom *L. extremiorientale* (Japanese name, Akayamadori) belongs to the
14 genus *Leccinum* in the family Boletaceae, and it can be seen from summer to autumn. The
15 mushroom has a red brown areolate cap and distributes mainly in the northern temperate zone.
16 The exhaustive studies on chemical constituents of some species in this family Boletaceae
17 have been carried out (Kovganko et al. 1999; Hellwig et al. 2002; Kim et al. 2006;
18 Kukovinets et al. 2006). In our previous research on the bioactive compounds from the
19 mushroom, two sterols showed the ability to suppress the formation of osteoclasts, and
20 leccinine A (**2**) showed protective activity against the endoplasmic reticulum stress-dependent
21 cell death (Choi et al. 2010c, 2011). In order to find plant growth regulatory compounds from
22 the mushroom, we carried out a chemical investigation of the mushroom.

23 Here, we describe the isolation, structural identification, and growth regulatory activity
24 against lettuce of compounds **1** and **2**.

25 **Extraction and isolation**

26 The fresh fruiting bodies of *L. extremiorientale* were collected at Narusawa village,
27 Yamanashi Prefecture in Japan, in Aug 2007. The fresh fruiting bodies of *L. extremiorientale*
28 (14.9 kg) were extracted with EtOH (20 L, three times) and then with acetone (10 L, once).
29 After the solution was combined and evaporated under reduced pressure, the concentrate of
30 the extracts was divided into *n*-hexane-soluble, EtOAc-soluble and water-soluble parts. The *n*-
31 hexane-soluble part (58.0 g) and the EtOAc-soluble part (32.4 g) showed the growth
32 regulatory activity against lettuce. In this study, the *n*-hexane soluble part was fractionated by
33 silica gel flash column chromatography (50%, 20% *n*-hexane/CH₂Cl₂; 90%, 50%
34 CH₂Cl₂/acetone; acetone and MeOH, 2.0 L each) to obtain 16 fractions (fractions 1 to 16).
35 Fraction 14 (814 mg) was further separated by reverse-phase HPLC (Develosil C30-UG-5;
36 column size, ϕ 20 \times 250 mm; UV wavelength, 210 nm; flow rate, 5 mL/min; eluent, 95%
37 MeOH) and 21 fractions were obtained (fractions 14-1 to 14-21). Fraction 14-8 (42.8 mg;
38 retention time, 16 min) was subjected to recrystallization in diethyl ether and *n*-hexane, and
39 compound **1** (24.2 mg) was obtained from this fraction.

40 **Identification of active compounds**

41 ¹H NMR spectra (one- and two-dimensional) were recorded on a JEOL lambda-500
42 spectrometer (JEOL Ltd., Tokyo, Japan) at 500 MHz, while ¹³C NMR spectra were recorded
43 by the same instrument at 125 MHz. HRESIMS data were measured by a JMS-T100LC mass
44 spectrometer (JEOL Ltd., Tokyo, Japan). HPLC separation was performed with a JASCO
45 Gulliver system (JASCO Co., Tokyo, Japan) using a reverse-phase HPLC column (Develosil
46 C30-UG-5, ϕ 20 \times 250 mm, Nomura Chemical, Seto, Japan). IR spectrum was recorded on a
47
48
49
50

1 FT/IR-4100 (JASCO Co., Tokyo, Japan). The specific rotation value was measured with a
2 JASCO DIP-1000 polarimeter (JASCO Co., Tokyo, Japan). Silica cartridges and C18
3 cartridges (Nihon Waters K.K., Tokyo, Japan) were used in the pro-processing of the samples.
4 Silica gel plate (TLC Silica gel 60 F₂₅₄, Merck KGaA, Darmstadt, Germany) and silica gel
5 60N (Kanto Chemical Co., Inc., Tokyo, Japan) were used for analytical TLC and for flash
6 column chromatography, respectively.

7 Compound **1** was purified as colorless crystals. We measured the NMR spectra including
8 DEPT, COSY, HMQC and HMBC. The structure of compound **1** was identified to be (8*E*,
9 12*Z*)-10,11-dihydroxyoctadeca-8,12-dienoic acid by comparison of their spectroscopic data
10 with those that were reported previously (Fig. 1). This compound has been isolated from
11 *Aspergillus flavus*, and exhibited low inhibitory activity to acetylcholinesterase that is
12 involved in progressing Alzheimer's disease (Lopez et al. 2002; Qiao et al. 2011). The value
13 of specific rotation of compound **1** (+ 37.0) was similar to that reported previously (+ 28.2)
14 and the NMR data of compound **1** were also identical to that in the previous report,
15 suggesting that absolute configuration of compound **1** was the same as that in the report (Qiao
16 et al. 2011). However, the absolute configurations at C-10 and -11 of compound **1** has been
17 still unknown. There is no other report of isolation of compound **1** from natural sources.

18 Compound **1**: colorless crystals; ESIMS *m/z* 335 [M+Na]⁺; [α]_D²⁸ + 37.0 (*c* 0.10,
19 MeOH); ¹H NMR (in CD₃OD) δ_H: 0.86 (t, 7.0, 3H), 1.27 (m), 1.27 (m), 1.35 (m), 1.37 (m),
20 1.37 (m), 1.37 (m), 1.61 (m, 2H), 2.03 (m, 2H), 2.06 (m, 2H), 2.31 (t, 7.3, 2H), 4.04 (dd, 7.3,
21 4.3, 1H), 4.42 (dd, 8.5, 4.3, 1H), 5.37 (dd, 11.3, 8.5, 1H), 5.46 (dd, 15.6, 7.3, 1H), 5.61 (m,
22 1H), 5.71 (m, 1H); ¹³C NMR (in CD₃OD) δ_C: 14.6, 22.5, 24.5, 28.0, 28.5, 28.6, 28.7, 29.3,
23 31.5, 32.2, 33.9, 70.5, 75.7, 127.3, 127.7, 135.1, 135.2, 178.8.

24 Leccinine A (**2**) has been isolated as an endoplasmic reticulum stress-suppressive
25 compound from *L. extremiorientale* by us, however, activity towards plant growth has not
26 been examined (Choi et al. 2011).

27 **Plant growth regulatory activity**

28 Lettuce seeds (*Lactuca sativa* L. cv. Cisko; Takii Co., Ltd., Tokyo, Japan) were put on filter
29 paper (Advantec No. 2, φ 55 mm; Toyo Roshi Kaisha, Japan), soaked in distilled water in a
30 Petri dish (φ 60 × 20 mm), and incubated in a growth chamber in the dark at 23 °C for 1 d.
31 Compounds **1**, **2** and 2,4-dichlorophenoxyacetic acid (2,4-D, positive control) were dissolved
32 in 1 mL of dichloromethane (1, 10, 100 and 1000 nmol/mL) and then poured on filter paper (φ
33 55 mm) in a Petri dish (φ 60 × 20 mm). After the sample-loaded paper had been air-dried, 1
34 mL of distilled water was poured on the sample-loaded paper or intact filter paper (control).
35 The preincubated lettuces (n = 23 in each Petri dish) were transferred onto the sample-loaded
36 filter paper or control filter paper and incubated in a growth chamber in the dark at 23 °C for 3
37 d. The lengths of the hypocotyl and the root were measured using a ruler.

38 Effect of the compounds on the growth of lettuce was examined (Fig. 2). 2,4-D inhibited
39 the root and hypocotyl growth of lettuce dose-dependently. Compound **1** promoted the root
40 growth as low as 100 nmol/paper dose-dependently and inhibited the hypocotyl growth at
41 1000 nmol/paper. Compound **2** inhibited the root and hypocotyl growth at 1000 nmol/paper.

42 We isolated (8*E*, 12*Z*)-10,11-dihydroxyoctadeca-8,12-dienoic acid (**1**) from *L.*
43 *extremiorientale*. Although compound **1** was a known compound, this was the first time to
44 isolate from mushroom. In the previous reports, (12*Z*,15*Z*)-9-hydroxy-10-oxo-octadeca-
45 12,15-dienoic acid (KODA) was isolated as a stress-mediated compound from *Lemna*
46 *paucicostata*. The reaction products of KODA with norepinephrine or epinephrine showed
47 strong flower-inducing activities toward *Lemna* (Yokoyama et al. 2000, 2009; Murata et al.
48 2012). As this example shows, oxidized fatty acids show various activities towards plants.

1 Although leccinine A (2) has been isolated from this mushroom, this is the first report of
2 biological activity of the compound towards plant growth. The taxon-specific germination-
3 inducing factor (GIF) was purified from the ectomycorrhizal fungus *Leccinum aurantiacum*
4 (Bjurman et al. 1984). Mushrooms belonging to the genus *Leccinum* may be good sources of
5 plant growth regulators. Our result will provide useful information for the development of
6 plant-growth regulators.

7 8 Disclosure

9
10 The authors declare no conflict of interest. All the experiments undertaken in this study
11 comply with the current laws of Japan.

12 13 REFERENCES

- 14
15 Asai T, Choi JH, Ikka T, Fushimi K, Abe N, Tanaka H, Yamakawa Y, Kobori H, Kiriiwa Y,
16 Motohashi R, Deo VP, Asakawa T, Kan T, Morita A, Kawagishi H, 2015. Effect of 2-
17 azahypoxanthine (AHX) produced by the fairy-ring-forming fungus on the growth and
18 the grain yield of rice. *Japan Agricultural Research Quarterly* 49: 45–49;
19 <http://dx.doi.org/10.6090/jarq.49.45>.
- 20 Bjurman J, Fries N, 1984. Purification and properties of the germination-inducing factor in
21 the ectomycorrhizal fungus *Leccinum aurantiacum* (Boletaceae). *Physiologia Plantarum*
22 62: 465–471; <http://dx.doi.org/10.1111/j.1399-3054.1984.tb04604.x>.
- 23 Choi JH, Abe N, Tanaka H, Fushimi K, Nishina Y, Morita A, Kiriiwa Y, Motohashi R,
24 Hashizume D, Koshino H, Kawagishi H, 2010a. Plant-growth regulator, imidazole-4-
25 carboxamide produced by fairy-ring forming fungus *Lepista sordida*. *Journal of*
26 *Agricultural and Food Chemistry* 58: 9956–9959; <http://dx.doi.org/10.1021/jf101619a>.
- 27 Choi JH, Fushimi K, Abe N, Tanaka H, Maeda S, Morita A, Hara M, Motohashi R,
28 Matsunaga J, Eguchi Y, Ishigaki N, Hashizume D, Koshino H, Kawagishi H, 2010b.
29 Disclosure of the “fairy” of fairy-ring forming fungus *Lepista sordida*. *ChemBioChem* 11:
30 1373–1377; <http://dx.doi.org/10.1002/cbic.201000112>.
- 31 Choi JH, Ozawa N, Masuda K, Koyama T, Yazawa K, Kawagishi H, 2010c. Suppressing the
32 formation of osteoclasts using bioactive components of the edible mushroom *Leccinum*
33 *extremiorientale* (L. Vass.) Singer (Agaricomycetideae). *International Journal of*
34 *Medicinal Mushrooms* 12: 401–406; <http://dx.doi.org/10.1615/IntJMedMushr.v12.i4.70>.
- 35 Choi JH, Ozawa N, Yamakawa Y, Nagai K, Hirai H, Kawagishi H, 2011. Leccinine A, an
36 endoplasmic reticulum stress-suppressive compound from the edible mushroom
37 *Leccinum extremiorientale*. *Tetrahedron* 67: 6649–6653;
38 <http://dx.doi.org/10.1016/j.tet.2011.05.052>.
- 39 Fushimi K, Anzai K, Tokuyama S, Kiriiwa Y, Matsumoto N, Sekiya A, Hashizume D,
40 Nagasawa K, Hirai H, Kawagishi H, 2012. Agrocybynes A–E from the culture broth
41 *Agrocybe praecox*. *Tetrahedron* 68: 1262–1265;
42 <http://dx.doi.org/10.1016/j.tet.2011.11.049>.
- 43 Hellwig V, Dasenbrock J, Graf C, Kahner L, Schumann S, Steglich W, 2002. Calopins and
44 cyclocalopins-bitter principles from *Boletus calopus* and related mushrooms. *European*
45 *Journal of Organic Chemistry* 17: 2895–2904; [http://dx.doi.org/10.1002/1099-0690\(200209\)2002:17<2895::AID-EJOC2895>3.0.CO;2-S](http://dx.doi.org/10.1002/1099-0690(200209)2002:17<2895::AID-EJOC2895>3.0.CO;2-S).
- 46 Kim JW, Yoo ID, Kim WG, 2006. Free radical-scavenging δ -lactones from *Boletus calopus*.
47 *Planta Medica* 72: 1431–1432; <http://dx.doi.org/10.1055/s-2006-951722>.
- 48 Kovganko NV, Kashkan ZhN, 1999. Sterol glycosides and acylglycosides. *Chemistry of*
49 *Natural Compounds* 35: 479–497; <http://dx.doi.org/10.1007/BF02323277>.
- 50

- 1 Kukovincts OS, Zainullin RA, Kislitsyn MI, 2006. Natural arylterpenes and their biological
2 activity. *Chemistry of Natural Compounds* 42: 1–15; [http://dx.doi.org/10.1007/s10600-](http://dx.doi.org/10.1007/s10600-006-0025-6)
3 006-0025-6.
- 4 Lopez S, Bastida J, Viadomat F, Codina C, 2002. Acetylcholinesterase inhibitory activity of
5 some amaryllidaceae alkaloids and *Narcissus* extracts. *Life Sciences* 71: 2521–2529;
6 [http://dx.doi.org/10.1016/S0024-3205\(02\)02034-9](http://dx.doi.org/10.1016/S0024-3205(02)02034-9).
- 7 Matsuzaki N, Wu J, Kawaide M, Choi JH, Hirai H, Kawagishi H, 2016. Plant growth
8 regulatory compounds from the mushroom *Russula vinosa*. *Mycoscience* 57: 404–407;
9 <http://dx.doi.org/10.1016/j.myc.2016.07.001>.
- 10 Murata A, Kai K, Tsutsui K, Takeuchi J, Todoroki Y, Furihata K, Yokoyama M, Baldermann S,
11 Watanabe N, 2012. Enantio-selective reduction of the flowering related compound
12 KODA and its analogues in *Pharbitis nil* cv. Violet. *Tetrahedron* 68: 5582–5589;
13 <http://dx.doi.org/10.1016/j.tet.2012.04.077>.
- 14 Qiao MF, Ji NY, Miao FP, Yin XL, 2011. Steroids and an oxylipin from an algicolous isolate
15 of *Aspergillus flavus*. *Magnetic Resonance in Chemistry* 49: 366–369;
16 <http://dx.doi.org/10.1002/mrc.2748>.
- 17 Tobina H, Choi JH, Asai T, Kiriiwa Y, Asakawa T, Kan T, Morita A, Kawagishi H, 2014.
18 2-Azahypoxanthine and imidazole-4-carboxamide produced by the fairy-ring-forming
19 fungus increase yields of wheat. *Field Crops Research* 162: 6–11.
- 20 Wu J, Tokunaga T, Kondo M, Ishigami K, Tokuyama S, Suzuki T, Choi JH, Hirai H,
21 Kawagishi H, 2015. Erinaceolactones A to C, from the culture broth of *Hericium*
22 *erinaceus*. *Journal of Natural Products* 78: 155–158;
23 <http://dx.doi.org/10.1021/np500623s>.
- 24 Yokoyama M, Yamaguchi S, Inomata S, Komatsu K, Yoshida S, Iida T, Yokokawa Y,
25 Yamaguchi M, Kaihara S, Takimoto A, 2000. Stress-induced factor involved in flower
26 formation of *Lemna* is an α -ketol derivative of linolenic acid. *Plant & Cell Physiology*
27 41: 110–113; <http://dx.doi.org/10.1093/pcp/41.1.110>.
- 28 Yokoyama M, Yamaguchi S, Suzuki M, Tanaka O, Kobayashi K, Yanaki T, Watanabe N, 2009.
29 Identification of reaction products of 9-hydroxy-10-oxo-12(Z),15(Z)-octadecadienoic
30 acid (KODA) and norepinephrine that strongly induce flowering in *Lemna*. *Plant*
31 *Biotechnology* 25: 547–551; <http://dx.doi.org/10.5511/plantbiotechnology.25.547>.

33 **Figure legends**

34 Fig. 1– Structures of compounds **1** and **2**.

35 Fig. 2– Effect of compounds **1** and **2** on the growth of lettuce. Lettuce seedlings were treated
36 with compounds **1** and **2**. Respective length of growth compared with the control \pm standard
37 deviation (* $p < 0.01$ vs control, $n = 21$ –23).

Highlights

- Two compounds were isolated from the fruiting bodies of *Leccinum extremiorientale*.
- This is the first reported isolation of compound **1** from mushroom.
- Compound **1** strongly promoted the root growth of lettuce.
- Compound **2** strongly inhibited the root and hypocotyl growth of lettuce.
- The biological activity of the compounds against plant was first found in this study.

Figure

[Click here to download high resolution image](#)





Fig. 1 Ito *et al.*

Figure

[Click here to download high resolution image](#)




Fig. 2 Ito *et al.*