

## -L-Fucosidase from *Bombyx mori* has broad substrate specificity and hydrolyzes core fucosylated N-glycans

著者	Nakamura Shuntaro, Miyazaki Takatsugu, Park Enoch Y.
journal or publication title	Insect Biochemistry and Molecular Biology
page range	103427
year	2020-09
出版者	Elsevier
権利	(C) 2020. This manuscript version is made available under the CC-BY-NC-ND 4.0 license <a href="http://creativecommons.org/licenses/by-nc-nd/4.0/">http://creativecommons.org/licenses/by-nc-nd/4.0/</a>
URL	<a href="http://hdl.handle.net/10297/00027529">http://hdl.handle.net/10297/00027529</a>

doi: 10.1016/j.ibmb.2020.103427

1  **$\alpha$ -L-Fucosidase from *Bombyx mori* has broad substrate specificity and**  
2 **hydrolyzes core fucosylated *N*-glycans**

3

4 Shuntaro Nakamura<sup>1</sup>, Takatsugu Miyazaki<sup>1,2,\*</sup> and Enoch Y. Park<sup>1,2</sup>

5

6 <sup>1</sup>Department of Agriculture, Graduate School of Integrated Science and Technology,

7 Shizuoka University, 836 Ohya, Suruga-ku, Shizuoka, 422-8529, Japan.

8

9 <sup>2</sup>Green Chemistry Research Division, Research Institute of Green Science and Technology,

10 Shizuoka University, 836 Ohya, Suruga-ku, Shizuoka, 422-8529, Japan.

11

12 **\*Correspondence:** Takatsugu Miyazaki, Green Chemistry Research Division, Research

13 Institute of Green Science and Technology, Shizuoka University, 836 Ohya, Suruga-ku,

14 Shizuoka, 422-8529, Japan; Tel.: +81-54-238-4886; E-mail:

15 miyazaki.takatsugu@shizuoka.ac.jp

16

17

18

19 **Abstract**

20 *N*-glycans play a role in physiological functions, including glycoprotein conformation,  
21 signal transduction, and antigenicity. Insects display both  $\alpha$ -1,6- and  $\alpha$ -1,3-linked fucose  
22 residues bound to the innermost *N*-acetylglucosamine of *N*-glycans whereas core  $\alpha$ -1,3-  
23 fucosylated *N*-glycans are not found in mammals. Functions of insect core-fucosylated glycans  
24 are not clear, and no  $\alpha$ -L-fucosidase related to the *N*-glycan degradation has been identified. In  
25 the genome of the domestic silkworm, *Bombyx mori*, a gene for a protein, BmFucA, belonging  
26 to the glycoside hydrolase family 29 is a candidate for an  $\alpha$ -L-fucosidase gene. In this study,  
27 BmFucA was cloned and recombinantly expressed as a glutathione-*S*-transferase tagged protein  
28 (GST-BmFucA). Recombinant GST-BmFucA exhibited broad substrate specificity and  
29 hydrolyzed *p*-nitrophenyl  $\alpha$ -L-fucopyranoside, 2'-fucosyllactose, 3-fucosyllactose, 3-fucosyl-  
30 *N,N'*-diacetylchitobiose, and 6-fucosyl-*N,N'*-diacetylchitobiose. Further, GST-BmFucA  
31 released fucose from both pyridylaminated complex-type and paucimannose-type glycans that  
32 were core- $\alpha$ -1,6-fucosylated. GST-BmFucA also shows hydrolysis activity for core-fucosylated  
33 glycans attached to phospholipase A<sub>2</sub> from bee venom. BmFucA may be involved in the  
34 catabolism of core-fucosylated *N*-glycans in *B. mori*.

35

36 **Keywords:** *Bombyx mori* / fucose / glycoprotein / glycoside hydrolase family 29 / *N*-glycan

37

38 **Abbreviations:** ER, endoplasmic reticulum; Glc, glucose; GlcNAc, *N*-acetylglucosamine; Fuc,  
39 fucose; Man, mannose; GH29, Glycoside hydrolase family 29; FucA,  $\alpha$ -L-fucosidase; FucT,  
40 fucosyltransferase; 2'-FL, 2'-fucosyllactose; 3-FL, 3-fucosyllactose; 6-FGn<sub>2</sub>, GlcNAc $\beta$ (1-  
41 4)[Fuc $\alpha$ (1-6)]GlcNAc; 3-FGn<sub>2</sub>, GlcNAc $\beta$ (1-4)[Fuc $\alpha$ (1-3)]GlcNAc; PA, pyridylamino;  
42 PNGase F, peptide:*N*-glycanase F; pNP, *p*-nitrophenyl; PLA<sub>2</sub>, Phospholipase A<sub>2</sub>; AAL, *Aleuria*  
43 *aurantia* lectin

44

## 45 **1. Introduction**

46 Protein glycosylation is a common post-translational modification (Varki, 2017).  
47 Glycoproteins have various glycan structures that affect various physiological functions, such  
48 as protein folding, stability, transport, and activity (Moremen et al., 2012; Ohtsubo and Marth,  
49 2006; Varki et al., 2009b). *N*-glycans are attached to an asparagine residue at the Asn-X-Ser/Thr  
50 sequon of glycoproteins, where X is any amino acid residue except Pro, and preferably at an  
51 Asn-X-Ser sequon rather than an Asn-X-Thr sequon (Reddy et al., 1999; Zielinska et al., 2010).  
52 *N*-glycans are composed mainly of fucose (Fuc), galactose, glucose (Glc), *N*-acetylglucosamine  
53 (GlcNAc), *N*-acetylgalactosamine, mannose (Man), and sialic acid (Moremen et al., 2012).

54 *N*-glycan structure differs between insects and mammals. Mammalian cells produce  
55 complex-type *N*-glycans, and insect cells mainly produce paucimannose-type *N*-glycans.  
56 Biosynthesis of mammalian complex-type *N*-glycans involves an *N*-glycan precursor  
57 (Glc<sub>3</sub>Man<sub>9</sub>GlcNAc<sub>2</sub>) first synthesized as a lipid-linked oligosaccharide in the endoplasmic  
58 reticulum (ER). This precursor is transferred to an asparagine residue in the sequon and is  
59 trimmed and elongated by enzymes located in the ER and Golgi apparatus.  
60 GlcNAcMan<sub>3</sub>GlcNAc<sub>2</sub>, an intermediate in *N*-glycan synthesis, is extended by Golgi-resident  
61 glycosyltransferases (Moremen et al., 1994). Insects display similar *N*-glycan biosynthesis; a  
62 GlcNAcMan<sub>3</sub>GlcNAc<sub>2</sub> glycan intermediate is also formed cultured cells. Insect cells generally  
63 show low activity for  $\beta$ -1,2-*N*-acetylglucosaminyltransferase II (GnT-II) and  
64 galactosyltransferase and no detectable activity for sialyltransferase (Altman et al., 1993;  
65 Geisler and Jarvis, 2012; van Die et al., 1996; Walski et al., 2017b). Also, insect cells possess  
66 *N*-acetylglucosaminidase (fused lobes) that removes terminal GlcNAc residues from  
67 GlcNAcMan<sub>3</sub>GlcNAc<sub>2</sub>, a process required to synthesize paucimannose-type *N*-glycans (Geisler  
68 and Jarvis, 2008). Homologs of mammalian glycosyltransferases involved in complex glycan  
69 biosynthesis have been identified in insects (Geisler and Jarvis, 2012; Haines and Irvine, 2005;  
70 Kajiura et al., 2015; Koles et al., 2004; Miyazaki et al., 2019a; Miyazaki et al., 2019b; Vadaie  
71 and Jarvis, 2004), and complex glycans other than paucimannose-type are found in *N*-glycomes  
72 of some insects and their culture cells (Hykollari et al., 2019; Koles et al., 2007; Kurz et al.,  
73 2015; Stanton et al., 2017). Sialylated *N*-glycans were detected in some insect cells and tissues  
74 (Aoki et al., 2007; Aoki and Tiemeyer, 2010; Cime-Castillo et al., 2015; Walski et al., 2017a)  
75 and in the envelop protein of mature dengue type 2 virus derived from insect cells (Lei et al.,  
76 2015). Thus, the *N*-glycan pathway in insects is complicated and is not completely understood.

77 Some insect *N*-glycans can be further processed with  $\alpha$ -1,3- and/or  $\alpha$ -1,6-linked fucose  
78 residues on the innermost GlcNAc residue of the core structure.  $\alpha$ -1,6-Fucosylation is catalyzed  
79 by  $\alpha$ -1,6-fucosyltransferase (FucT6), and core  $\alpha$ -1,6-fucosylated *N*-glycans are further  
80 processed by core  $\alpha$ -1,3-fucosyltransferase (FucTA) to produce core  $\alpha$ -1,6- and  $\alpha$ -1,3-  
81 fucosylated glycans unique to insects and nematodes but absent in mammals (Aoki et al., 2007;  
82 Fabini et al., 2001; Minagawa et al., 2015; Yan et al., 2013). FucTA fucosylates non- $\alpha$ -1,6-  
83 fucosylated glycans, but FucT6 cannot transfer fucose to core  $\alpha$ -1,3-fucosylated glycans  
84 (Paschinger et al., 2005). FucT6 and FucTA require a GlcNAc residue on the non-reducing end  
85 of *N*-glycans (Fabini et al., 2011; Paschinger et al., 2005). Analyses of *N*-glycan structures  
86 revealed that the core-fucosylation is universal among insects, including beetles (Walski et al.,  
87 2016; Liu et al., 2019), moths (Kajiura et al., 2015; Mabashi-Asazuma et al., 2015; Soya et al.,  
88 2016; Stanton et al., 2017), and bees (Kubelka et al., 1993; Kubelka et al., 1995). The levels of  
89 core fucosylation were reported to be different between males and females of invertebrates  
90 including insects (Scheys et al., 2019; Scheys et al., 2020). Core  $\alpha$ -1,6-fucosylated glycans in  
91 mammals affect tissue development and signal transduction, but the role of insect core-  
92 fucosylated glycans is less clear (Becker and Lowe, 2003; Miyoshi et al., 2008; Walski et al.,  
93 2017b). Further, enzymes related to the degradation of insect *N*-glycans are poorly  
94 characterized.

95  $\alpha$ -L-Fucosidase is classified as a member of the glycoside hydrolase family (GH) 29, 95,  
96 141, and 151 according to the CAZy database (<http://www.cazy.org/>, Katayama et al., 2004;  
97 Lombard et al., 2014; Ndeh et al., 2017; Sela et al., 2012) GH29 and GH95 account for most  
98  $\alpha$ -L-fucosidases currently registered. Hydrolysis mechanisms of GH29 and GH95 enzymes are  
99 retaining and inverting, respectively (Nagae et al., 2007; Sulzenbacher et al., 2004). GH29 is  
100 comprised of  $\alpha$ -L-fucosidases from archaea, bacteria, fungi, plants, and mammals, whereas  
101 GH95  $\alpha$ -L-fucosidases are found in bacteria, fungi, and plants but not in animals (Intra et al.,  
102 2006; Pogorelko et al., 2016). Structural differences in catalytic domains are observed; GH29  
103 enzymes adopt a  $(\beta/\alpha)_8$ -barrel architecture, and domains of GH95 enzymes show an  $(\alpha/\alpha)_6$ -  
104 barrel fold (Nagae et al., 2007; Sakurama et al., 2012). GH29  $\alpha$ -L-fucosidases are divided into  
105 two subfamilies, GH29-A and GH29-B, based on differences in substrate specificity (Grootaert  
106 et al., 2020; Sakurama et al., 2012). GH29-B enzymes are more specific for  $\alpha$ -1,3/4-fucosylated  
107 substrates than GH29-A enzymes. GH29-A  $\alpha$ -L-fucosidases have been isolated and identified  
108 from a wide range of organisms, including bacteria (Intra et al., 2007).

109 The physiological function of two arthropodal GH29  $\alpha$ -L-fucosidases has been reported.  
110 *Drosophila melanogaster*  $\alpha$ -L-fucosidase (DmFucA) is expressed in testis, accessory gland,  
111 midgut, and brain (Pasini et al., 2008). DmFucA on the sperm plasma membrane participates,  
112 during fertilization, in gametophyte recognition possibly through interaction with glycoside  
113 ligands on egg surfaces (Intra et al., 2006). In contrast, an  $\alpha$ -L-fucosidase from the spider,  
114 *Nephilingis cruentata* (NcFuc), is expressed in midgut diverticula (Perrella et al., 2018a).  
115 NcFuc is associated with digestive processes and likely removes fucose residues from food  
116 sources for metabolic purposes (Fuzita et al., 2016; Perrella et al., 2018a) Physiological  
117 functions and localization of arthropodal  $\alpha$ -L-fucosidases are reported, but substrate specificity  
118 of insect  $\alpha$ -L-fucosidases and their relationship with *N*-glycan metabolism remain unknown. In  
119 this study, GH29  $\alpha$ -L-fucosidase (named BmFucA) from *Bombyx mori* is first cloned, deeply  
120 investigated for substrate specificity, and identified as an enzyme that hydrolyzes both the core  
121  $\alpha$ -1,3- and  $\alpha$ -1,6-fucoside linkages of *N*-glycans.

122  
123

## 124 **2. Materials and methods**

### 125 **2.1. Chemical**

126 *p*-Nitrophenyl  $\alpha$ -L-fucopyranoside (pNP- $\alpha$ -L-Fuc) was purchased from Merck (Darmstadt,  
127 Germany), 2'-fucosyllactose (2'-FL) and 3-fucosyllactose (3-FL) were from Carbosynth  
128 Limited (Berkshire, UK), and GlcNAc $\beta$ (1-4)[Fuc $\alpha$ (1-6)]GlcNAc (6-FGn<sub>2</sub>) and GlcNAc $\beta$ (1-  
129 4)[Fuc $\alpha$ (1-3)]GlcNAc (3-FGn<sub>2</sub>) were from Tokyo Chemical Industry (Tokyo, Japan).  
130 Fucosylated substrate structures were described in Supplementary data, Fig. S1. The  
131 pyridylaminated (PA) glycans were purchased from Masuda Chemical Industries (Takamatsu,  
132 Japan). All other reagents were of analytical grade and purchased from FUJIFILM Wako Pure  
133 Chemical (Osaka, Japan) or Merck unless otherwise stated.

134

### 135 **2.2. Cloning, expression, and purification of BmFucA**

136 Total RNA of a *B. mori* fifth-instar larva was extracted using Trizol reagent (Thermo Fisher  
137 Scientific, Waltham, MA, USA) and cDNA synthesized using a PrimeScript RT reagent kit  
138 (Takara Bio, Kusatsu, Japan). DNA encoding BmFucA without the signal sequence (Met1–  
139 Gly15), predicted using SignalP (<http://www.cbs.dtu.dk/services/SignalP/>, Petersen et al.,  
140 2011) as amplified by PCR using cDNA as a template and a set of primers, 5'-  
141 TTTCCATGGCCATACACAATCGTCGTGATC-3' and 5'-  
142 TTTTCTCGAGTTATGTGGTCGCGAGTTTGAGG-3'. Amplified DNA was subcloned into

143 a pET41a (+) vector (Merck) at NcoI and XhoI restriction enzyme sites, and *E. coli* BL21 (DE3)  
144 was transformed with the plasmid to generate a recombinant protein containing N-terminal  
145 glutathione *S*-transferase- and His-tags (GST-BmFucA). *E. coli* BL21(DE3) harboring the  
146 expression plasmid was incubated at 37°C in Luria–Bertani medium containing 50 µg/mL  
147 kanamycin, until optical density reached 0.6–0.8. At this point, enzyme synthesis was induced  
148 with 0.1 mM isopropyl β-D-1-thiogalactopyranoside for 24 h at 20°C. The cells were harvested  
149 by centrifugation (5,000 × g, 4°C, 5 min) and resuspended in 50 mM sodium phosphate buffer  
150 (pH 7.5) containing 300 mM NaCl and 20 mM imidazole. Cells were disrupted by sonication  
151 on ice for 20 min. After centrifugation (20,640 × g, 4°C, 15 min) to remove the insoluble  
152 fraction, the supernatant was loaded onto a Ni-Sepharose excel column (GE Healthcare,  
153 Chicago, USA) pre-equilibrated with the same buffer. The column was washed with buffer, and  
154 recombinant protein was then eluted with 50 mM sodium phosphate buffer (pH 7.5) containing  
155 300 mM NaCl and 250 mM imidazole. Protein purity was confirmed by SDS-PAGE and  
156 Coomassie brilliant blue staining. Fractions containing protein were dialyzed against 20 mM  
157 sodium citrate buffer (pH 5.5) containing 300 mM NaCl. GST-BmFucA concentration was  
158 determined by absorbance at 280 nm based on a molar absorptivity calculated using ExPASy  
159 ProtParam (<http://web.expasy.org/protparam/>).

160

### 161 **2.3. Enzyme assay**

162 Enzyme activity for pNP-α-L-Fuc was measured in 50 µL reaction mixtures containing  
163 1 µg/mL of GST-BmFucA and 1 mM pNP-α-L-Fuc. Reactions were stopped by two volumes  
164 of 1 M Na<sub>2</sub>CO<sub>3</sub>, and liberated *p*-nitrophenol was quantified by absorbance at 405 nm. pH  
165 dependence was investigated at 30°C in 50 mM sodium citrate buffer (pH 3.0–8.0), 50 mM  
166 MES–NaOH buffer (pH 6.0–6.5), and 50 mM sodium phosphate buffer (pH 6.5–8.0).  
167 Temperature dependence was assayed at various temperatures (25°C–80°C) in 50 mM sodium  
168 citrate buffer (pH 5.0). Kinetic parameters were determined using GST-BmFucA (60 nM) and  
169 various concentrations (0.1–2 mM) of pNP-α-L-Fuc in sodium citrate buffer (pH 5.5) at 30°C.  
170 Kinetic parameters were calculated using non-linear regression analysis with Kaleida Graph  
171 (Synergy Software, Kentucky, USA).

172 When 2'-FL, 3-FL, and 6-FGn<sub>2</sub> were used as substrates, hydrolysis activity of GST-  
173 BmFucA was analyzed by TLC with reaction condition as 10 mM substrates, 95 µg/mL GST-  
174 BmFucA, 30°C, and 17 h. Reaction solution and authentic standards (fucose, lactose, and  
175 chitobiose) as controls were spotted on TLC aluminum sheet silica gel 60 F254 and developed

176 with 1-butanol:acetic acid:water = 2:1:1. To calculate specific activity of GST-BmFucA for 2'-  
177 FL, 3-FL, 6-FGn<sub>2</sub>, and 3-FGn<sub>2</sub>, fucose release was quantified using a K-FUCOSE kit  
178 (Megazyme, Dublin, Ireland). Enzyme reaction volume was 50  $\mu$ L containing 95  $\mu$ g/mL GST-  
179 BmFucA and 1 mM each substrate. Reaction mixtures were incubated at 30°C.

180 To measure hydrolytic activity toward free core  $\alpha$ -1,6-fucosylated *N*-glycans, reaction  
181 mixtures containing 50  $\mu$ g/mL GST-BmFucA, 1  $\mu$ M PA-glycan (MMF<sup>6</sup>-PA, GnGnF<sup>6</sup>-PA, or  
182 NaNaF<sup>6</sup>-PA), and 50 mM sodium citrate buffer (pH 5.0) were prepared and incubated at 30°C.  
183 Reaction products were separated and detected by reverse phase high-performance liquid  
184 chromatography (HPLC) using a TKSgel ODS-80<sub>TM</sub> column (4.6 mm  $\times$  250 mm, Tosho, Tokyo,  
185 Japan) as described previously (Miyazaki et al., 2019a; Miyazaki et al., 2019b).

186

## 187 **2.4. Lectin blotting**

188 Phospholipase A<sub>2</sub> (PLA<sub>2</sub>) from *Apis mellifera* bee venom (0.5 mg/mL, Cayman Chemical  
189 Co., Michigan, USA) was dissolved in sodium citrate buffer (pH 5.0) containing 10 mM  
190 dithiothreitol and denatured at 95°C for 15 min, followed by addition of 1  $\mu$ g of BmFucA and  
191 further incubation at 30°C for 40 h. PLA<sub>2</sub> samples were treated with peptide:*N*-glycanase F  
192 (PNGaseF, Takara Bio, Kusatsu, Japan) at 37°C for 17 h and proteins separated by SDS-PAGE  
193 and electroblotted onto polyvinylidene fluoride membranes with a Mini Trans-Blot  
194 Electrophoretic Transfer Cell (Bio-Rad, Hercules, CA, USA). Lectin blotting used a fluorescein  
195 isothiocyanate-conjugated *Aleuria aurantia* lectin (AAL, Vector Laboratories Inc., Burlingame,  
196 USA) to evaluate defucosylation ratio after  $\alpha$ -L-fucosidase treatment. AAL band intensity was  
197 quantified by ImageJ software, and significant differences were calculated by the Student's t-  
198 test (Schneider et al., 2012). Lectin blotting was performed in triplicate.

199

## 200 **3. Results**

### 201 **3.1. Expression and general properties of recombinant BmFucA**

202 A BLAST search was performed using amino acid sequences of characterized  $\alpha$ -L-  
203 fucosidases to identify an  $\alpha$ -L-fucosidase (BmFucA) gene in the *B. mori* genome. BmFucA was  
204 found and has 51% and 53% sequence identities with *Homo sapiens*  $\alpha$ -L-fucosidase (HsFucA1)  
205 and DmFucA, respectively. The open reading frame of BmFucA encodes 477 amino acid  
206 residues with a predicted signal peptide of 15 amino acids. A phylogenetic analysis exhibited  
207 that BmFucA is classified in subfamily GH29-A (Fig. 1). Recombinant BmFucA without the  
208 signal peptide was successfully expressed in *Escherichia coli* as a GST-BmFucA and purified



209 to homogeneity (Fig. 2A). The molecular weight of GST-BmFucA estimated from SDS-PAGE  
210 analysis is 89 kDa, which was almost identical kDa calculated from its amino acid sequence  
211 (82 kDa).

212 GST-BmFucA exhibited hydrolytic activity for pNP- $\alpha$ -L-Fuc. Optimum pH of GST-  
213 BmFucA was pH 4.5–5.0, and optimum temperature was 45°C–60°C (Fig. 2B, C). Kinetic  
214 parameters were  $K_m = 0.2 \pm 0.02$  mM,  $k_{cat} = 12.5 \pm 0.5$  s<sup>-1</sup>, and  $k_{cat}/K_m = 62.7$  s<sup>-1</sup> mM<sup>-1</sup> (Table  
215 1 and Fig. S2).

216

### 217 **3.2. Specificity of GST-BmFucA toward fucose-containing oligosaccharides**

218 Commercially available fucose-containing oligosaccharides, 2'-FL, 3-FL, 6-FGn<sub>2</sub>, and 3-  
219 FGn<sub>2</sub>, were tested for hydrolysis by GST-BmFucA. Thin-layer chromatography (TLC) analysis  
220 showed release of fucose from 2'-FL, 3-FL, and 6-FGn<sub>2</sub> after incubation with the enzyme (Fig.  
221 3). All oligosaccharides including 3-FGn<sub>2</sub> were hydrolyzed using a K-FUCOSE kit (see  
222 Materials and Methods). Specific activities of GST-BmFucA for 2'-FL, 3-FL, 6-FGn<sub>2</sub>, and 3-  
223 FGn<sub>2</sub> were  $168.4 \pm 9.9$  nmol min<sup>-1</sup> mg<sup>-1</sup>,  $35.3 \pm 8.7$  nmol min<sup>-1</sup> mg<sup>-1</sup>,  
224  $127.3 \pm 12.2$  nmol min<sup>-1</sup> mg<sup>-1</sup>, and  $19.9 \pm 3.3$  nmol min<sup>-1</sup> mg<sup>-1</sup>, respectively. GST-BmFucA  
225 hydrolysis the showed substrate preferences as 2'-FL > 6-FGn<sub>2</sub> > 3-FL > 3-FGn<sub>2</sub>. Activity  
226 toward these substrates was less than activity of pNP $\alpha$ -L-Fuc (Table 2).

227

### 228 **3.3. Activity of GST-BmFucA toward core-fucosylated glycans**

229 GST-BmFucA was assayed with core  $\alpha$ -1,6-fucosylated fluorescent PA glycans, and the  
230 reaction products were analyzed using reversed-phase HPLC. The retention times of glycan  
231 substrates, MMF<sup>6</sup>-PA, GnGnF<sup>6</sup>-PA, and NaNaF<sup>6</sup>-PA, were 15, 19, and 22 min, respectively  
232 (Fig. 4). Incubation with substrates induced new peaks with shorter retention times, increasing  
233 peak areas with time, indicating that GST-BmFucA hydrolyzed  $\alpha$ -1,6-fucosyl linkages (Fig. 4).  
234 Specific activities with MMF<sup>6</sup>-PA, GnGnF<sup>6</sup>-PA, and NaNaF<sup>6</sup>-PA were 26.9, 24.4, and  
235 26.4 pmol min<sup>-1</sup> mg<sup>-1</sup>, respectively.

236 GST-BmFucA activity on core  $\alpha$ -fucosylated glycoproteins was assessed using lectin  
237 blotting. PLA<sub>2</sub> from *Apis mellifera* bee venom (15.7 kDa), which contains core  $\alpha$ -1,3-  
238 fucosylated and core  $\alpha$ -1,6-fucosylated *N*-glycans, and PNGaseF were used as a model  
239 glycoprotein in the assay (Li et al., 2018). PNGaseF is active on non-fucosylated or core  $\alpha$ -1,6-  
240 fucosylated glycans but not on  $\alpha$ -1,3-fucosylated glycans (Tretter et al., 1991). Using AAL,  
241 which has a high affinity for fucose linked to  $\alpha$ -1,3- and  $\alpha$ -1,6-GlcNAc (Yamashita et al., 1985),

242 PLA<sub>2</sub> was found to be fucosylated (Fig. 4D). Band intensity of AAL for PLA<sub>2</sub> was reduced by  
243 55% after GST-BmFucA treatment. Thus, GST-BmFucA hydrolyzed core  $\alpha$ -1,3- and/or  $\alpha$ -1,6-  
244 fucosylated *N*-glycans of PLA<sub>2</sub>. When PLA<sub>2</sub> was treated with GST-BmFucA and PNGaseF,  
245 band intensity of PLA<sub>2</sub> was reduced by approximately 22% compared with the PNGaseF  
246 treatment alone, suggesting that GST-BmFucA might remove core  $\alpha$ -1,3-fucose from *N*-  
247 glycans of PLA<sub>2</sub>.

248

#### 249 **4. Discussion**

250 In this study, BmFucA was cloned and enzymatically characterized to reveal its substrate  
251 specificity and function. The optimal pH of GST-BmFucA is like that of DmFucA and NcFuc  
252 (Intra et al., 2006; Perrella et al., 2018b) as well as rat and human lysosomal fucosidases  
253 (Dawson and Tsay, 1977; Opheim and Touster, 1977). Kinetic parameters for pNP- $\alpha$ -L-Fuc are  
254 comparable with those of HsFucA1, NcFuc, and TmFuc belonging to GH29-A (Table 1)  
255 (Dawson and Tsay, 1977; Perrella et al., 2018b; Tarling et al., 2003). Moreover, GST-BmFucA  
256 activity differs from GH29-B enzymes because GST-BmFucA efficiently hydrolyzes pNP- $\alpha$ -  
257 L-Fuc compared with GH29-B  $\alpha$ -L-fucosidases from *Bifidobacterium bifidum*, *Bacteroides*  
258 *thetaiotaomicron*, and *Arabidopsis thaliana* (Ashida et al., 2009; Kato et al., 2018; Sakurama  
259 et al., 2012). These latter enzymes reportedly do not act on pNP- $\alpha$ -L-Fuc (Table 1). The broad  
260 specificity of GST-BmFucA for various fucosyl linkages and the greatest activity for pNP- $\alpha$ -  
261 L-Fuc among tested substrates are similar to GH29-A  $\alpha$ -L-fucosidases.

262 2'-FL and 3-FL, found in human milk, are good substrates for GH95  $\alpha$ -L-fucosidase and  
263 GH29-B  $\alpha$ -1,3/4-fucosidase from bifidobacteria, respectively (Ashida et al., 2009; Sakurama et  
264 al., 2012; Sela et al., 2012). These bifidobacterial  $\alpha$ -L-fucosidases seem to release fucose from  
265 fucose-containing oligosaccharides for subsequent use in cellular metabolism. *Elizabethkingia*  
266 *meningoseptica*  $\alpha$ -L-fucosidase cFase I, whose amino acid sequence identity to BmFucA is  
267 25%, has high activity for the core  $\alpha$ -1,3-fucosyl linkage in *N*-glycans as well as for 3-FL (Li  
268 et al., 2017). To date, oligosaccharides having 2'-FL and 3-FL components have not been  
269 identified in *B. mori* and other insects yet. Further, few reports are available of  $\alpha$ -L-fucosidases  
270 exhibiting hydrolytic activity toward trisaccharides 6-FGn<sub>2</sub> and 3-FGn<sub>2</sub>, though some studies  
271 report that disaccharides Fuca-1,6-GlcNAc and Fuca-1,3-GlcNAc are substrates. *Lactobacillus*  
272 *casei*  $\alpha$ -L-fucosidases, AlfC and AlfB, show high activity for Fuca-1,6-GlcNAc and Fuca-1,3-  
273 GlcNAc, respectively. They also release core fucose from *N*-glycans contained in human  
274 mucosal surfaces and breast milk. *L. casei* may utilize this fucose (Becerra et al., 2020;

275 Rodríguez-Díaz et al., 2011). GST-BmFucA activity on trisaccharides, 6-FGn<sub>2</sub> and 3-FGn<sub>2</sub>, was  
276 lower than AlfB and AlfC activity toward disaccharides. HsFucA1, located in lysosomes, also  
277 hydrolyzes Fuc $\alpha$ -1,6-GlcNAc. HsFucA1 also has broad substrate specificity for 2'-FL and  
278 oligosaccharides containing fucose with  $\alpha$ -1,3- and  $\alpha$ -1,4-linkages (Dawson and Tsay, 1977).  
279 GST-BmFucA shows similar properties as HsFucA1 on various linked fucosyloligosaccharides  
280 (Table 2).

281 To date, few studies of  $\alpha$ -L-fucosidase using core-fucosylated *N*-glycans as substrates are  
282 available. *Bacteroides fragilis*  $\alpha$ -L-fucosidase (BfFucH) was used with fluorescent 2-  
283 aminobenzamide-labeled core  $\alpha$ -1,6-fucosylated *N*-glycans. BfFucH releases fucose from  
284 paucimannose and complex *N*-glycans, but it removed only 26%–35% of fucose in 1 h.  
285 *Omnitrophica bacterium*  $\alpha$ -L-fucosidase releases fucose from core  $\alpha$ -1,3- and  $\alpha$ -1,6-fucosylated  
286 glycans, respectively (Tsai et al., 2017; Vainauskas et al., 2018). In contrast,  $\alpha$ -L-fucosidase  
287 from *Streptomyces* sp. 142 did not release fucose from core  $\alpha$ -1,6-fucosylated complex *N*-  
288 glycans (Sano et al., 1992). Compared with BfFucH activity, GST-BmFucA displays higher  
289 activity on core  $\alpha$ -1,6-fucosylated *N*-glycans. The present study indicates that the nonreducing  
290 end structure of *N*-glycans does not affect GST-BmFucA activity. GST-BmFucA represents the  
291 first case of activity of an arthropodal  $\alpha$ -L-fucosidase for core-fucosylated *N*-glycans. PLA<sub>2</sub> has  
292 a single sequon (Asn13-Lys14-Ser15) with  $\alpha$ -1,6 or  $\alpha$ -1,3 core fucose (Staudacher et al., 1992).  
293 As described above, insects also exhibit core- $\alpha$ -1,3-fucosylated glycans (Fabini et al., 2001;  
294 Kajihura et al., 2015; Kubelka et al., 1993, Kubelka et al., 1995; Liu et al., 2019; Mabashi-  
295 Asazuma et al., 2015; Minagawa et al., 2015; Soya et al., 2016; Stanton et al., 2017; Walski et  
296 al., 2016). Tomato and *Arabidopsis*  $\alpha$ -fucosidases classified as GH29-B are involved in *N*-  
297 glycan degradation and hydrolyze  $\alpha$ -1,3-fucosyl linkages of 3-FGn<sub>2</sub> but not 3-fucosyl-*N*-  
298 acetylglucosamine and longer core- $\alpha$ -1,3-fucosyl *N*-glycan substrates (Kato et al., 2018;  
299 Rahman et al., 2017; Rahman et al., 2018). In addition, almond  $\alpha$ -L-fucosidase can hydrolyze  
300 3-FL and lacto-*N*-fucopentanose II but not showed activity core- $\alpha$ -1,3-fucosylated *N*-glycans  
301 (Zeleny et al., 2006). Since GST-BmFucA exhibited hydrolytic activity on core- $\alpha$ -1,6-  
302 fucosylated *N*-glycans and might act on core- $\alpha$ -1,3-fucosylated *N*-glycans (Fig. 4), BmFucA is  
303 likely to be involved in the catabolism of core-fucosylated *N*-glycans in *B. mori* cells.  $\alpha$ -L-  
304 Fucosidase activity on both core  $\alpha$ -1,3- and  $\alpha$ -1,6-fucosylated *N*-glycans is reported in bacteria  
305 (Vainauskas et al., 2018) but not for eukaryotic  $\alpha$ -L-fucosidases.

306 Catabolism of *N*-glycan of glycoproteins occurs mainly in lysosomes. pH in lysosomes is  
307 low, which may partially denature glycoproteins (Winchester, 2005). Glycoproteins are

308 degraded by various proteases and digested into polypeptides. *N*-glycans bound to polypeptides  
309 are cleaved with glycosylasparaginase to release the sugar chain (Aronson, 1999).  $\alpha$ -L-  
310 Fucosidase,  $\alpha$ -mannosidase, and  $\beta$ -*N*-acetylglucosaminidase act on released sugar chains to  
311 produce monosaccharides. These sugars are then used for re-synthesis of new *N*-glycans  
312 (Abraham et al., 1983; Barker et al., 1988; Baussant et al., 1986; Johnson and Alhadeff, 1991;  
313 Kuranda and Aronson, 1986). Considering the pH dependence of GST-BmFucA together with  
314 the above discussion, BmFucA is proposed to act on free *N*-glycans in lysosomes and release  
315 fucose for *N*-glycan turnover. This study provides new insights into carbohydrate metabolism  
316 in insects. Also, BmFucA may be used in glycoengineering to remove fucose from core-  
317 fucosylated *N*-glycans of glycoproteins, such as immunoglobulins, that are expressed in both  
318 insect and mammalian cells.

319

### 320 **Acknowledgment**

321 This work was supported in part by Japan Society for the Promotion of Science KAKENHI  
322 (grant No. 19K15748). We thank Enago ([www.enago.jp](http://www.enago.jp)) for the English language review.

323

324 **References**

325 Abraham, D, Blakemore, W.F., Jolly, R.D., Sidebotham, R., Winchester, B., 1983. The  
326 catabolism of mammalian glycoproteins. Comparison of the storage products in bovine, feline  
327 and human mannosidosis. *Biochem. J.* 215, 573–579.

328  
329 Altmann, F., Kornfeld, G., Dalik, T., Staudacher, E., Glössl, J., 1993. Processing of asparagine-  
330 linked oligosaccharides in insect cells. *N*-acetylglucosaminyltransferase I and II activities in  
331 cultured lepidopteran cells. *Glycobiology* 3, 619–625.

332  
333 Aoki, K., Perlman, M., Lim, J.M., Cantu, R., Wells, L., Tiemeyer, M., 2007. Dynamic  
334 developmental elaboration of *N*-linked glycan complexity in the *Drosophila melanogaster*  
335 embryo. *J. Biol. Chem.* 282, 9127–9142.

336  
337 Aoki, K., Tiemeyer, M., 2010. The glycomics of glycan glucuronylation in *Drosophila*  
338 *melanogaster*. *Methods Enzymol.* 480, 297–321.

339  
340 Aronson, N.N. Jr., 1999. Aspartylglycosaminuria: biochemistry and molecular biology.  
341 *Biochim. Biophys. Acta.* 1455, 139–154.

342  
343 Ashida, H., Miyake, A., Kiyohara, M., Wada, J., Yoshida, E., Kumagai, H., Katayama, T.,  
344 Yamamoto, K., 2009. Two distinct  $\alpha$ -L-fucosidases from *Bifidobacterium bifidum* are essential  
345 for the utilization of fucosylated milk oligosaccharides and glycoconjugates. *Glycobiology* 9,  
346 1010–1017.

347  
348 Ashida, H., Fujimoto, T., Kurihara, S., Nakamura, M., Komeno, M., Huang, Y., Katayama, T.,  
349 Kinoshita, T., Takegawa, K., 2020. 1,6- $\alpha$ -L-Fucosidases from *Bifidobacterium longum* subsp.  
350 *infantis* ATCC 15697 involved in the degradation of core-fucosylated *N*-Glycan. *J. App.*  
351 *Glycosci.* 67, 23–29.

352  
353 Barker, C., Dell, A., Rogers, M., Alhadeff, J.A., Winchester, B., 1988. Canine  $\alpha$ -L-fucosidase  
354 in relation to the enzymic defect and storage products in canine fucosidosis. *Biochem. J.* 254,  
355 861–868

356

357 Baussant, T., Strecker, G., Wieruszeski, J.M., Montreuil, J., Michalski, J.C., 1986. Catabolism  
358 of glycoprotein glycans. Characterization of a lysosomal endo-*N*-acetyl- $\beta$ -D-glucosaminidase  
359 specific for glycans with a terminal chitobiose residue. *Eur. J. Biochem.* 159, 381–385  
360

361 Becerra, J.E., Rodríguez-Díaz, J., Gozalbo-Rovira, R., Palomino-Schätzlein, M., Zúñiga, M.,  
362 Monedero, V., Yebra, M.J., 2020. Unique Microbial Catabolic Pathway for the Human Core  
363 N-Glycan Constituent Fucosyl- $\alpha$ -1,6-*N*-Acetylglucosamine-Asparagine. *mBio* 11, e02804-  
364 e02819.  
365

366 Becker, D.J., Lowe, J.B., 2003. Fucose: biosynthesis and biological function in mammals.  
367 *Glycobiology* 13, 41R–53R.  
368

369 Bjellqvist, B., Basse, B., Olsen, E., Celis, J.E., 1994. Reference points for comparisons of two-  
370 dimensional maps of proteins from different human cell types defined in a pH scale where  
371 isoelectric points correlate with polypeptide compositions. *Electrophoresis* 15, 529–539.  
372

373 Cime-Castillo, J., Delannoy, P., Mendoza-Hernández, G., Monroy-Martínez, V., Harduin-  
374 Lepers, A., Lanz-Mendoza, H., Hernández-Hernández Fde, L., Zenteno, E., Cabello-Gutiérrez,  
375 C., Ruiz-Ordaz, B.H., 2015. Sialic acid expression in the mosquito *Aedes aegypti* and its  
376 possible role in dengue virus-vector interactions. *Biomed Res. Int.* 2015, 504187.  
377

378 Dawson, G., Tsay, G., 1977. Substrate specificity of human  $\alpha$ -L-fucosidase. *Arch. Biochem.*  
379 *Biophys.* 184, 12–23.  
380

381 Fabini, G., Freilinger, A., Altmann, F., Wilson, I.B.H., 2001. Identification of core  $\alpha$ -1,3-  
382 fucosylated glycans and cloning of the requisite fucosyltransferase cDNA from *Drosophila*  
383 *melanogaster*. Potential basis of the neural anti-horseradish peroxidase epitope. *J. Biol. Chem.*  
384 276, 28058–28067.  
385

386 Fuzita, F.J., Pinkse, M.W., Patane, J.S., Verhaert, P.D., Lopes, A.R., 2016. High throughput  
387 techniques to reveal the molecular physiology and evolution of digestion in spiders. *BMC*  
388 *Genomics* 17, 716.  
389

390 Geisler, C., Aumiller, J.J., Jarvis, D.L., 2008. A fused lobes gene encodes the processing  $\beta$ -N-  
391 acetylglucosaminidase in Sf9 cells. *J. Biol. Chem.* 283, 11330–11339.

392

393 Geisler, C., Jarvis, D.L., 2012. Substrate specificities and intracellular distributions of three N-  
394 glycan processing enzymes functioning at a key branch point in the insect N-glycosylation  
395 pathway. *J. Biol. Chem.* 287, 7084–7097.

396

397 Grootaert, H., Van Landuyt, L., Hulpiau, P., Callewaert, N., 2020. Functional exploration of  
398 the GH29 fucosidase family. *Glycobiology*, in press, doi: 10.1093/glycob/cwaa023.

399

400 Haines, N., Irvine, K.D., 2005. Functional analysis of *Drosophila*  $\beta$ 1,4-N-  
401 acetylgalactosaminyltransferases. *Glycobiology* 15, 335–346.

402

403 Hykollari, A., Malzl, D., Stanton, R., Eckmair, B., Paschinger, K., 2019. Tissue-specific  
404 glycosylation in the honeybee: Analysis of the N-glycomes of *Apis mellifera* larvae and venom.  
405 *Biochim Biophys Acta. Gen. Subj.* 1863, 129409.

406

407 Intra, J., Cenni, F., Perotti, M.E., 2006. An  $\alpha$ -L-fucosidase potentially involved in fertilization  
408 is present on *Drosophila* spermatozoa surface. *Mol. Reprod. Dev.* 73, 1149–1158.

409

410 Intra, J., Perotti, M.E., Pavesi, G., Horner, D., 2007. Comparative and phylogenetic analysis of  
411  $\alpha$ -L-fucosidase genes. *Gene* 392, 34–46.

412

413 Johnson, S.W., Alhadeff, J.A., 1991. Mammalian  $\alpha$ -L-fucosidases. *Comp Biochem. Physiol. B.*  
414 99, 479–488.

415

416 Kajiura, H., Hamaguchi, Y., Mizushima, H., Misaki, R., Fujiyama, K., 2015. Sialylation  
417 potentials of the silkworm, *Bombyx mori*; *B. mori* possesses an active  $\alpha$ 2,6-sialyltransferase.  
418 *Glycobiology* 12, 1441–1453.

419

420 Katayama, T., Sakuma, A., Kimura, T., Makimura, Y., Hiratake, J., Sakata, K., Yamanoi, T.,  
421 Kumagai, H., Yamamoto, K., 2004. Molecular cloning and characterization of *Bifidobacterium*

422 *bifidum* 1,2- $\alpha$ -L-fucosidase (AfcA), a novel inverting glycosidase (Glycoside Hydrolase Family  
423 95). J. Bacteriol. 186, 4885–4893.  
424

425 Kato, S., Hayashi, M., Kitagawa, M., Kajiura, H., Maeda, M., Kimura, Y., Igarashi, K.,  
426 Kasahara, M., Ishimizu, T., 2018. Degradation pathway of plant complex-type *N*-glycans:  
427 identification and characterization of a key  $\alpha$ 1,3-fucosidase from glycoside hydrolase family  
428 29. Biochem. J. 475, 305–317.  
429

430 Koles, K., Irvine, K.D., Panin, V.M., 2004. Functional characterization of *Drosophila*  
431 sialyltransferase. J. Biol. Chem. 279, 4346–4357.  
432

433 Koles, K., Lim, J.M., Aoki, K., Porterfield, M., Tiemeyer, M., Wells, L., Panin, V., 2007.  
434 Identification of *N*-glycosylated proteins from the central nervous system of *Drosophila*  
435 *melanogaster*. Glycobiology 17, 1388–1403.  
436

437 Kubeka, V., Altmann, F., Staudacher, E., Tretter, V., März, L., Hård, K., Kamerling, JP.,  
438 Vliegenthart, JF., 1993. Primary structures of the *N*-linked carbohydrate chains from honeybee  
439 venom phospholipase A<sub>2</sub>. Eur. J. Biochem. 213, 1193–1204.  
440

441 Kubelka, V., Altmann, F., März, L., 1995. The asparagine-linked carbohydrate of honeybee  
442 venom hyaluronidase. Glycoconj. J. 12, 77–83.  
443

444 Kumar, S., Stecher, G., Tamura, K., 2016. MEGA7: Molecular Evolutionary Genetics Analysis  
445 Version 7.0 for Bigger Datasets. Mol. Biol. Evol. 23, 1870–1874.  
446

447 Kuranda, M.J., Aronson, N.N. Jr., 1986. A di-*N*-acetylchitobiase activity is involved in the  
448 lysosomal catabolism of asparagine-linked glycoproteins in rat liver. J. Biol. Chem. 261, 5803–  
449 5809.  
450

451 Kurz, S., Aoki, K., Jin, C., Karlsson, N.G., Tiemeyer, M., Wilson, I.B.H., Paschinger, K., 2015.  
452 Targeted release and fractionation reveal glucuronylated and sulphated *N*- and *O*-glycans in  
453 larvae of dipteran insects. J. Proteomics 126, 172–188.  
454



455 Lei, Y., Yu, H., Dong, Y., Yang, J., Ye, W., Wang, Y., Chen, W., Jia, Z., Xu, Z., Li, Z., Zhang,  
456 F., 2015. Characterization of N-glycan structures on the surface of mature dengue 2 virus  
457 derived from insect cells. PLoS One. 10, e0132122.  
458

459 Li, T., Li, M., Hou, L., Wang, L., Sun, G., Chen, L., 2018. Identification and characterization  
460 of a core fucosidase from the bacterium *Elizabethkingia meningoseptica*. J. Biol. Chem. 293,  
461 1243–1258.  
462

463 Liu, S., Chen, C., Chang, S., Mong, K., Lin, C., Chang, C., Tang, C., Li, Y., 2009. Identification  
464 of essential residues of human  $\alpha$ -L-fucosidase and tests of its mechanism. Biochemistry 48,  
465 110–120.  
466

467 Liu, D., De Schutter, K., Smargiasso, N., De Pauw, E., Van Damme, E.J.M., Smagghe, G.,  
468 2019. The N-glycan profile of the peritrophic membrane in the colorado potato beetle larva  
469 (*Leptinotarsa decemlineata*). J. Insect Physiol. 115, 27–32.  
470

471 Lombard, V., Golaconda, Ramulu, H., Coutinho, P.M., Henrissat, B., 2014. The carbohydrate-  
472 active enzyme database (CAZy) in 2013. Nucleic Acids Res. 42, 490–495.  
473

474 Mabashi-Asazuma, H., Sohn, B.H., Kim, Y.S., Kuo, C.W., Khoo, K.H., Kucharski, C.A., Fraser,  
475 M.J. Jr, Jarvis, D.L., 2015. Targeted glycoengineering extends the protein N-glycosylation  
476 pathway in the silkworm silk gland. Insect Biochem. Mol. Biol. 65, 20–27.  
477

478

479 Minagawa, S., Sekiguchi, S., Nakaso, Y., Tomita, M., Takahisa, M., Yasuda, H., 2015.  
480 Identification of core alpha 1,3-fucosyltransferase gene from silkworm: an insect popularly  
481 used to express mammalian proteins. J. Insect. Sci. 15, 110.  
482

483 Miyazaki, T., Miyashita, R., Mori, S., Kato, T., Park, E.Y., 2019a. Expression and  
484 characterization of silkworm *Bombyx mori*  $\beta$ -1,2-N-acetylglucosaminyltransferase II, a key  
485 enzyme for complex-type N-glycan biosynthesis. J. Biosci. Bioeng. 127, 273–280.  
486

487 Miyazaki, T., Miyashita, R., Nakamura, S., Ikegaya, M., Kato, T., Park, E.Y., 2019b.  
488 Biochemical characterization and mutational analysis of silkworm *Bombyx mori*  $\beta$ -1,4-*N*-  
489 acetylgalactosaminyltransferase and insight into the substrate specificity of  $\beta$ -1,4-  
490 galactosyltransferase family enzymes. *Insect. Biochem. Mol. Biol.* 115, 103254.  
491  
492 Miyoshi, E., Nakano, M., 2008. Fucosylated haptoglobin is a novel marker for pancreatic  
493 cancer: detailed analyses of oligosaccharide structures, *Proteomics* 16, 3257–3262.  
494  
495 Moremen, K.W., Trimble, R.B., Herscovics, A., 1994. Glycosidases of the asparagine-linked  
496 oligosaccharide processing pathway. *Glycobiology* 4, 113–125.  
497  
498 Moremen, K.W., Tiemeyer, M., Nairn, A.V., 2012. Vertebrate protein glycosylation: diversity,  
499 synthesis and function. *Nat. Rev. Mol. Cell Biol.* 13, 448–462.  
500  
501 Nagae, M., Tsuchiya, A., Katayama, T., Yamamoto, K., Wakatsuki, S., Kato, R., 2007.  
502 Structural basis of the catalytic reaction mechanism of novel 1,2- $\alpha$ -L-fucosidase from  
503 *Bifidobacterium bifidum*. *J. Biol. Chem.* 282, 18497–18509.  
504  
505 Ndeh, D., Rogowsk, A., Cartmell, A., Luis, A.S., Basle, A., Gray, J., Venditto, I., Briggs, J.,  
506 Zhang, X., Labourel, A., Terrapon, N., Buffetto, F., Nepogodiev, S., Xiao, Y., Field, R.A., Zhu,  
507 Y., O’Neil, M.A., Urbanowicz, B.R., York, W.S., Davies, G.J., Abbott, D.W., Ralet, M.C.,  
508 Martens, E.C., Henrissat, B., Gilbert, H.J., 2017. Complex pectin metabolism by gut bacteria  
509 reveals novel catalytic function. *Nature* 544, 65–70.  
510  
511 Ohtsubo, K., Marth, J.D., 2006. Glycosylation in cellular mechanisms of health and disease.  
512 *Cell* 126, 855–867.  
513  
514 Opheim, D.J., Touster, O., 1977. The purification and characterization of rat liver lysosomal  $\alpha$ -  
515 L-fucosidase. *J. Biol. Chem.* 252, 739–743.  
516  
517 Paschinger, K., Staudacher, E., Stemmer, U., Fabini, G., Wilson, I.B.H., 2005.  
518 Fucosyltransferase substrate specificity and the order of fucosylation in invertebrates.  
519 *Glycobiology* 15, 463–474.

520  
521 Pasini, M.E., Intra, J., Pavesi, G., 2008. Expression study of an  $\alpha$ -L-fucosidase gene in the  
522 *Drosophilidae* family. *Gene* 420, 23–33.  
523  
524 Perrella, N.N., Fuzita, F.J., Moreti, R., Verhaert, P.D.E.M., Lopes, A.R., 2018a. First  
525 characterization of fucosidases in spiders. *Arch. Insect. Biochem. Physiol.* 98, e21462.  
526  
527 Perrella, N.N., Withers, S.G., Lopes, A.R., 2018b. Identity and role of the non-conserved  
528 acid/base catalytic residue in the GH29 fucosidase from spider *Nephilingis cruentata*.  
529 *Glycobiology* 28, 925–932.  
530  
531 Petersen, T.N., Brunak, S., von, Heijne, G., Nielsen, H., 2011. SignalP 4.0: discriminating  
532 signal peptides from transmembrane regions. *Nat. Methods* 8, 785–786.  
533  
534 Pogorelko, G.V., Reem, N.T., Young, Z.T., Chambers, L., Zobotina, O.A., 2016. Post-synthetic  
535 defucosylation of AGP by *Aspergillus nidulans*  $\alpha$ -1,2-fucosidase expressed in *Arabidopsis*  
536 apoplast induces compensatory upregulation of  $\alpha$ -1,2-fucosyltransferases. *PLoS One.* 7,  
537 e0159757.  
538  
539 Rahman, M.D., Maeda, M., Itano, S., Hossain, M.A., Ishimizu, T., Kimura, Y., 2017. Molecular  
540 characterization of tomato  $\alpha$ 1,3/4-fucosidase, a member of glycosyl hydrolase family 29  
541 involved in the degradation of plant complex type *N*-glycans. *J. Biochem.* 161, 421–432.  
542  
543 Rahman, M.D., Tsujimori, Y., Maeda, M., Hossain, M.A., Ishimizu, T., Kimura, Y., 2018.  
544 Molecular characterization of second tomato  $\alpha$ 1,3/4-fucosidase ( $\alpha$ -Fuc'ase SI-2), a member of  
545 glycosyl hydrolase family 29 active toward the core  $\alpha$ 1,3-fucosyl residue in plant *N*-glycans. *J.*  
546 *Biochem.* 164, 53–63.  
547  
548 Reddy, A., Gibbs, B.S., Liu, Y.L., Coward, J.K., Changchien, L.M., Maley, F., 1999.  
549 Glycosylation of the overlapping sequons in yeast external invertase: effect of amino acid  
550 variation on site selectivity in vivo and in vitro. *Glycobiology* 6, 547–555.  
551

552 Rodríguez-Díaz, J., Monedero, V., Yebra, M.J., 2011. Utilization of natural fucosylated  
553 oligosaccharides by three novel  $\alpha$ -L-fucosidases from a probiotic *Lactobacillus casei* strain.  
554 Appl. Environ. Microbiol. 77, 703–705.  
555

556 Sano, M., Hayakawa, K., Kato, I., 1992. Purification and characterization of  $\alpha$ -L-fucosidase  
557 from *Streptomyces* species. J. Biol. Chem. 267, 1522–1527.  
558

559 Sakurama, H., Tsutsumi, E., Ashida, H., Katayama, T., Yamamoto, K., Kumagai, H., 2012.  
560 Differences in the Substrate Specificities and Active-site Structures of Two  $\alpha$ -L-Fucosidases  
561 (Glycoside Hydrolase Family 29) from *Bacteroides thetaiotaomicron*. Biosci. Biotechnol.  
562 Biochem. 76, 1022–1024.  
563

564 Scheys, F., De Schutter, K., Shen, Y., Yu, N., Smargiasso, N., De Pauw, E., Van Damme,  
565 E.J.M., Smaghe, G., 2019. The N-glycome of the hemipteran pest insect *Nilaparvata lugens*  
566 reveals unexpected sex differences. Insect Biochem. Mol. Biol. 107, 39–45.  
567

568 Scheys, F., Van Damme, E.J.M., Smaghe, G., 2020. Let's talk about sexes: sex-related N-  
569 glycosylation in ecologically important invertebrates. Glycoconj. J. 37, 41–46.  
570

571 Schneider, C.A., Rasband, W.S., Eliceiri, K.W., 2012. NIH Image to ImageJ: 25 years of image  
572 analysis. Nat. Methods 9, 671–675.  
573

574 Sela, D.A., Garrido, D., Lerno, L., Wu, S., Tan, K., Eom, H.J., Joachimiak, A., Lebrila, C.B.,  
575 Mills, D.A., 2012. *Bifidobacterium longum* subsp. *Infantis* ATCC 15697  $\alpha$ -fucosidases are  
576 active on fucosylated human milk oligosaccharides. Appl. Environ. Microbiol. 78, 765–803.  
577

578 Sievers, F., Wilm, A., Dineen, D., Gibson, T.J., Karplus, K., Li, W., Lopez, R., McWilliam, H.,  
579 Remmert, M., Söding, J., Thompson, J.D., Higgins, D.G., 2011. Fast, scalable generation of  
580 high-quality protein multiple sequence alignments using Clustal Omega. Mol. Syst. Biol. 7, 539.  
581

582 Soya, S., Şahar, U., Karaçalı, S., 2016. Monosaccharide profiling of silkworm (*Bombyx mori*  
583 L.) nervous system during development and aging. Invert. Neurosci. 16, 8.  
584

585 Stanton, R., Hykollari, A., Eckmair, B., Malzl, D., Dragosits, M., Palmberger, D., Wang, P.,  
586 Wilson, I.B.H., Paschinger, K., 2017. The underestimated N-glycomes of lepidopteran species.  
587 Biochim Biophys Acta. Gen. Subj. 1861, 699–714.  
588  
589 Staudacher, E., Altmann, F., März, L., Hård, K., Kamerling, J.P., Vliegenthart, J.F., 1992.  $\alpha$  1-  
590 6( $\alpha$  1-3)-difucosylation of the asparagine-bound *N*-acetylglucosamine in honeybee venom  
591 phospholipase A<sub>2</sub>. Glycoconj. J. 2, 82–85.  
592  
593 Sulzenbacher, G., Bignon, C., Nishimura, T., Tarling, C.A., Withers, S.G., Henrissat, B.,  
594 Bourne, Y., 2004. Crystal structure of *Thermotoga maritima*  $\alpha$ -L-fucosidase. Insights into the  
595 catalytic mechanism and the molecular basis for fucosidosis. J. Biol. Chem. 279, 13119–13128.  
596  
597 Tarling, C.A., He, S., Sulzenbacher, G., Bignon, C., Bourne, Y., Henrissat, B., Withers, S.G.,  
598 2003. Identification of the catalytic nucleophile of the family 29  $\alpha$ -L-fucosidase from  
599 *Thermotoga maritima* through trapping of a covalent glycosyl-enzyme intermediate and  
600 mutagenesis. J. Biol. Chem. 278, 47394–47399.  
601  
602 Tretter, V., Altmann, F., März, L., 1991. Peptide-*N*<sup>4</sup>-(*N*-acetyl- $\beta$ -glucosaminyl) asparagine  
603 amidase F cannot release glycans with fucose attached  $\alpha$  1→3 to the asparagine-linked *N*-  
604 acetylglucosamine residue. Eur. J. Biochem. 199, 647–652.  
605  
606 Tsai, T., Li, S.T., Liu, C.P., Chen, K.Y., Shivatare, S.S., Lin, C.W., Liao, S.F., Hsu, T.L., Wu,  
607 Y.T., Tsai, M.H., Lai, M.Y., Lin, N.H., Wu, C.Y., Wong, C.H., 2017. An Effective Bacterial  
608 Fucosidase for Glycoprotein Remodeling. ACS Chem. Biol. 20, 63–72.  
609  
610 Vadaie, N., Jarvis, D.L., 2004. Molecular cloning and functional characterization of a  
611 Lepidopteran insect  $\beta$ 4-*N*-acetylgalactosaminyltransferase with broad substrate specificity, a  
612 functional role in glycoprotein biosynthesis, and a potential functional role in glycolipid  
613 biosynthesis. J. Biol. Chem. 279, 33501–33518.  
614  
615 Vainauskas, S., Kirk, C.H., Petralia, L., Guthrie, E.P., McLeod, E., Bielik, A., Luebbers, A.,  
616 Foster, J.M., Hokke, C.H., Rudd, P.M., Shi, X., Taron, C.H., 2018. A novel broad specificity

617 fucosidase capable of core  $\alpha$ 1-6 fucose release from *N*-glycans labeled with urea-linked  
618 fluorescent dyes. *Sci. Rep.* 8, 9504.  
619  
620 van Die, I., van Tetering, A., Bakker, H., van den Eijnden, D.H., Joziase, D.H., 1996.  
621 Glycosylation in lepidopteran insect cells: identification of a  $\beta$ 1  $\rightarrow$  4-*N*-  
622 acetylgalactosaminyltransferase involved in the synthesis of complex-type oligosaccharide  
623 chains. *Glycobiology* 6, 157–164.  
624  
625 Varki, A., Cummings, R.D., Esko, J.D., Freeze, H.H., Stanley, P., Marth, J.D., Bertozzi, C.R.,  
626 Hart, G.W., Etzler, M.E., 2009a. Symbol nomenclature for glycan representation. *Proteomics*  
627 9, 5398–5399.  
628  
629 Varki, A., Lowe, J.B., Cummings, R.D., Esko, J.D., Freeze, H.H., Stanley, P., Bertozzi, R.C.,  
630 Hart, W.G., Etzler, E.M., 2009b. *Essentials of Glycobiology*, second ed. Cold Spring Harbor  
631 Laboratory Press.  
632  
633 Varki, A., 2017. Biological roles of glycans. *Glycobiology* 27, 3–49.  
634  
635 Walski, T., De Schutter, K., Cappelle, K., Van Damme, E.J.M., Smagghe, G., 2017a.  
636 Distribution of glycan motifs at the surface of midgut cells in the cotton leafworm (*Spodoptera*  
637 *littoralis*) demonstrated by lectin binding. *Front. Physiol.* 8, 1020.  
638  
639 Walski, T., De Schutter, K., Van Damme, E.J., Smagghe, G., 2017b. Diversity and functions of  
640 protein glycosylation in insects. *Insect. Biochem. Mol. Biol.* 83, 21–34.  
641  
642 Walski, T., Van Damme, E.J., Smargiasso, N., Christiaens, O., De Pauw, E., Smagghe, G., 2016.  
643 Protein *N*-glycosylation and *N*-glycan trimming are required for postembryonic development  
644 of the pest beetle *Tribolium castaneum*. *Sci. Rep.* 6, 35151.  
645  
646 Wilson, I.B.H., Rendić, D., Freilinger, A., Dumić, J., Altmann, F., Mucha, J., Müller, S., Hauser,  
647 M.T., 2001. Cloning and expression of cDNAs encoding  $\alpha$ 1,3-fucosyltransferase homologues  
648 from *Arabidopsis thaliana*. *Biochim. Biophys. Acta.* 1527, 88–96.  
649

650 Winchester, B., 2005. Lysosomal metabolism of glycoproteins. *Glycobiology* 15, 1R–15R.  
651

652 Yamashita, K., Kochibe, N., Ohkura, T., Ueda, I., Kobata, A., 1985. Fractionation of L-fucose-  
653 containing oligosaccharides on immobilized *Aleuria aurantia* lectin. *J. Biol. Chem.* 260, 4688–  
654 4693.  
655

656 Yan, S., Serna, S., Reichardt, N.C., Paschinger, K., Wilson, I.B.H., 2013. Array-assisted  
657 characterization of a fucosyltransferase required for the biosynthesis of complex core  
658 modifications of nematode *N*-glycans. *J. Biol. Chem.* 288, 21015–21028.  
659

660 Zeleny, R., Leonard, R., Dorfner, G, Dalik, T., Kolarich, D., Altmann, F., 2006. Molecular  
661 cloning and characterization of a plant  $\alpha$ 1,3/4-fucosidase based on sequence tags from almond  
662 fucosidase I. *Phytochemistry* 67, 641–648.  
663

664 Zielinska, D.F., Gnad, F., Wisniewski, J.R., Mann, M., 2010. Precision mapping of an in vivo  
665 *N*-glycoproteome reveals rigid topological and sequence constraints. *Cell* 141, 897–907.  
666

667 **Legends to figures**

668 **Figure 1. Phylogenetic tree of BmFucA and characterized GH29  $\alpha$ -L-fucosidases.**

669 Amino acid sequences of all enzymes were obtained from GenBank. Sequence alignment  
670 was performed using Clustal Omega (Sievers et al., 2011) and a phylogenetic tree generated  
671 using MEGA 7 (Kumar et al., 2016). Sequences used were from *Bombyx mori*  
672 (XP\_004923234.1), *Drosophila melanogaster* (AAM50292.1), *Homo sapiens* (AAA52481.1),  
673 *Nephilingis cruentata* (Perrella et al., 2018a), *Omnitrophica bacterium* (KXK31601.1),  
674 *Xanthomonas campestris pv. campestris* (AAM42160.1), *F. graminearum* (AFR68935.1) *Bi.*  
675 *longum* subsp. *infantis* (ACJ51546.1), *L. casei* (CAQ67877.1), *L. casei* (CAQ67984.1),  
676 *Thermotoga maritima* (AAD35394.1), *Ba. thetaiotaomicron* (AAO78076.1), *E. anopheles*  
677 (WP\_047034007.1), *A. thaliana* (NP\_180377.2), *Ba. thetaiotaomicron* (AAO77299.1), and *S.*  
678 *pneumoniae* (AAK76203.1). Bootstrap values are indicated, and the scale bar shows the number  
679 of assumed amino acid replacements per site.

680

681 **Figure 2. General properties of GST-BmFucA.**

682 (A) SDS-PAGE analysis of purified GST-BmFucA with CBB staining. Lane 1, molecular  
683 weight marker; lane 2, supernatant of sonicated cell extract; and lane 3, purified GST-BmFucA.  
684 Black arrow indicates GST-BmFucA (89 kDa). pH dependence (B) and temperature  
685 dependence (C) of GST-BmFucA. pH dependence was measured at 30°C using 50 mM sodium  
686 citrate buffer (pH 3.0–6.0, circles), MES–NaOH buffer (pH 6.0–6.5, diamonds), or sodium  
687 phosphate buffer (pH 6.5–8.0, squares). Temperature dependence was measured at 25°C–80°C  
688 using 50 mM sodium citrate buffer (pH 5.0).

689

690 **Figure 3. Specificity of GST-BmFucA toward fucose-containing oligosaccharides.**

691 The substrates were assayed with GST-BmFucA and analyzed by TLC. Abbreviations used:  
692 Fuc, L-fucose; 2'-FL, 2'-fucosyllactose; 3-FL, 3-fucosyllactose; Lac, lactose; 6-FGn<sub>2</sub>,  
693 GlcNAc $\beta$ (1-4)[Fuc $\alpha$ (1-6)]GlcNAc. Black arrows indicate standards of L-fucose and lactose.

694

695 **Figure 4. Activity of GST-BmFucA toward core fucosylated N-glycans.**

696 (A–C) HPLC analysis of the reaction products generated by GST-BmFucA. Reaction  
697 mixtures containing MMF<sup>6</sup>-PA (A), GnGnF<sup>6</sup>-PA (B), and NaNaF<sup>6</sup>-PA (C) as substrates and  
698 GST-BmFucA were incubated in 50 mM sodium citrate buffer (pH 5.0) for 0, 20, and 60 min  
699 and analyzed using reversed-phase HPLC (see Materials and Methods). Substrates are shown



700 using symbols from the literature (Varki et al., 2009a). (D) Defucosylation of *N*-glycans  
701 attached to PLA<sub>2</sub> by GST-BmFucA. PLA<sub>2</sub> was incubated with GST-BmFucA and PNGaseF  
702 and analyzed using SDS-PAGE and lectin blotting (see Materials and Methods). Intact and  
703 deglycosylated PLA<sub>2</sub> are indicated with black and white arrows, respectively. Error bars  
704 indicate standard deviation. Asterisks indicate significant differences based on t-tests (\**p* < 0.05  
705 and \*\**p* < 0.01).  
706  
707

708 **Table 1. Kinetic parameters of GST-BmFucA and GH29  $\alpha$ -L-fucosidases from different**  
709 **species for pNP- $\alpha$ -L-Fuc.**

710

Species (Enzyme)	$k_{cat}$ ( $s^{-1}$ )	$K_m$ (mM)	$k_{cat}/K_m$ ( $s^{-1} mM^{-1}$ )
<i>Bombyx mori</i> (BmFucA)	12.5 $\pm$ 0.5	0.2 $\pm$ 0.02	62.7
<i>Nephilingis cruentata</i> (NcFuc) <sup>a</sup>	11 $\pm$ 1	0.4 $\pm$ 0.01	27.5
<i>Homo sapiens</i> (HsFucA1) <sup>b</sup>	17.1 $\pm$ 0.3	0.28 $\pm$ 0.02	61.1
<i>Thermotoga maritima</i> (TmFuc) <sup>c</sup>	14.3 $\pm$ 0.3	0.05 $\pm$ 0.003	280
<i>Elizabethkingia meningoseptica</i> (cFase I) <sup>d</sup>	8.4 $\pm$ 0.2	0.6 $\pm$ 0.05	13.9


711 <sup>a</sup>Perrella et al. (2018). <sup>b</sup>Liu et al. (2009). <sup>c</sup>Sulzenbacher et al. (2004). <sup>d</sup>Li et al., (2018).

712


713 **Table 2. Hydrolytic activity of GST-BmFucA and other  $\alpha$ -L-fucosidases for fucosylated**  
 714 **substrates.**  
 715

Substrate	Relative activity (%)									
	GH29-A						GH29-B			GH95
	BmFucA	HsFucA1 <sup>a</sup>	Blon_0426 <sup>b</sup>	AlfB <sup>c</sup>	AlfC <sup>c</sup>	BT_2970 <sup>d</sup>	cFaseI <sup>e</sup>	BT_2192 <sup>d</sup>	SpGH29 <sup>f</sup>	AfcA <sup>g</sup>
pNP- $\alpha$ -L-Fuc	100	–	100	100	100	100	7.53	0.025	–	–
4MU $\alpha$ Fuc	–	100	–	–	–	–	–	–	–	–
2'-FL	10.5	28.8	0.56	1.2	N.D.	0	N.D.	0.067	N.D.	100
3-FL	2.2	1	–	N.D.	N.D.	0	100	100	+	<1
6-FGn <sub>2</sub>	7.9	–	–	–	–	–	N.D.	–	–	N.D.
Fuc $\alpha$ 1-6GlcNAc	–	6.0	37.8	N.D.	61.3	0	N.D.	0	N.D.	–
3-FGn <sub>2</sub>	1.2	–	–	–	–	–	–	–	–	–
Fuc $\alpha$ 1-3GlcNAc	–	–	–	458.3	0.0018	5.3	–	0	N.D.	–


716 <sup>a</sup>Dawaon and Tsay. (1977). <sup>b</sup>Ashida et al., (2020). <sup>c</sup>Rodríguez-Díaz et al. (2011). <sup>d</sup>Sakurama et  
 717 al. (2012). <sup>e</sup>Li et al. (2018). <sup>f</sup>Ashida et al. (2009). <sup>g</sup>Katayama et al. (2004). +, detected on TLC;  
 718 –, not tested; N.D., not detected. The highest activity was defined as 100% for each enzyme.  
 719



0.2




**A**



BmFucA	-	-	+	-	+	-
		2'-FL		3-FL		Lac
	Fuc					

**B**



BmFucA	-	-	+
		6-FGn <sub>2</sub>	
	Fuc		

← Fuc

← Fuc

← Lac

