

A unique mechanism of successful fertilization in a domestic bird

著者	Sasanami Tomohiro, Izumi Shunsuke, Sakurai Naoki, Hirata Toshifumi, Mizushima Shusei, Matsuzaki Mei, Hiyama Gen, Yarinaga Eriko, Yoshimura Takashi, Ukena Kazuyoshi, Tsutsui Kazuyoshi
journal or publication title	Scientific Reports
volume	5
page range	7700
year	2015-01-09
出版者	Nature Publishing Group
権利	http://creativecommons.org/licenses/by/4.0/
URL	http://hdl.handle.net/10297/8009

doi: 10.1038/srep07700

A unique mechanism of successful fertilization in a domestic bird

Tomohiro Sasanami¹, Shunsuke Izumi², Naoki Sakurai³, Toshifumi Hirata², Shusei Mizushima¹, Mei Matsuzaki¹, Gen Hiyama¹, Eriko Yorinaga⁴, Takashi Yoshimura⁴, Kazuyoshi Ukena⁵, Kazuyoshi Tsutsui^{6*}

¹Department of Applied Biological Chemistry, Faculty of Agriculture, Shizuoka University, 836 Ohya, Shizuoka 422-8529, Japan.

²Department of Mathematical and Life Sciences, Graduate School of Science, Hiroshima University, Higashi-Hiroshima 739-8526, Japan.

³Department of Environmental Dynamics and Management, Graduate School of Biosphere Science, Hiroshima University, Higashi-Hiroshima 739-8526, Japan.

⁴Graduate School of Bioagricultural Sciences, Nagoya University, Furo-cho, Chikusa-ku, Nagoya 464-8601, Japan.

⁵Section of Behavioural Sciences, Division of Human Sciences, Graduate School of Integrated Arts and Sciences, Hiroshima University, Higashi-Hiroshima 739-8526, Japan.

⁶Department of Biology and Center for Medical Life Science, Waseda University, 2-2 Wakamatsu-cho, Shinjuku, Tokyo 162-8480, Japan.

*Corresponding to: Kazuyoshi Tsutsui, Department of Biology and Center for Medical Life Science, Waseda University, 2-2 Wakamatsu-cho, Shinjuku, Tokyo 162-8480, Japan. Email: k-tsutsui@waseda.jp

Supplementary Information

Table S1: Free monosaccharide composition from water soluble extracts of the cloacal gland and testis.

Figure S1: Effects of AL8110 on the PGF_{2α}-induced spontaneous contractions of the isolated vagina.

Figure. S2: Expression of the receptor for PGF_{2α} in the utero-vaginal junction (UVJ).

Caption to Movie S1: Appearance of cloacal gland secretion (CGS).

Data of Structural analysis

Table S1. Free monosaccharide composition from water soluble extracts of the cloacal gland and testis.

Gland	Sugar content					
	Gal	Fru	Glc	Gal	Fru	Glc
	(µg/g)			(µg/gland)		
Cloacal gland	115	43	85	75	28	55
Testis	45	49	0*	136	148	0*

Free fructose (Fru) content in the extracts was determined by an HPLC system equipped with a pulsed amperometric detector, and free galactose (Gal) and glucose (Glc) were reduced and acetylated to alditol acetate form and were determined by GLC using inositol as an internal standard. *, Free glucose (Glc) in the testis was not detected by gas chromatography.




Figure S1 | Effects of AL 8110 on the PGF_2 -induced spontaneous contractions of the isolated vagina. The upward arrow indicated application of each chemical.




Figure S2 | Expression of the receptor for PGF_{2α}. **a**, Autoradiograms of the UVJ sections after hybridization with ³³P-labeled antisense probe specific for the receptor for PGF_{2α} (upper panel) or sense probe (lower panel) are shown. Representative results of two experiments are shown (n = 2). **b**, RT-PCR analysis of the receptor for PGF_{2α}. The infundibulum (Inf), magnum (Magn), isthmus (Isthm), uterus (Uter), UVJ or vagina (Vag) isolated from females at 14 h after egg-laying were processed for RT-PCR analysis using gene-specific primers. Data were normalized with S17 ribosomal protein and expressed as the mean ± SEM of three independent experiments. **c**, RT-PCR analysis of the receptor for PGF_{2α} during ovulatory cycle. The UVJ isolated from females at 8, 14, 20 or 25 h after egg-laying were processed for RT-PCR analysis using gene-specific primers. Data were normalized with S17 ribosomal protein and expressed as the mean ± SEM of three independent experiments. **d**, Comparison of sperm filling rate during ovulatory cycle. Females were mated at 8, 14, 20 or 25 h after egg-laying. The utero-vaginal junction mucosa was isolated from the bird, and SST was observed under fluorescence microscope. Sperm filling rate was calculated and expressed as the mean ± SEM. 3–5 birds were used within each treatment.




Figure S3 | Purification scheme of bioactive substance from cloacal gland extracts. TFA: trifluoroacetic acid , RM: retained materials, ACN: acetonitrile, PB: phosphate buffer.

Caption to Movie S1

Appearance of cloacal gland secretion (CGS).

The cloacal gland (CG) of male quail ejects CGS as a meringue-like foam by pushing lightly with the thumb and index finger.

Data of structural analysis

The MS, IR and ¹H NMR spectra of the isolated bioactive substance were as follows:

MS m/z 355 (M+1), 337 [(M-H₂O)+1], 319 [(M-2H₂O)+1] and 301 [(M-3H₂O)+1]; IR (Liq. film) ν 3400-3300 (OH) and 1708 cm⁻¹ (COOH); ¹H NMR δ 0.89 (3H, t, J=6.5 Hz, 20-H₃), 4.02 (1H, m, 11-H), 4.22 (1H, q, J=6.2 Hz, 15-H), 4.25 (1H, dd, J=5.4 and 4.5 Hz, 9-H), 5.38 (1H, bq, J=7.5 Hz, 5-H), 5.53 (1H, bq, J=7.5 Hz, 6-H), 5.59 and 5.60 (2H, m, 13- and 14-H). The spectroscopic data of the isolated bioactive substance were found to be prostaglandin F₂α (PGF₂α) as compared with the spectra of authentic specimens, such as PGF₂α, (15R)-PGF₂α and (13Z, 15R)-PGF₂α (Cayman Chemical).