

## Purification and characterization of a lectin from the mushroom *Hypsizygus marmoreus*

|                              |  |
|------------------------------|--|
| 著者                           | Suzuki Tomohiro, Abe Tomoya, Umehara Kanako, Choi Jae-Hoon, Hirai Hirofumi, Dohra Hideo, Kawagishi Hirokazu  |
| journal or publication title | Mycoscience  |
| volume                       | 56   |
| number                       | 4  |
| page range                   | 359-363  |
| year                         | 2015-07  |
| 出版者                          | Elsevier   |
| 権利                           | Copyright (C) 2014 The Mycological Society of Japan. Published by Elsevier Japan KK All rights reserved.   |
| 注記                           | NOTICE: this is the author's version of a work that was accepted for publication in Mycoscience. Changes resulting from the publishing process, such as peer review, editing, corrections, structural formatting, and other quality control mechanisms may not be reflected in this document. Changes may have been made to this work since it was submitted for publication. A definitive version was subsequently published in Mycoscience, Volume 56, Issue 4, July 2015, doi:10.1016/j.myc.2014.11.001 |
| URL                          | <a href="http://hdl.handle.net/10297/8160">http://hdl.handle.net/10297/8160</a>  |

1 Note

2 Title: Purification and characterization of a lectin from the mushroom *Hypsizigus marmoreus*

3

4 Tomohiro Suzuki <sup>a</sup>, Tomoya Abe <sup>b</sup>, Kanako Umehara <sup>b</sup>, Jae-Hoon Choi <sup>b</sup>, Hirofumi Hirai <sup>b</sup>,

5 Hideo Dohra <sup>a</sup>, Hirokazu Kawagishi <sup>a, b, c\*</sup>

6

7 <sup>a</sup> Research Institute of Green Science and Technology, Shizuoka University, 836 Ohya,

8 Suruga-ku, Shizuoka 422-8529, Japan

9 <sup>b</sup> Graduate School of Agriculture, Shizuoka University, 836 Ohya, Suruga-ku, Shizuoka

10 422-8529, Japan

11 <sup>c</sup> Graduate School of Science and Technology, Shizuoka University, 836 Ohya, Suruga-ku,

12 Shizuoka 422-8529, Japan

13

14 \*Corresponding author:

15 H. Kawagishi

16 Tel: +81-54-238-4885

17 Fax: +81-54-238-4885

18 E-mail: [achkawa@ipc.shizuoka.ac.jp](mailto:achkawa@ipc.shizuoka.ac.jp)

19

20 Text: 9 pages; tables: 3; figures: 3

21

1 Abstract

2 HML (*Hypsizigus marmoreus* lectin) was isolated from the mushroom *Hypsizigus*  
3 *marmoreus* using CM cation exchange, bovine submaxillary gland mucin affinity column  
4 and TSK-GEL G3000SW gel filtration chromatography. The results of SDS-PAGE,  
5 MALDI-TOF MS and gel filtration analysis of HML indicated that the lectin was a dimer  
6 with each subunit of 9.5 kDa. The partial amino acid sequences of HML were determined by  
7 N-terminal sequencing of peptides obtained by trypsin or Glu-C endopeptidase digest of the  
8 lectin. In the hemagglutination inhibition assay, HML did not bind to any mono- or  
9 oligo-saccharides tested. Among the glycoproteins examined, asialo-fetuin was the strongest  
10 inhibitor.

11

12 Keywords: Carbohydrate-binding specificity; *Lyophyllaceae* family; Partial amino acid  
13 sequences

14

15

16

1 Lectins are proteins or glycoproteins that are capable of specific recognition and binding to  
2 carbohydrate moieties of glycoproteins or glycolipids (Goldstein et al. 1980). Thus, lectins  
3 are used in studies of cell surface carbohydrate structures and oligosaccharides structures of  
4 glycoconjugates (Sharon and Lis 2004; Sharon 2007). Recently, new technologies such as  
5 lectin microarray have been in focus, therefore lectins with ever-unknown specific sugar  
6 binding properties are useful as a molecule that can be spotted on the array (Hirabayashi  
7 2004; Kuno et al. 2005; Meany et al. 2011). Among lectins, mushroom lectins represent a  
8 class of intensely studied biomolecules, and more than 50 mushroom lectins have been  
9 purified to date and have attracted wide attention due to their biological activities  
10 (Kawagishi 1995; Kobayashi et al. 2004a,b; Kobayashi et al. 2005; Horibe et al. 2010;  
11 Kobayashi et al. 2012; Suzuki et al. 2012; Xu et al. 2014).

12 *Hypsizigus marmoreus* (bunashimeji in Japanese) is as popular as *Lentinula edodes*  
13 (shiitake) and *Flammulina velutipes* (enokitake) as an edible mushroom in Japan and readily  
14 available at supermarkets nationwide. Besides its good taste, many biological activities have  
15 been discovered in *H. marmoreus*: antioxidant (Matsuzawa et al. 1998; Lee et al. 2007),  
16 antiproliferative (Lam and Ng 2001), ribosome inactivating (Lam and Ng 2001; Wong et al.  
17 2008), antiviral (Akihisa et al. 2005; Suzuki et al. 2011), anticancer (Chang et al. 2004;  
18 Chang et al. 2009) and ribonuclease (Guan et al. 2007) activities. However, there are no  
19 reported lectin or hemagglutinin from this mushroom.

20 In the course of our continuing screening for new mushroom lectins, we found lectin  
21 activity in the extract of *H. marmoreus*. Here, we describe the isolation and characterization  
22 of the lectin from this mushroom.

23

#### 24 *Purification of the lectin*

25 The purification procedure is summarized in Table 1. In the present study, bovine

1 submaxillary gland mucin (BSM)-Toyopearl was chosen as the affinity support of choice as  
2 the hemagglutination activity of the crude extract from the mushroom was inhibited by BSM  
3 (Kobayashi 2004b). Initially, lyophilized fruiting bodies of *H. marmoreus* (JA Zennou) were  
4 homogenized and extracted with 50 mM acetate buffer, pH 5.0, overnight. The homogenate  
5 was centrifuged at 8,500g for 25 min and the resulting supernatant was applied to a column  
6 (5 × 13 cm) of CM-Toyopearl 650M (Tosoh, Tokyo, Japan) equilibrated with the buffer.  
7 After unbound materials were washed with the buffer, the bound fraction was eluted with 0.5  
8 M NaCl in the buffer. The eluates were concentrated and equilibrated with 50 mM acetate  
9 buffer, pH 5.0, by ultrafiltration, and applied to a BSM-Toyopearl column (2.5 × 20 cm)  
10 equilibrated in the same buffer. After all unbound substances were removed by washing the  
11 column with the buffer, the bound fraction was then eluted by 0.5 M acetic acid. The  
12 lectin-containing fraction was dialyzed against 10 mM phosphate-buffered saline, pH 7.4  
13 (PBS), containing 0.1% (w/v) sodium dodecyl sulfate (SDS), and applied to a TSK  
14 G3000SW column (7.8 × 300 mm; Tosoh, Tokyo, Japan) equilibrated in the same buffer.  
15 The eluant was dialyzed extensively against distilled water and ultra-filtered to give *H.*  
16 *marmoreus* lectin (HML).

17

#### 18 *Molecular properties of HML*

19 HML gave a single band with an apparent mass of 10 kDa on SDS-PAGE (Laemmli 1970)  
20 in the presence of 2-mercaptoethanol (Fig. 1). Gel filtration for determination of its  
21 molecular mass by HPLC was carried out on a TSK-gel G3000SW<sub>XL</sub> column (7.8 × 300  
22 mm) operating at room temperature in PBS at a flow rate of 0.5 ml/min. The molecular mass  
23 was calibrated with the standard proteins (Sigma, MO, USA). HML gave a single  
24 symmetrical peak at an elution volume corresponding to a molecular mass of about 18 kDa  
25 (Fig. 2). Matrix-assisted laser desorption ionization-time-of-flight mass spectrometry

1 (MALDI-TOF MS) analysis was performed on AutoFlex (Bruker Daltonics, MA, USA),  
2 using 10 mg/ml super 2-hydroxy-5-methoxybenzoic acid (Bruker Daltonics), as a matrix.  
3 The MALDI-TOF mass spectrum of the lectin gave a main peak at  $m/z$  9,576 and a small  
4 peak at  $m/z$  19,251 (Fig. 3). The results of SDS-PAGE, gel filtration, and MALDI-TOF MS  
5 analysis indicated that the molecular mass of HML was 19 kDa and it was composed of two  
6 identical 9.5 kDa-subunits.

7 *N*-terminal amino acid sequence analysis of intact HML suggested that the amino  
8 terminus of HML was blocked. Thus, HML was digested with sequencing grade modified  
9 trypsin (Promega, WI, USA) or Endoproteinase Glu-C from *Staphylococcus aureus* strain V8  
10 (Sigma), and the resulting peptides were isolated by reverse-phase HPLC. Each purified  
11 peptide sequence was analyzed on a PPSQ-21A protein sequencer (Shimadzu, Kyoto, Japan).  
12 As a result, three partial amino acid sequences of HML were determined as shown in Table 2.  
13 The partial amino acid sequences of these three peptides obtained by protein sequencing  
14 were very short, but covered about 40% of HML judging from its molecular mass. These  
15 sequences show similarity to those of many proteins by BLASTP search against  
16 non-redundant (nr) protein sequences and Swiss-Prot database. However, no known lectins  
17 were found by the homology search, indicating that HML might be a new type of lectin.  
18 Further amino acid sequencing analysis or molecular cloning of HML is needed to conclude  
19 it.

20

#### 21 *Properties of HML as a lectin*

22 The hemagglutinating activity of the lectin was determined by a two-fold serial dilution  
23 procedure using Pronase-treated rabbit erythrocytes. The hemagglutination titer was defined  
24 as a reciprocal of the highest dilution exhibiting hemagglutination. Inhibition was expressed  
25 as the minimum concentration of each sugar or glycoprotein required for inhibition of

1 hemagglutination of titer 4 of the lectin. Table 3 shows the inhibition of hemagglutination  
2 activity of HML by various monosaccharides, oligosaccharides, and glycopeptides. HML did  
3 not bind to any mono- or oligo-saccharides tested. Among the glycoproteins examined, the  
4 most potent inhibitor was asialo-fetuin (minimum inhibitory concentration (MIC), 7.81  
5  $\mu\text{g/ml}$ ). Thyroglobulin, fetuin, BSM and  $\alpha_1$ -acid glycoprotein were also inhibitory at higher  
6 concentrations, 15.6, 15.6, 31.3 and 125  $\mu\text{g/ml}$ , respectively, than asialo-fetuin. Asialo-fetuin  
7 that carries both *N* and *O* linked oligosaccharides was the most potent inhibitor of the  
8 hemagglutination activity. Fetuin has a triantennary structure with terminal Gal $\beta$ 1-4GlcNAc  
9 and three *O*-linked structure (Townsend et al. 1986). Thyroglobulin contains mainly  
10 diantennary Gal $\beta$ 1-4GlcNAc *N*-glycans (Takeya et al. 2007). Mucin contains several  
11 Gal $\beta$ 1-3GalNAc $\alpha$ 1 *O*-glycans. HML showed high affinity for fetuin and thyroglobulin. The  
12 affinity of HML towards fetuin and thyroglobulin suggested that the Gal $\beta$ 1-4GlcNAc  
13 *N*-glycans structure was important for the interaction with HML. In basidiomycetes, lectins  
14 from *Volvariella volvacea* (VVL) (She et al. 1998), *Mycocleptodonoids aitchisonii* (MAL)  
15 (Kawagishi et al. 2001) and *Ganoderma lucidum* (Thakur et al. 2007) have been reported  
16 with hemagglutination activity inhibited only by glycoproteins. Hemagglutinating activity of  
17 VVL was inhibited by thyroglobulin and MAL exhibited strong sugar binding specificity  
18 towards asialo BSM. In the case of *G. lucidum* lectin, asialo-fetuin was most potent inhibitor.  
19 The results of the hemagglutination inhibition test of HML suggest that the  
20 carbohydrate-binding specificity of HML was most similar to that of *G. lucidum* lectin. At  
21 present, we cannot explain the reason why the hemagglutination mediated by HML was  
22 inhibited by BSM, but asialo-BSM, PSM (porcine stomach mucin) and asialo-PSM. More  
23 detailed analysis of its binding specificity is necessary.

24 Some lectins can specifically bind to fucose, galactose/*N*-acetylgalactosamine,  
25 mannose/glucose, *N*-acetylglucosamine, sialic acid or complex oligosaccharides

1 (Guzman-Partida et al. 2004). Therefore, such lectins have become invaluable tools for  
2 chemical and biological research of carbohydrate-containing compounds as well as for  
3 clinical uses (Van Damme et al. 1997; Loris et al. 1998; Lam and Ng 2011). The result of the  
4 hemagglutination inhibition test of HML suggested that the lectin belongs to the group  
5 termed “complex” with specificity towards *N*-glycans. FVA-L from *Flammulina velutipes*  
6 (Ng et al. 2006), lectin IV from *Griffonia simplicifolia* (Delbaere et al. 1993), PHA-L from  
7 *Phaseolus vulgaris* (Loris et al. 1998), PF2 from *Olneya tesota* (Guzman-Partida et al. 2004)  
8 and VL1 from *Acacia constricta* (Guzman-Partida et al. 2004) also belong to this group. To  
9 our knowledge, this is the first report of isolation and characterization of the lectin from  
10 *Hypsizigus marmoreus*.

11



## 1 **Disclosure**

2 The authors declare no conflict of interest. All the experiments undertaken in this study  
3 comply with the current laws of Japan.

4

## 5 **Acknowledgments**

6 This work was partially supported by a Grant-in-Aid for Scientific Research on Innovative  
7 Areas “Chemical Biology of Natural Products” from MEXT (Grant Number 24102513).

8

9

## 10 **References**

- 11 Akihisa T, Franzblau SG, Tokuda H, Tagata M, Ukiya M, Matsuzawa T, Metori K, Kimura Y, Suzuki  
12 T, Yasukawa K, 2005. Antitubercular activity and inhibitory effect on Epstein-Barr virus activation  
13 of sterols and polyisoprenepolyols from an edible mushroom, *Hypsizigus marmoreus*. *Biological  
14 and Pharmaceutical Bulletin* 28: 1117–1119.
- 15 Chang JS, Bae JT, Oh EJ, Kim JY, Park SH, Lee KR, 2009. Cancer preventive potential of methanol  
16 extracts of *Hypsizigus marmoreus*. *Journal of Medicinal Food* 12: 493–500; doi:  
17 10.1089/jmf.2008.1050.
- 18 Chang JS, Son JK, Li G, Oh EJ, Kim JY, Park SH, Bae JT, Kim HJ, Lee IS, Kim OM, Kozukue N,  
19 Han JS, Hirose M, Lee KR, 2004. Inhibition of cell cycle progression on HepG2 cells by  
20 hypsiziprenol A9, isolated from *Hypsizigus marmoreus*. *Cancer Letters* 212: 7–14.
- 21 Delbaere LT, Vandonselaar M, Prasad L, Quail JW, Wilson KS, Dauter Z, 1993. Structures of the  
22 lectin IV of *Griffonia simplicifolia* and its complex with the Lewis b human blood group  
23 determinant at 2.0 Å resolution. *Journal of Molecular Biology* 230: 950–965.
- 24 Goldstein IJ, Hughes RC, Monsigny M, Osawa T, Sharon N, 1980. What should be called a lectin?  
25 *Nature* 285: 66.
- 26 Guan GP, Wang HX, Ng TB, 2007. A novel ribonuclease with antiproliferative activity from fresh  
27 fruiting bodies of the edible mushroom *Hypsizigus marmoreus*. *Biochimica et Biophysica Acta*  
28 1770: 1593–1597.
- 29 Guzman-Partida AM, Robles-Burgueno MR, Ortega-Nieblas M, Vazquez-Moreno I, 2004.  
30 Purification and characterization of complex carbohydrate specific isolectins from wild legume  
31 seeds: *Acacia constricta* is (vinorama) highly homologous to *Phaseolus vulgaris* lectins.  
32 *Biochimie* 86: 335–342.
- 33 Hirabayashi J, 2004. Lectin-based structural glycomics: glycoproteomics and glycan profiling.  
34 *Glycoconjugate Journal* 21: 35–40; doi: 10.1023/B:GLYC.0000043745.18988.a1.
- 35 Horibe M, Kobayashi Y, Dohra H, Morita T, Murata T, Usui T, Nakamura-Tsuruta S, Kamei M,

- 1 Hirabayashi J, Matsuura M, Yamada M, Saikawa Y, Hashimoto K, Nakata M, Kawagishi H, 2010.  
2 Toxic isolectins from the mushroom *Boletus venenatus*. *Phytochemistry* 71: 648–657; doi:  
3 10.1016/j.phytochem.2009.12.003.
- 4 Kawagishi H, 1995. Mushroom lectins. *Food Reviews International* 11: 63–68.
- 5 Kawagishi H, Takagi J-i, Taira T, Murata T, Usui T, 2001. Purification and characterization of a lectin  
6 from the mushroom *Mycocleptodonoides aitchisonii*. *Phytochemistry* 56: 53–58; doi:  
7 10.1016/S0031-9422(00)00351-4.
- 8 Kobayashi Y, Ishizaki T, Kawagishi H, 2004a. Screening for lectins in wild and cultivated  
9 mushrooms from Japan and their sugar-binding specificities. *International Journal of Medicinal*  
10 *Mushrooms* 6: 117–130.
- 11 Kobayashi Y, Kobayashi K, Umehara K, Dohra H, Murata T, Usui T, Kawagishi H, 2004b.  
12 Purification, characterization, and sugar binding specificity of an *N*-Glycolylneuraminic  
13 acid-specific lectin from the mushroom *Chlorophyllum molybdites*. *Journal of Biological*  
14 *Chemistry* 279: 53048–53055.
- 15 Kobayashi Y, Nakamura H, Sekiguchi T, Takanami R, Murata T, Usui T, Kawagishi H, 2005.  
16 Analysis of the carbohydrate binding specificity of the mushroom *Pleurotus ostreatus* lectin by  
17 surface plasmon resonance. *Analytical Biochemistry* 336: 87–93.
- 18 Kobayashi Y, Tateno H, Dohra H, Moriwaki K, Miyoshi E, Hirabayashi J, Kawagishi H, 2012. A  
19 novel core fucose-specific lectin from the mushroom *Pholiota squarrosa*. *Journal of Biological*  
20 *Chemistry* 287: 33973–33982; doi: 10.1074/jbc.M111.327692.
- 21 Kuno A, Uchiyama N, Koseki-Kuno S, Ebe Y, Takashima S, Yamada M, Hirabayashi J, 2005.  
22 Evanescent-field fluorescence-assisted lectin microarray: a new strategy for glycan profiling.  
23 *Nature Methods* 227: 680–685; doi:10.1038/nmeth803.
- 24 Laemmli UK, 1970. Cleavage of structural proteins during the assembly of the head of bacteriophage  
25 T4. *Nature* 227: 680–685.
- 26 Lam SK, Ng TB, 2001. Hypsin, a novel thermostable ribosome-inactivating protein with antifungal  
27 and antiproliferative activities from fruiting bodies of the edible mushroom *Hypsizigus marmoreus*.  
28 *Biochemical and Biophysical Research Communications* 285: 1071–1075.
- 29 Lam SK, Ng TB, 2011. Lectins: production and practical applications. *Applied Microbiology and*  
30 *Biotechnology* 89: 45–55; doi: 10.1007/s00253-010-2892-9.
- 31 Lee Y, Yen M, Mau J, 2007. Antioxidant properties of various extracts from *Hypsizigus marmoreus*.  
32 *Food Chemistry* 104: 1–9; doi: 10.1016/j.foodchem.2006.10.063.
- 33 Loris R, Hamelryck T, Bouckaert J, Wyns L, 1998. Legume lectin structure. *Biochimica et*  
34 *Biophysica Acta* 1383: 9–36.
- 35 Matsuzawa T, Saitoh H, Sano M, Tomita I, Ohkawa M, Ikekawa T, 1998. Studies on antioxidant  
36 effects of *Hypsizigus marmoreus*. II. Effects of *Hypsizigus marmoreus* for antioxidant activities of  
37 tumor-bearing mice. *Yakugaku Zasshi* 118: 476–481.
- 38 Meany DL, Hackler L, Zhang H, Chan DW, 2011. Tyramide signal amplification for

1 antibody-overlay lectin microarray: A strategy to improve the sensitivity of targeted glycan  
2 profiling. *Journal of Proteome Research* 10: 1425–1431; doi:10.1021/pr1010873.

3 Ng TB, Ngai PH, Xia L, 2006. An agglutinin with mitogenic and antiproliferative activities from the  
4 mushroom *Flammulina velutipes*. *Mycologia* 98: 167–171.

5 Sharon N, 2007. Lectins: carbohydrate-specific reagents and biological recognition molecules.  
6 *Journal of Biological Chemistry* 282: 2753–2764.

7 Sharon N, Lis H, 2004. History of lectins: from hemagglutinins to biological recognition molecules.  
8 *Glycobiology* 14: 53R–62R.

9 She QB, Ng TB, Liu WK, 1998. A novel lectin with potent immunomodulatory activity isolated from  
10 both fruiting bodies and cultured mycelia of the edible mushroom *Volvariella volvacea*.  
11 *Biochemical and Biophysical Research Communications* 247: 106–111; doi:  
12 10.1006/bbrc.1998.8744.

13 Suzuki T, Sugiyama K, Hirai H, Ito H, Morita T, Dohra H, Murata T, Usui T, Tateno H, Hirabayashi J,  
14 Kobayashi Y, Kawagishi H, 2012. Mannose-specific lectin from the mushroom *Hygrophorus*  
15 *russula*. *Glycobiology* 22: 616–629; doi: 10.1093/glycob/cwr187.

16 Suzuki T, Umehara K, Tashiro A, Kobayashi Y, Dohra H, Hirai H, Kawagishi H, 2011. An antifungal  
17 protein from the culinary-medicinal beech mushroom, *Hypsizygus marmoreus* (Peck) Bigel.  
18 (Agaricomycetidae). *International Journal of Medicinal Mushrooms* 13: 27–31.

19 Takeya A, Hosomi O, Nishijima H, Ohe Y, Sugahara K, Sagi M, Yamazaki K, Hayakawa H,  
20 Takeshita H, Sasaki C, Kogure T, Mukai T, 2007. Presence of  $\beta$ -linked GalNAc residues on  
21 N-glycans of human thyroglobulin. *Life Sciences* 80: 538–545.

22 Thakur A, Rana M, Lakhanpal TN, Ahmad A, Khan MI, 2007. Purification and characterization of  
23 lectin from fruiting body of *Ganoderma lucidum*: Lectin from *Ganoderma lucidum*. *Biochimica et*  
24 *Biophysica Acta* 1770: 1404–1412; doi: 10.1016/j.bbagen.2007.05.009.

25 Townsend RR, Hardy MR, Wong TC, Lee YC, 1986. Binding of N-linked bovine fetuin  
26 glycopeptides to isolated rabbit hepatocytes: Gal/GalNAc hepatic lectin discrimination between  
27 Gal  $\beta$ (1,4)GlcNAc and Gal  $\beta$ (1,3)GlcNAc in a triantennary structure. *Biochemistry* 25:  
28 5716–5725.

29 Van Damme EJ, Van Leuven F, Peumans WJ, 1997. Isolation, characterization and molecular cloning  
30 of the bark lectins from *Maackia amurensis*. *Glycoconjugate Journal* 14: 449–456.

31 Wong JH, Wang HX, Ng TB, 2008. Marmorin, a new ribosome inactivating protein with  
32 antiproliferative and HIV-1 reverse transcriptase inhibitory activities from the mushroom  
33 *Hypsizygus marmoreus*. *Applied Microbiology and Biotechnology* 81: 669–674; doi:  
34 10.1007/s00253-008-1639-3.


35 Xu CJ, Wang YX, Niu BN, Liu B, Li YB, Wang XM, Lu SL, 2014. Isolation and characterization of a  
36 novel lectin with mitogenic activity from *Pleurotus ferulae*. *Pakistan journal of pharmaceutical*  
37 *sciences* 27: 983–989.

38

1

2 **Figure legends**

3



4

5 Fig. 1 – SDS-PAGE of HML. Lane M, marker proteins; lane 1, HML reduced with

6 2-mercaptoethanol. Samples were heated in the presence of 2-mercaptoethanol for 10 min at

7 100°C and SDS-PAGE (12% polyacrylamide gel) was done by the Laemmli method


8 (Laemmli 1970), with Coomassie Brilliant Blue staining. The molecular mass on SDS-PAGE

9 was estimated using recombinant ladder markers (10 to 100 kDa; XL-Ladder Low; APRO

10 Life Science Institute, Tokushima, Japan).

11

12



1

2 Fig. 2 – HPLC profile of HML. Gel filtration by HPLC was carried out on a TSK-gel

3 G3000SW<sub>XL</sub> column (7.8 × 300 mm) operating at room temperature in PBS at a flow rate of

4 0.5 mL/min. Fractions were collected by monitoring absorbance at 280 nm. The molecular

5 mass was calibrated with the following standard proteins (Sigma); bovine thyroglobulin (669

6 kDa), horse spleen apoferritin (443 kDa), sweet potato β-amylase (200 kDa), yeast alcohol

7 dehydrogenase (150 kDa), bovine serum albumin (67 kDa), carbonic anhydrase (29 kDa)

8 and cytochrome c (12.4 kDa).

9


10

11

12

13

14



1

2 Fig. 3 – MALDI-TOF Mass Spectrometry of HML. The MALDI-TOF mass spectrum was  
3 acquired on an AutoFlex. Each sample was dissolved in 0.1% trifluoroacetic acid  
4 (TFA):acetonitrile (2:1 v/v) and mixed with the matrix solution (1:4 v/v). The mixture (1  $\mu$ l)  
5 was put on a stainless steel target and crystallized at room temperature. The spectrum was  
6 measured in linear mode using 20-kV ion acceleration without post-acceleration. The  
7 spectrum was recorded at a detector voltage of 1.65 kV, and was expressed as the averaged  
8 results of at least 300 laser shots.

Table 1 – Purification of HML from 100 g of the fruiting bodies of *Hypsizigus marmoreus*.

| Fraction                          | Total protein (mg) | Total agglutination (titer) <sup>a</sup> | Specific agglutination (titer/mg protein) | Recovery of activity (%) |
|-----------------------------------|--------------------|--|---|--------------------------|
| Extract                           | $1.1 \times 10^3$  | $7.0 \times 10^4$                        | 64  | 100                      |
| CM cation-exchange chromatography | 20                 | $4.1 \times 10^4$                        | $2.0 \times 10^3$                         | 59                       |
| BSM-affinity chromatography       | 0.090              | $1.4 \times 10^3$                        | $1.6 \times 10^4$                         | 2.3                      |
| Gel filtration chromatography     | 0.045              | $8.8 \times 10^2$                        | $2.2 \times 10^4$                         | 1.2                      |

<sup>a</sup> The hemagglutinating activity was determined by a 2-fold serial dilution procedure using rat erythrocytes. Titer was defined as the reciprocal of the end-point dilution exhibiting the hemagglutination.

Table 2 – Partial amino acid sequences of HML.

| Peptide                                       | Sequence      |
|---|---------------|
| Trypsin-1 ( <i>m/z</i> 2,425; calcd, 1,478.6) | LTDPTMIFDSQPN |
| Glu-1 ( <i>m/z</i> 1,358; calcd, 1,283.3)     | DLGSNCGGVFDAE |
| Glu-2 ( <i>m/z</i> 1,331; calcd, 1,238.2)     | AHMTVSFDSDE   |



Table 3 – Inhibition of hemagglutination activity by HML.

| Inhibitors <sup>a</sup>       | Minimum inhibitory concentration <sup>b</sup><br>( $\mu\text{g/ml}$ ) |
|-------------------------------|---|
| Asialo-fetuin                 | 7.81  |
| Thyroglobulin                 | 15.6  |
| Fetuin                        | 15.6  |
| BSM <sup>c</sup>              | 31.3  |
| $\alpha_1$ -acid glycoprotein | 125   |

<sup>a</sup> Glucose, galactose, mannose, L-fucose, arabinose, L-arabinose, raffinose, saccharose, lactose, methyl  $\alpha$ -glucoside, methyl  $\beta$ -glucoside, methyl  $\beta$ -galactoside, methyl  $\alpha$ -mannoside, melibiose, 2-deoxyglucose and 2-deoxyribose did not inhibit at all at concentrations up to 0.4 M. Asialo-BSM, PSM<sup>d</sup> and asialo-PSM did not inhibit at concentrations up to 1 mg/ml.

<sup>b</sup> These are minimum inhibitory concentrations required for inhibition 4 hemagglutination doses of the lectin.

<sup>c</sup> BSM : Bovine submaxillary gland mucin.

<sup>d</sup> PSM : Porcine stomach mucin.