

# Bioconversion of AHX to AOH by resting cells of *Burkholderia contaminans* CH-1

著者	Choi Jae-Hoon, Kikuchi Ayaka, Hirai Hirofumi, Tokuyama Shinji, Kawagishi Hirokazu
journal or publication title	Bioscience, Biotechnology, and Biochemistry
volume	80
page range	2045-2050
year	2016-07-07
出版者	Taylor & Francis
権利	Copyright (C) JSBBA All Rights Reserved.
注記	This is an Accepted Manuscript of an article published by Taylor & Francis Group in Bioscience, Biotechnology, and Biochemistry on 07/07/2016, available online: <a href="http://www.tandfonline.com/10.1080/09168451.2016.1189314">http://www.tandfonline.com/10.1080/09168451.2016.1189314</a> .
URL	<a href="http://hdl.handle.net/10297/10349">http://hdl.handle.net/10297/10349</a>

doi: 10.1080/09168451.2016.1189314

1 **Bioconversion of AHX to AOH by resting cells of *Burkholderia contaminans* CH-1**

2

3 Jae-Hoon Choi<sup>1,2</sup>, Ayaka Kikuchi<sup>2</sup>, Hirofumi Hirai<sup>1,2,3</sup>, Shinji Tokuyama<sup>1,2,\*</sup> and

4 Hirokazu Kawagishi<sup>1,2,3,4,\*</sup>

5

6 <sup>1</sup> *College of Agriculture, Academic Institute, Shizuoka University, Shizuoka, Japan;*

7 <sup>2</sup>*Graduate School of Agriculture, Shizuoka University, Shizuoka, Japan;* <sup>3</sup>*Research*

8 *Institute of Green Science and Technology, Shizuoka University, Shizuoka, Japan;*

9 <sup>4</sup>*Graduate School of Science and Technology, Shizuoka University, Shizuoka, Japan*

10

11 \*Corresponding authors. Email: acstoku@ipc.shizuoka.ac.jp (S.T.);

12 kawagishi.hirokazu@shizuoka.ac.jp (H.K.)

13

1 **Abstract**

2 Fairy rings are zones of stimulated grass growth owing to the interaction between a  
3 fungus and a plant. We previously reported the discovery of two novel plant-growth  
4 regulating compounds that related to forming fairy rings, 2-azahypoxanthine (AHX) and  
5 2-aza-8-oxohypoxanthine (AOH). In this study, a bacterial strain CH-1 was isolated  
6 from an airborne-contaminated nutrient medium containing AHX. The strain converted  
7 AHX to AOH and identified as *Burkholderia contaminans* based on the partial gene  
8 sequence of its 16S rDNA. The quantitative production of AOH by resting cells of the  
9 strain was achieved. Among seven *Burkholderia* species, two bacteria and two yeasts  
10 tested, *B. contaminans* CH-1 showed the highest rate of conversion of AHX to AOH.  
11 By batch system, up to 10.6 mmol AHX was converted to AOH using the resting cells.  
12 The yield of this process reached at 91%.

13

14 **Key words:** bioconversion; *Burkholderia contaminans*; fairy chemicals; fairy ring;  
15 plant growth regulator

16

## 1 **Introduction**

2 Fairy rings are zones of stimulated grass growth due to the interaction between a  
3 fungus and a plant. Since the first scientific article about “fairy rings” in 1675 and  
4 subsequent studies reviewed in Nature in 1884,<sup>1)</sup> this phenomenon had been attributed  
5 to “fairy” before our study in 2010.<sup>2-5)</sup> In that year, we reported that isolation and  
6 identification of the fairy-ring causing principle produced by a fairy-ring forming  
7 fungus *Lepista sordida*, 2-azahypoxanthine (AHX) (Fig. 1).<sup>6)</sup> After that, it was found  
8 out that AHX was absorbed into plants from their roots and then was converted into a  
9 common metabolite, 2-aza-8-oxohypoxanthine (AOH) (Fig. 1).<sup>7)</sup> On the other hand, this  
10 fungus occasionally causes growth-suppressed rings and we purified the suppressor  
11 from the fungus and identified it as imidazole-4-carboxamide (ICA) (Fig. 1A).<sup>8)</sup>  
12 Furthermore, we proved the existence of endogenous AHX and AOH in plants and  
13 found a novel purine pathway producing them in plants (Fig. 1B). We named the three  
14 compounds “fairy chemicals” after the title of the article in Nature that covered our  
15 research about them (Fig. 1A).<sup>9)</sup> Fairy chemicals regulated growth of all the plant tested  
16 and increased the yields of rice, wheat and various other crops in pot and/or field  
17 experiments, suggesting that they might find practical application in agriculture.<sup>10,11)</sup>  
18 For practical use of the compounds, field experiments are necessary and large amounts  
19 of them are required for them. AHX and ICA are chemically synthesized readily. On the  
20 other hand, only method for synthesis of AOH is enzymatic conversion of AHX to it by  
21 xanthine oxidase (XOD) at present (Fig. 2).<sup>12)</sup> However, commercially available XOD  
22 is inappropriate for large-scale preparation of AOH because the enzyme is very  
23 expensive and the enzymatic reaction should be carried out with low concentration of  
24 AHX (Fig. 2A).

1 During searching for better methods to synthesize AOH, we happened to notice that  
2 aqueous solution of AHX was accidentally contaminated by microorganisms and HPLC  
3 analysis of the solution indicated that AHX in the solution was converted to AOH.  
4 Therefore, we tried to isolate the airborne microorganism that produced AOH.

5 Biotransformation to produce value-added substances from natural resources have  
6 been given attention as an alternative tool to replace conventional chemical synthesis of  
7 the substrate.<sup>13,14)</sup> The use of microorganisms as catalysts in the manufacture of  
8 economically important substances is a rapidly developing field of biotechnology.

9 On the other hand, resting cells are non-growing but metabolically active. The use  
10 of resting cells in a bioconversion process would be more advantageous than that of  
11 growing cells.<sup>15)</sup> Production with resting cells often allows easier and more flexible  
12 operation of reaction systems, which results in higher yield.<sup>16,17)</sup>

13 In the present study, we succeeded in isolation of a strain of a bacterium that  
14 converted high concentrations of AHX to AOH with high yield, and resting cells of the  
15 strain were applied to production of AOH from AHX.

16

## 17 **Materials and methods**

18 *Materials.* Strains (*Burkholderia caryophilli* NBRC 13591, *B. capaci* NBRC  
19 14595, *B. caledonica* NBRC 102488, *B. fungorum* NBRC 102489, *B. ferrariae* NBRC  
20 106233, *B. ginsengisoli* NBRC 100965, *B. mimosarum* NBRC 106338, *Bacillus subtilis*  
21 NBRC 13719, *Saccharomyces cerevisiae* NBRC 2377 and *Candida albicans* NBRC  
22 1060) were purchased from the National Institute of Technology and Evaluation (Chiba,  
23 Japan). Xanthine oxidase from butter milk (XOD; 0.35 U/mg powder) was obtained

1 from Oriental Yeast Co. (Tokyo, Japan). All solvents used throughout the experiments  
2 were obtained from Kanto Chemical Co. (Tokyo, Japan).

3

4 *General.*  $^{13}\text{C}$  NMR spectra were recorded on a JEOL lambda spectrometer at  
5 67.5 MHz. HPLC separations were performed with a Jasco Gulliver system using  
6 reverse-phase HPLC columns (Develosil C30-UG-5, Develosil ODS-UG-5, Nomura  
7 chemical Co., Ltd, Japan).

8

9 *Isolation of bacterial strain.* A nutrient solution (for plant culture; 1 mM  
10  $\text{NH}_4\text{NO}_3$ , 0.6 mM  $\text{Na}_2\text{HPO}_4$ , 0.3 mM  $\text{K}_2\text{SO}_4$ , 0.4 mM  $\text{MgCl}_2$ , 0.2 mM  $\text{CaCl}_2$ , 45  $\mu\text{M}$   
11 Fe-ethylenediaminetetraacetic acid, 50  $\mu\text{M}$   $\text{H}_3\text{BO}_3$ , 9  $\mu\text{M}$   $\text{MnSO}_4$ , 0.3  $\mu\text{M}$   $\text{CuSO}_4$ , 0.7  
12  $\mu\text{M}$   $\text{ZnSO}_4$  and 0.1  $\mu\text{M}$   $\text{Na}_2\text{MoO}_4$ ) was used to isolate airborne bacterial strains.  
13 Uncovered petri dish (90 mm) containing a twenty milliliter of AHX solution (0.1 mM  
14 in the nutrient solution) was incubated at room temperature for 3 days. The culture  
15 solution was spread on nutrient agar plates. One colony capable of converting AHX to  
16 AOH was isolated.

17

18 *Culture conditions.* Strain CH-1 was grown in TSB medium containing  
19 polypeptone, 17 g, soy peptone, 3 g, NaCl, 5 g,  $\text{K}_2\text{HPO}_4$ , 2.5 g, and glucose, 2.5 g per  
20 liter. pH of the medium was adjusted to 7.3 with 2 N NaOH. Cultures were inoculated  
21 at an  $\text{OD}_{600}$  of 3.0 and incubated at 30 °C at 180 rpm. Bacterial growth was measured at  
22 600 nm using a Hitachi U-2001 spectrophotometer.

23

1            *Measurement of AOH conversion activity by xanthine oxidase.*        Xanthine  
2    oxidase (XOD, 0.08 unit or 0.5 unit) was dissolved in 10 mM phosphate buffered saline  
3    (PBS, pH 7.4, 900  $\mu$ L). One hundred  $\mu$ L of 0.36 mM (or 11.5 mM) AHX was added to  
4    the buffer containing 0.08 unit (or 0.5 unit) of XOD and reacted in a test tube. After the  
5    reaction, the reaction mixture was analyzed by HPLC (column, C30-UG-5; mobile  
6    phase, 0.02% TFA in H<sub>2</sub>O; flow rate, 0.8 mL/min; wavelength, 254 nm).

7  
8            *Measurement of AOH conversion activity by resting cells.*        The strain was  
9    grown in TSB medium 5 mL (test tube  $\phi$ 18 mm, 30°C, 180 rpm). Cells were harvested  
10   by centrifugation at 11,228 $\times$ g for 5 min. The cell pellet was rinsed and resuspended in  
11   10 mM PBS (900  $\mu$ L). Various concentrations of AHX (100  $\mu$ L) were added to the  
12   suspension and reacted in a test tube. After the reaction, the reaction mixture was heated  
13   (98°C, 5 min) and centrifuged at 11,228 $\times$ g for 5 min. Concentration of AHX and AOH  
14   in the supernatant was determined by HPLC (column, C30-UG-5; mobile phase, 0.02%  
15   TFA in H<sub>2</sub>O; flow rate, 0.8 mL/min; wavelength, 254 nm).

16        For a conversion of AHX at a large scale, the strain was precultured in TSB  
17   medium (5 mL, test tube  $\phi$ 18 mm, 30°C, 180 rpm, 16 h) and then the precultured cells  
18   were grown in 100 mL of TSB medium in 500 mL-Erlenmeyer flask (30°C, 180 rpm, 6  
19   h). The cells were harvested by centrifugation at 15,652 $\times$ g for 5 min. The obtained cell  
20   pellet was rinsed and resuspended in 10 mM PBS (20 mL). Various concentration of  
21   AHX (180 mL) were added and reacted in the flask (40°C, 80 rpm). Concentrations of  
22   AHX and AOH were determined by HPLC.

23        In order to examine whether further metabolism of AOH occurs by resting cells of  
24   CH-1, the strain was grown in 5 mL of TSB medium (test tube  $\phi$ 18 mm, 30°C, 180 rpm).

1 Cells are harvested by centrifugation at 11,228×g for 5 min. The obtained cell pellet  
2 was rinsed and resuspended in 10 mM PBS (900 μL). One hundred μL of 0.46 mM  
3 AOH was added to the suspension and reacted in a test tube. After the reaction, a  
4 concentration of AOH was determined by HPLC.

5  
6 *Purification of AOH.* AOH produced by the resting cell method was purified  
7 as follows. After the reaction at the 500 mL scale, the reaction mixture was heated at  
8 98°C for 5 min, and the cells were removed by centrifugation (39,120×g, 10 min). Then,  
9 the solution was concentrated 7-fold by an evaporator and AOH was crystallized at 4°C  
10 for 24 h and the obtained crystals were collected by filtration and dried in a vacuum  
11 desiccator.

12  
13 *Measurement of xanthine oxidase activity of resting cells of strain CH-1.* The  
14 strain was grown in 5 mL of TSB medium (test tube φ18 mm, 30°C, 180 rpm). Cells are  
15 harvested by centrifugation at 11,228×g for 5 min. The obtained cell pellet was washed  
16 and resuspended in 50 mM Tris-HCl buffer (pH 7.5, 1.45 mL). Fifty μL of 10 mM  
17 xanthine was added to the suspension and the reaction mixture was incubated in a test  
18 tube. After incubation, the reaction mixture was heated at 98°C for 5 min and was  
19 centrifuged at 11,228×g for 5 min. Concentration of xanthine and uric acid in the  
20 supernatant was determined by HPLC (column, ODS-UG-5; mobile phase, 10 mM  
21 potassium phosphate buffer, pH 5.5; flow rate, 1 mL/min; wavelength, 260 nm).

22  
23 *Bacterial identification by sequence analysis of 16S rDNA.* To extract the  
24 genomic DNA, bacterial pellet was suspended in 0.2 mL of Insta Gene Matrix (Bio-Rad,



1 USA), incubated at 56°C for 30 min and then heated at 100°C for 8 min. After heating,  
2 the supernatant was used for the PCR reaction. Bacterial 16S rDNAs were amplified by  
3 using two universal primers, 9 F (5'-GAGTTTGATCCTGGCTCAG-3') and 1541 R  
4 (5'- AAGGAGGTGATCCAGCC-3'). Polymerase chain reaction was performed using  
5 a kit with Hot Star tag DNA polymerase (Qiagen). The PCR program used for  
6 amplification was as follows: 95 °C for 15 min, followed by 25 cycles consisting of  
7 94°C for 30 sec, 50°C for 1 min, and 72°C for 1 min, and a single final extension step  
8 consisting of 72°C for 10 min. The sequencing analysis of amplified products was  
9 performed with four universal primers, 9 F (5'-GAGTTTGATCCTGGCTCAG-3'), 785  
10 F (5'-GGATTAGATACCCTGGTAGTC-3'), 926 R  
11 (5'-CCGTCAATTCCTTTGAGTTT-3') and 1541 R (5'-  
12 AAGGAGGTGATCCAGCC-3') by Fasmac Co., Ltd.

13

## 14 **Results**

### 15 *Isolation and identification of AOH-forming bacterium*

16 Since AHX in the nutrition medium contaminated by airborne microorganisms was  
17 converted to AOH, we were triggered to investigate airborne microorganisms for AOH  
18 production from AHX. An uncovered petri dish containing AHX solution was left alone  
19 at room temperature for 3 days. Conversion of AHX to AOH in the solution was  
20 analyzed by HPLC and part of AHX was converted to AOH (Fig. 3). Therefore, we  
21 tried to isolate the microorganism that converted AHX to AOH. The culture solution  
22 was spread on nutrient agar plates. As a result, one colony capable of converting AHX  
23 to AOH was isolated and we named the strain CH-1.

1 The partial sequence (1423 bp) of its 16S rDNA showed that the strain CH-1 is  
2 *Burkholderia contaminans* with similarity of 99.9% to the type strain (Table 1).

3

#### 4 *Bioconversion of AOH by resting cells of Burkholderia contaminans CH-1*

5 After cultivation, the isolated *B. contaminans* CH-1 cells were harvested via  
6 centrifugation, washed, and then resuspended in the buffer containing AHX. AOH yield  
7 was greatly improved with the increase in reaction time with the resting cells (Fig. 4).  
8 Fig. 4A shows that AHX (initial concentration of 0.5 mM) was completely  
9 biotransformed to AOH by the cells at 24 h. The produced AOH was stable with the  
10 cells without further degradation even after 98 h of reaction (Fig. 4C).

11

#### 12 *Bioconversion of AOH by resting cells of various microorganisms*

13 We compared the ability of the conversion of the obtained strain, CH-1, with other  
14 *Burkholderia* species and other microorganisms. For the comparison, reaction time was  
15 set to be 24 h, because the conversion with CH-1 almost reached maximum at 24 h.  
16 Seven *Burkholderia* species (*B. caryophilli* NBRC 13591, *B. capaci* NBRC 14595, *B.*  
17 *caledonica* NBRC 102488, *B. fungorum* NBRC 102489, *B. ferrariae* NBRC 106233, *B.*  
18 *ginsengisoli* NBRC 100965, and *B. mimosarum* NBRC 106338), other two bacteria  
19 (*Escherichia coli* JM 109 and *Bacillus subtilis* NBRC 13719), and two yeasts  
20 (*Saccharomyces cerevisiae* NBRC 2377 and *Candida albicans* NBRC 1060) were  
21 tested (Fig. 5A). Among them, only two species were able to produce AOH from AHX  
22 quantitatively like CH-1 (Fig. 5A). As shown in Fig. 5B-D, when the reaction time was  
23 shortened to 6 h from 24 h, AOH ratio between the contents of AHX and AOH  
24 produced by CH-1, *B. cepacia*, and *B. fungorum* were 89%, 70%, and 16%, respectively.

1 As a consequence, *B. contaminans* CH-1 exhibited the highest converting ability of  
2 AHX to AOH among all the strains tested.

3

#### 4 *Production of AOH by the resting cells of B. contaminans CH-1*

5 The results mentioned above indicated that AOH could be produced from AHX by  
6 the resting cell method and that 0.5 mM AHX was completely converted to AOH. To  
7 confirm the result at higher concentrations of AHX, the resting cell method was carried  
8 out at a larger scale. 14.6 mM AHX was completely converted to AOH within 24 h of  
9 incubation (Fig. 4B). AOH was then purified by centrifugation and crystallization from  
10 reaction mixture of 11.6 mmol AHX, and we obtained 1.63 g (10.6 mmol) of purified  
11 AOH in 91% yield. The purified AOH showed the same retention time in HPLC and the  
12 same signals in <sup>13</sup>C NMR as authentic AOH (Fig. 6).

13

#### 14 **Discussion**

15 AHX and AOH are useful chemicals with potential applications in a wide range of  
16 fields.<sup>10,11)</sup> A production of AOH from AHX has been reported by xanthine oxidase.<sup>7)</sup>  
17 However, this production method is insufficient for industrial use owing to several  
18 disadvantages such as low yields and slow reaction rates.<sup>12)</sup>

19 In the current study, *B. contaminans* CH-1 with high AOH-producing ability was  
20 screened and identified (Fig. 3). *B. contaminans* is a member of the *Burkholderia*  
21 *cepacia* complex, a group of closely related gram-negative.<sup>18-20)</sup> Characterization of this  
22 strain showed that it inhibited the growth of a broad range of fungal pathogens.<sup>21-23)</sup> It  
23 has been reported about a degradation of 3-nitrotyrosine, 2-naphthoate and lactose by  
24 *Burkholderia* sp.<sup>24-26)</sup>

1        Biotransformation of resting cells has some advantages such as short reaction time,  
2 non-sterile reaction conditions and convenient product separation. In the enzymatic  
3 reaction of AOH from AHX by XOD, high concentration of AHX strongly inhibited the  
4 activity of XOD (Fig. 2B). On the other hand, the resting cells almost completely  
5 converted AHX to AOH. These results indicated that the cells produced more active  
6 and/or more stable enzyme(s) or larger amounts of the enzyme(s) than the commercially  
7 available XOD. Considering the diverse environment of biotransformation between the  
8 resting cells and XOD, the resting cell method for AOH production is attractive in  
9 aspects of bioconversion time, substrate concentration and purification. To investigate  
10 whether the enzyme that produced AOH from AHX in the resting cells of CH-1 is XOD  
11 or not, we used xanthine as the substrate and xanthine was completely converted into  
12 uric acid within 4 h of incubation (Fig 7). This result suggested that AOH might be  
13 produced by XOD in the resting cells. We are now trying to purify the enzyme that  
14 catalyzes the reaction from AHX to AOH, because the enzyme would be useful for  
15 further improvement of the present biochemical method of AOH production and for  
16 investigation of the biosynthetic route from AHX to AOH in microorganisms and/or  
17 plants. A production level of AOH (10.6 mmol) was achieved using the bioconversion  
18 that successively used resting cells, which is a novel procedure for large scale  
19 production of AOH. Highly purified AOH is easily obtained through a simple  
20 purification procedure of centrifugation and crystallization. This suggests that AOH  
21 may be easily produced by our method to practical use in agriculture.

22

23 **Author contributions**

1 J.-H.C., S.T., and H.K. designed the experiments. J.-H.C. and A.K. performed the  
2 experiments. H.H., S.T. and H.K. assisted with experiments and contributed to  
3 discussions. J.-H.C. and H.K. wrote the manuscript.

#### 4 5 **Acknowledgments**

6 We would like to thank K., Okamoto (Ushio ChemiX Co. Ltd.) for providing  
7 AHX.

#### 8 9 **Funding**

10 This work was partially supported by Science and Technology Research Promotion  
11 Program for Agriculture, Forestry, Fisheries and Food Industry from MAFF.

#### 12 13 **References**

- 14 [1] Evershed H. Fairy rings. *Nature*. 1884;29:384–385.
- 15 [2] Couch HB. Diseases of turfgrasses, Krieger, Malabar, 1995, pp. 181–186.
- 16 [3] Smith JD, Jackson N, Woolhouse AR. Fungal diseases of amenity turf grasses, E. &  
17 F. N. Spon, London, 1989, pp. 339–353.
- 18 [4] Shantz HL, Piemeisel RL. Fungus fairy rings in eastern Colorado and their effect on  
19 vegetation. *J. Agric. Res.* 1917;11:191–245.
- 20 [5] Ramsbottom J. Rate of growth of fungus rings. *Nature*. 1926;117:158–159.
- 21 [6] Choi J-H, Fushimi K, Abe N, et al. Disclosure of the “fairy” of fairy-ring-forming  
22 fungus *Lepista sordida*. *ChemBioChem*. 2010;11:1373–1377.
- 23 [7] Choi J-H, Ohnishi T, Yamakawa Y, et al. The source of “fairy rings”:  
24 2-azahypoxanthine and its metabolite found in a novel purine metabolic pathway in

- 1 plants. *Angew. Chem. Int. Ed.* 2014;53:1552–1555.
- 2 [8] Choi J-H, Abe N, Fushimi K, et al. Plant-growth regulator,  
3 imidazole-4-carboxamide produced by fairy-ring forming fungus *Lepista sordida*. *J.*  
4 *Agric. Food Chem.* 2010;58:9956–9959.
- 5 [9] Mitchinson A. Fairy chemicals. *Nature.* 2014;505:298.
- 6 [10] Tobina H, Choi J-H, Asai T, et al. 2-Azahypoxanthine and  
7 imidazole-4-carboxamide produced by the fairy-ring-forming fungus increase yields  
8 of wheat. *Field Crop Res.* 2014;162:6–11.
- 9 [11] Asai T, Choi J-H, Ikka T, et al. Effect of 2-azahypoxanthine (AHX) produced by  
10 the fairy-ring-forming fungus on the growth and the grain yield of rice. *Jpn. Agric.*  
11 *Res. Quart.* 2015;49:45–49.
- 12 [12] Ikeuchi K, Fujii R, Sugiyama S, et al. Practical synthesis of natural plant-growth  
13 regulator 2-azahypoxanthine, its derivatives, and biotin-labeled probes. *Org. Biomol.*  
14 *Chem.* 2014;12:3813–3815.
- 15 [13] Shimoni E, Baasov T, Ravid U, et al. Biotransformations of propenylbenzenes by  
16 an *Arthrobacter* sp. and its *t*-anethole blocked mutants. *J. Biotechnol.*  
17 2003;105:61–70.
- 18 [14] Tripathi U, Rao RS, Ravishankar GA. Biotransformation of phenylpropanoid  
19 compounds to vanilla flavor metabolites in cultures of *Haematococcus pluvialis*.  
20 *Process Biochem.* 2002;38:419–426.
- 21 [15] Carvalho F, Marques MPC, de Carvalho CCCR, et al. Sitosterol bioconversion  
22 with resting cells in liquid polymer based systems. *Bioresour. Technol.*  
23 2009;100:4050–4053.
- 24 [16] Wang Z. The potential of cloud point system as a novel twophase partitioning

- 1 system for biotransformation. *Appl. Microbiol. Biotechnol.* 2007;75:1–10.
- 2 [17] Saxena RK, Anand P, Saran S, et al. Microbial production of 1,3-propanediol:  
3 recent developments and emerging opportunities. *Biotechnol. Adv.*  
4 2009;27:895–913.
- 5 [18] Chiarini L, Bevivino A, Dalmastrì C, et al. *Burkholderia cepacia* complex species:  
6 health hazards and biotechnological potential. *Trends Microbiol.* 2006;14:277–286.
- 7 [19] Vanlaere E, Lipuma JJ, Baldwin A, et al. *Burkholderia latens* sp. nov.,  
8 *Burkholderia diffusa* sp. nov., *Burkholderia arboris* sp. nov., *Burkholderia*  
9 *seminalis* sp. nov. and *Burkholderia metallica* sp. nov., novel species within the  
10 *Burkholderia cepacia* complex. *Int. J. Syst. Evol. Microbiol.* 2008;58:1580–1590.
- 11 [20] Vanlaere E, Baldwin A, Gevers D, et al. Taxon K, a complex within the  
12 *Burkholderia cepacia* complex comprises at least two novel species: *Burkholderia*  
13 *contaminans* sp. nov. and *Burkholderia lata* sp. nov., *Int. J. Syst. Evol. Microbiol.*  
14 2009;59:102–111.
- 15 [21] Lu S-E, Woolfolk S, Caceres J. Isolation and identification and genetic analysis of  
16 rhizobacteria antagonistic to plant soilborne fungal pathogens. *Phytopathology.*  
17 2005;95:62–63.
- 18 [22] Gu GY, Smith L, Wang N, et al. Biosynthesis of an antifungal oligopeptide in  
19 *Burkholderia contaminans* strain MS14. *Biochem. Biophys. Res. Commun.*  
20 2009;380:328–332.
- 21 [23] Lu S-E, Novak J, Austin FW, et al. Occidiofungin, a unique antifungal  
22 glycopeptide produced by a strain of *Burkholderia contaminans*. *Biochemistry.*  
23 2009;48:8312–8321.
- 24 [24] Nishino SF, Spain JC. Biodegradation of 3-nitrotyrosine by resting cells of

- 1        *Burkholderia* sp. strain JS165 and *Variovorax paradoxus* JS171. Appl. Environ.  
2        Microbiol. 2006;72:1040–1044.
- 3        [25] Morawski B, Richard WE, John TR, et al. 2-Naphthoate catabolic pathway in  
4        *Burkholderia* strain JT1500. J. Bacteriol. 1997;179:115–121.
- 5        [26] Murakami H, Seko A, Azumi M, et al. Microbial conversion of lactose to  
6        lactobionic acid by resting cells of *Burkholderia cepacia* No. 24. J. Appl. Glycosci.  
7        2006;53:7–11.



## Figure captions

Fig. 1. Structures of fairy chemicals (A) and purine metabolic pathway in animals, plants and microorganisms (B).

Note: **1** : 2-azahypoxanthine (AHX); **2** : 2-aza-8-oxohypoxanthine (AOH); **3** : imidazole-4-carboxamide (ICA); **4** : 5-aminoimidazole-4-carboxamide (AICA); **5** : AICA ribotide; **6**:5-formyl AICAR; XOD: xanthine oxidase. The novel pathway in plants is in the framed rectangle.

Fig. 2. Time course of production of AOH from AHX by xanthine oxidase.

Note: Symbols, ● and ○, indicate AHX and AOH, respectively. Initial concentrations of AHX were 0.36 mM (A) and 11.5 mM (B), respectively.

Fig. 3. HPLC profile of authentic AHX and AOH, and culture broth of CH-1

Notes: A: authentic AHX; B: culture broth of CH-1; C: authentic AOH. Column: C30-UG-5; mobile phase: 0.02% TFA in H<sub>2</sub>O; flow rate: 0.8 mL/min; wavelength: 254 nm.

Fig. 4. Time course of AOH production (A, B) and AOH degradation (C) by resting cells of CH-1.

Note: Symbols, ● and ○, indicate AHX and AOH, respectively. Initial concentrations of AHX were 0.5 mM (A) and 14.6 mM (B), respectively.

Fig. 5. Concentration ratio of AHX and AOH in the reaction by resting cells of various microorganisms.

Notes: A: AHX (0.5 mM) was incubated for 24 h by resting cells of various microorganisms; conversion rate at each reaction time by resting cells of CH-1 (B), *B. cepacia* NBRC 14595 (C) and *B. fungorum* NBRC 102489 (D). Gray and black columns indicate AHX and AOH contents, respectively.

Fig. 6. Identification of the purified product.

Notes: HPLC chromatograms of AOH (A). Authentic AOH (above), purified AOH in this study (below). Column: C30-UG-5; mobile phase: 0.02% TFA in H<sub>2</sub>O; flow rate: 0.8 mL/min; wavelength: 254 nm. <sup>13</sup>C NMR spectra of AOH in DMSO-d<sub>6</sub> (B). Authentic AOH (above), purified AOH in this study (below).

Fig. 7. Time course of production of uric acid from xanthine by resting cells of CH-1

Note: Symbols, ● and ○, indicate xanthine and uric acid, respectively.




Fig. 1





Fig. 2




Fig. 3




Fig. 4




Fig. 5





Fig. 6






Fig. 7

Table 1. The 16S rDNA similarity between strain CH-1 and other *Burkholderia* genus

Number	Strains	Similarity (%) (Length)
1	<i>Burkholderia contaminans</i> LMG 23361 <sup>T</sup>	99.9 ( 1421/1423 )
2	<i>Burkholderia lata</i> 383 <sup>T</sup>	99.7 ( 1419/1423 )
3	<i>Burkholderia arboris</i> R-24201 <sup>T</sup>	99.6 ( 1418/1423 )
4	<i>Burkholderia cepacia</i> NBRC14074 <sup>T</sup>	99.4 ( 1413/1421 )
5	<i>Burkholderia cenocepacia</i> LMG 16656 <sup>T</sup>	99.2 ( 1410/1421 )