

# Cytotoxic compounds against cancer cells from *Bombyx mori* inoculated with *Cordyceps militaris*

|                              |  |
|------------------------------|--|
| 著者                           | Qiu Weitao, Wu Jing, Choi Jae-Hoon, Hirai Hirofumi, Nishida Hiroshi, Kawagishi Hirokazu  |
| journal or publication title | Bioscience, Biotechnology and Biochemistry   |
| volume                       | 81   |
| number                       | 6  |
| page range                   | 1224-1226  |
| year                         | 2017-02-09   |
| 出版者                          | Taylor & Francis   |
| 権利                           | (C) JSBBA All Rights Reserved.<br>This is an electronic version of an article published in Bioscience, Biotechnology, and Biochemistry [Volume 81, No.6, pp1224-1226, 2017]. Bioscience, Biotechnology, and Biochemistry is available online at:<br><a href="http://www.tandfonline.com/10.1080/09168451.2017.1289075">www.tandfonline.com/10.1080/09168451.2017.1289075</a> . |
| URL                          | <a href="http://hdl.handle.net/10297/00024648">http://hdl.handle.net/10297/00024648</a>  |

doi: 10.1080/09168451.2017.1289075

**Cytotoxic compounds against cancer cells from *Bombyx mori* inoculated with *Cordyceps militaris***

|                                    |   |
|------------------------------------|---|
| Journal:                           | <i>Bioscience, Biotechnology, and Biochemistry</i>  |
| Manuscript ID                      | BBB-160819.R1   |
| Manuscript Type:                   | Note  |
| Date Submitted by the Author:      | 16-Jan-2017   |
| Complete List of Authors:          | Qiu, Weitao ; Shizuoka University, Graduate School of Science and Technology<br>Wu, Jing ; Shizuoka University, Research Institute of Green Science and Technology<br>Choi, jaehoon; Shizuoka University, Research Institute of Green Science and Technology<br>HIRAI, Hirofumi; Shizuoka University, Agriculture<br>Nishida, Hiroshi; Niigata University of Pharmacy and Applied Life Sciences, Faculty of Applied Life Sciences<br>KAWAGISHI, Hirokazu; Shizuoka University, Research Institute of Green Science and Technology |
| Keywords:                          | Cytotoxic compound, <i>Bombyx mori</i> , <i>Cordyceps militaris</i>   |
| Subject Categories:                | Food & Nutrition Science  |
| Classification of Research Fields: | II - 2) Isolation and Structure Determination of Bioactive Substances < II. Organic Chemistry and Chemistry of Natural Products, V - 2) Chemistry and Biochemistry < V. Foods   |
|                                    |   |

1 **Cytotoxic compounds against cancer cells from *Bombyx mori* inoculated with**  
2 ***Cordyceps militaris***

3

4 Weitao Qiu<sup>1</sup>, Jing Wu<sup>2</sup>, Jae-Hoon Choi<sup>2,3</sup>, Hirofumi Hirai<sup>2,3</sup>, Hiroshi Nishida<sup>4,\*</sup>,  
5 Hirokazu Kawagishi<sup>1,2,3,\*</sup>

6

7 <sup>1</sup> Graduate School of Science and Technology, Shizuoka University, Shizuoka, Japan;

8 <sup>2</sup> Research Institute of Green Science and Technology, Shizuoka University, Shizuoka,  
9 Japan;

10 <sup>3</sup> College of Agriculture, Academic Institute, Shizuoka University, Shizuoka, Japan

11 <sup>4</sup> Faculty of Applied Life Sciences, Niigata University of Pharmacy and Applied Life  
12 Sciences, Niigata, Japan

13

14

15

16 \*Corresponding author.

17 E-mail: hnishida@nupals.ac.jp (H Nishida)

18 E-mail: kawagishi.hirokazu@shizuoka.ac.jp (H Kawagishi)

1 **Abstract**

2       Two compounds, 3'-deoxyinosine and **cordycepin**, were isolated from *Bombyx*  
3 *mori* inoculated with *Cordyceps militaris*. In the bioassay examining cytotoxicity  
4 against cancer cells, both compounds showed toxicity against A549, PANC-1 and  
5 MCF-7 cancer cells.

6

7 **Key words:** Cytotoxic compound; *Bombyx mori*; Structural identification; *Cordyceps*  
8 *militaris*

For Peer Review

1 The genus *Cordyceps* belongs to Ascomycota phylum, Sordariomycetes class,  
2 Hypocreales order, Clavicipitaceae family. *Cordyceps* fungi have parasitic nature on  
3 insects' larvae and pupae, or even on adult insects.<sup>1)</sup> Many species of *Cordyceps* have  
4 been used as traditional Chinese medicines from ancient times. In the studies about  
5 *Cordyceps* sp., a lot of bioactivities have been reported, such as immunomodulatory  
6 activity, antioxidant activity, antitumour activity and cancer cell cytotoxicity.<sup>2-17)</sup>

7 *Cordyceps militaris* is a valuable specie of the genus *Cordyceps*. It parasitizes  
8 lepidopteran pupa and forms fruiting bodies. Natural *C. militaris* is very expensive  
9 because of its less production. However, recently, artificial cultivation of the fruiting  
10 bodies was succeeded and since then the fungus has become a widely-used functional  
11 food and a subject of research. Although several biological and chemical studies about  
12 the mycelia and the fruiting bodies have been reported, there are few studies about its  
13 host, lepidopteran larvae infested with the fungus.<sup>4,7,8,10,12,13,15-20)</sup> In the screening  
14 experiments, we found the cytotoxicity of the crude extracts of *Bombyx mori* inoculated  
15 with *C. militaris* and started this study.

16 Here, we describe the isolation and identification of two cytotoxic compounds  
17 against cancer cells from the infested *B. mori*.

18 Larvae bodies of the infested *B. mori* were extracted with EtOH and then with  
19 acetone. After the solutions were combined and dried up under reduced pressure, the  
20 dried materials were extracted with *n*-hexane, EtOAc and MeOH, respectively. The  
21 EtOAc soluble part was fractionated by repeated chromatography. As a result, two  
22 compounds (**1** and **2**) were purified (Fig. 1).

23 Compound **1** was identified to be  
24 **(2*R*,3*R*,5*S*)-2-(6-aminopurin-9-yl)-5-(hydroxymethyl)oxolan-3-ol** (cordycepin), which  
25 was isolated from the cultured mycelia of *C. militaris* in 1950.<sup>21)</sup> Cordycepin is famous  
26 for its various bioactivities, such as antioxidative activity and cancer cell  
27 cytotoxicity.<sup>9,14,16,17,22,23)</sup> **All the data of **1** including the CD spectrum were identical with**  
28 **those of commercially available cordycepin.**

29 Compound **2** was identified as 3'-deoxyinosine that has been isolated from okra  
30 (*Abelmoschus esculentus*)<sup>24)</sup> and it was previously reported that this compound showed  
31 antiprotozoal activities.<sup>25,26)</sup> However, there is no other report of isolation of **2** from  
32 natural sources.

1 Cytotoxicity of compounds **1** and **2** against cancer cells (A549, PANC-1 and  
2 MCF-7) was tested (Fig. 2). Compounds **1** and **2** showed dose-dependent cytotoxicity  
3 against PANC-1 and MCF-7 cancer cells. On the other hand, **1** also showed  
4 dose-dependent cytotoxicity against A549 cell, but the cytotoxicity of **2** against the cells  
5 was weak and there was no dose-dependency. Both the two compounds showed the  
6 strongest cytotoxicity against MCF-7 cells at 30  $\mu$ M. In the previous studies, **1** has  
7 shown cytotoxicity against A549 and MCF-7 cancer cells.<sup>16,17,22,23</sup> The cytotoxicity of **1**  
8 against PANC-1 cancer cell is reported here for the first time. This is also the first time  
9 for reporting the cytotoxicity against cancer cells of **2**.

10 <sup>1</sup>H-NMR spectra (one-and two-dimensional) were recorded on a Jeol lambda-500  
11 spectrometer (Jeol Ltd., Tokyo, Japan) at 500 MHz, while <sup>13</sup>C-NMR spectra were  
12 recorded by the same instrument at 125 MHz. HRESIMS data were measured by a  
13 JMS-T100LC mass spectrometer (Jeol Ltd., Tokyo, Japan). HPLC separation was  
14 performed with a Jasco Gulliver system (Jasco Co., Tokyo, Japan) using a reverse-phase  
15 HPLC column (Cosmosil PBr,  $\phi$  20 $\times$ 250 mm, Nacalai tesque, Kyoto, Japan). C18  
16 cartridges (Nihon Waters K.K., Tokyo, Japan) were used in the pro-processing of the  
17 samples. Silica gel plate (TLC Silica gel 60 F<sub>254</sub>, Merck KGaA, Darmstadt, Germany)  
18 and silica gel 60N (Kanto Chemical Co.,Inc., Tokyo, Japan) were used for analytical  
19 TLC and for flash column chromatography, respectively.

20 Larvae of *B. mori*, which was inoculated with *C. militaris* by injection and then  
21 fruiting bodies of the fungus formed from the larvae. The larvae with the fruiting bodies  
22 were purchased from Nichihara Research & Development Laboratories, Inc. in 2014.  
23 Human adenocarcinoma A549, human pancreatic carcinoma PANC-1 and breast  
24 adenocarcinoma MCF-7 cells were obtained from ATCC cell line (VA, USA), and  
25 maintained in Dulbecco's modified eagle's medium (DMEM, Sigma-Aldrich)  
26 supplemented with 10% fetal bovine serum, 100  $\mu$ g/ml streptomycin, and 100 units/ml  
27 penicillin (all purchased from Invitrogen, CA, USA). All cultures were kept in a  
28 humidified atmosphere with 5% CO<sub>2</sub> at 37°C.

29 The larvae with the fruiting bodies (16.4 kg) was divided into the bodies of *B.*  
30 *mori* (14.3 kg) and the fruiting bodies (2.10 kg). The insect bodies were crushed and  
31 extracted with EtOH (40.0 L, 3 times) and then with acetone (20.0 L, 3 times). After the

1 solutions were combined and dried up under reduced pressure, the dried materials were  
2 extracted with *n*-hexane, EtOAc and MeOH for 2 times (2.0 L), respectively. The  
3 EtOAc-soluble part (45.0 g) was fractionated by silica gel flash column chromatography  
4 (99/1, 95/5, 80/20, 70/30, 50/50, 30/70 CH<sub>2</sub>Cl<sub>2</sub>/acetone; 99/1, 95/5, 80/20, 70/30, 50/50,  
5 30/70 CH<sub>2</sub>Cl<sub>2</sub>/MeOH; 2.0 L each) to obtain 24 fractions (fractions 1 to 24). Fraction 21  
6 (726 mg) was fractionated by C18 cartridges (50% MeOH, MeOH; 100 mL, 2 times,  
7 respectively) to obtain 3 fractions (fractions 21-1 to 21-3). Fraction 21-1 (520 mg) was  
8 further fractionated by reverse-phase HPLC (Cosmosil PBr,  $\phi$  20 $\times$ 250 mm, UV 250 nm,  
9 30% MeOH) to give 23 fractions (fractions 21-1-1 to 21-1-23), and fraction 21-1-17  
10 was compound **1** (41.0 mg). Fraction 21-1-11 (5.3 mg) was separated by reverse-phase  
11 HPLC (Cosmosil PBr,  $\phi$  20 $\times$ 250 mm, UV 250 nm, 20% MeOH) to give compound **2**  
12 (2.2 mg).

13 Compound **2**: white amorphous; ESIMS  $m/z$  275 [M+Na]<sup>+</sup>; <sup>1</sup>H-NMR (in  
14 CD<sub>3</sub>OD)  $\delta_{\text{H}}$ : 8.38 (s, H-2), 8.04 (s, H-8), 5.99 (d,  $J=2.1$  Hz, H-1'), 4.65 (m, H-2'), 4.51  
15 (m, H-4'), 3.90 (dd,  $J=12.2, 2.7$  Hz, H-5'a), 3.67 (dd,  $J=12.2, 3.7$  Hz, H-5'b), 2.33 (m,  
16 H-3'a), 2.02 (m, H-3'b), <sup>13</sup>C-NMR  $\delta_{\text{C}}$ : 159.4 (C-6), 149.4 (C-4), 147.1 (C-2), 140.3  
17 (C-8), 125.8 (C-5), 93.4 (C-1'), 82.8 (C-4'), 77.2 (C-2'), 63.9 (C-5'), 34.5 (C-3').

18 Compounds **1** and **2** were tested the cell viability against A549, PANC-1 and  
19 MCF-7 cancer cells through MTT assay.<sup>27)</sup> Cells were seeded at  $1 \times 10^4$  cells/cm<sup>2</sup> in  
20 96-well plates and cultured for 24 h. Cells were incubated with compounds **1** and **2** (1,  
21 10, 30  $\mu$ M, respectively), sulforaphane (SFN, 30  $\mu$ M) as indicated concentrations for 48  
22 h. Sulforaphane (SFN; 30  $\mu$ M) was used as positive control. After the incubation, the  
23 growth medium was removed and the cells were given 100  $\mu$ L of 0.05% MTT solution,  
24 then incubated for 4 h. After the cells were incubated with 100  $\mu$ L of lysis buffer [20%  
25 SDS, 50% *N,N*-dimethyl formamide (DMF), pH 4.7], absorbance was measured by a  
26 microplate reader (Bio-Rad Laboratories, CA, USA) at 595 nm. All incubations were  
27 carried out at 37°C in 5% CO<sub>2</sub>. All data are shown as means  $\pm$  SD. Differences among  
28 the all groups were evaluated using a 1-way analysis of variance (ANOVA) followed by  
29 the Dunnett. A *P* value less than 0.05 and 0.01 were considered statistically significant.

30

31 **Author contribution**

1 Hirokazu Kawagishi and Hiroshi Nishida designed the experiments. Weitao Qiu  
2 performed the experiments. Jing Wu, Jea-Hoon Choi, Hirofumi Hirai, Hiroshi Nishida  
3 and Hirokazu Kawagishi contributed to discussions. Weitao Qiu, Hiroshi Nishida and  
4 Hirokazu Kawagishi wrote the manuscript.

#### 6 **Disclosure statements**

7 No potential conflict of interest was reported by authors.

#### 9 **References**

- 10 [1] Xiao J, Qi Y, Xiong Q. Nucleosides, a valuable chemical marker for quality control  
11 intraditional Chinese medicine *Cordyceps*. Recent. Pat. Biotechnol.  
12 2013;7:153-166.
- 13 [2] Kuo Y, Weng S, Chou C, Chang T, Tsai W. Activation and proliferation signals in  
14 primary human T lymphocytes inhibited by ergosterol peroxide isolated from  
15 *Cordyceps cadae*. Br. J. Pharmacol. 2003;40:895-906.
- 16 [3] Kuo C, Chen C, Lin C, Jan M, Huang R, Luo Y, Chuang W, Sheu C, Lin Y.  
17 Abrogation of streptococcal pyrogenic exotoxin B-mediated suppression of  
18 phagocytosis in U937 cells by *Cordyceps sinensis* mycelium via production of  
19 cytokines. Food. Chem. Toxicol. 2007;45:278-285.
- 20 [4] Liu J, Feng C, Li X, Chang M, Meng J, Xu L. Immunomodulatory and  
21 antioxidative activity of *Cordyceps militaris* polysaccharides in mice. Int. J. Biol.  
22 Macromol. 2016;86:594-598.
- 23 [5] Li S, Zhao K, Ji Z, Song Z, Dong T, Lo C, Cheung J, Zhu S, Tsim K. A  
24 polysaccharide isolated from *Cordyceps sinensis*, a traditional Chinese medicine,  
25 protects PC12 cells against hydrogen peroxide-induced injury. Life Sci.  
26 2003;73:2503-2513.
- 27 [6] Olatunji OJ, Feng Y, Olantunji OO, Tang J, Ouyang Z, Su Z, Wang D, Yu X.  
28 Neuroprotective effects of adenosine isolated from *Cordyceps cicadae* against  
29 oxidative and ER stress damages induced by glutamate in PC12 cells. Environ.



- 1 Toxicol. Phar. 2016;44:53-61.
- 2 [7] Yu R, Yang W, Song L, Yan C, Zhang Z, Zhao Y. Structural characterization and  
3 antioxidant activity of a polysaccharide from the fruiting bodies of cultured  
4 *Cordyceps militaris*. Carbohydr. Polym. 2007;70:430-436.
- 5 [8] Zhu Z, Liu F, Gao H, Sun H, Meng M, Zhang Y. Synthesis, characterization and  
6 antioxidant activity of selenium polysaccharide from *Cordyceps militaris*. Int. J.  
7 Biol. Macromol. 2016;93:1090-1099.
- 8 [9] Olatunji OJ, Feng Y, Olatunji OO, Tang J, Ouyang Z, Su Z. Cordycepin protects  
9 PC12 cells against 6-hydroxydopamine induced neurotoxicity via its antioxidant  
10 properties. Biomed. Pharmacother. 2016;81:7-14.
- 11 [10] Yoo H, Shin J, Cho J, Son C, Lee Y, Park S, Cho C. Effects of *Cordyceps militaris*  
12 extract on angiogenesis and tumor growth. Acta Pharmacol. Sin. 2004;25:657-665.
- 13 [11] Zhang W, Yang J, Chen J, Hou Y, Han X. Immunomodulatory and anti-tumor  
14 effects of an exopolysaccharide fraction from cultivated *Cordyceps sinensis*  
15 (Chinese caterpillar fungus) on tumour-bearing mice. Biotechnol. Appl. Biochem.  
16 2005;42:9-15.
- 17 [12] Yang Q, Yin Y, Yu G, Jin Y, Ye X, Shrestha A, Liu W, Yu W, Sun H. A novel protein  
18 with anti-metastasis activity on 4T1 carcinoma from medicinal fungus *Cordyceps*  
19 *militaris*. Int. J. Biol. Macromol. 2015;80:385-391.
- 20 [13] Sun J, Chen Y, Wu Y, Zhang X, Jiang L, Zhang Y. Bioassay-Guided Separation and  
21 identification of a new anti-lung cancer compound from *Cordyceps militaris* by  
22 means of off-line two-dimensional preparative chromatography, real-time cell  
23 analysis, and X-ray single-crystal diffraction. Chromatographia. 2015;78:495-506.
- 24 [14] Baik J, Mum S, Kim K, Park S, Yoon H, Kim D, Park M, Kim C, Lee Y. Apoptotic  
25 effects of cordycepin through the extrinsic pathway and p38 MAPK activation in  
26 human glioblastoma U87MG cells. J. Microbiol. Biotechnol. 2016;26:309-314.
- 27 [15] Song J, Wang Y, Teng M, Zhang S, Yin M, Lu J, Liu Y, Lee R, Wang D, Teng L.  
28 *Cordyceps militaris* induces tumor cell death via the caspase-3 dependent

- 1 mitochondrial pathway in HepG2 and MCF-7 cells. Mol. Med. Report.  
2 2016;13:5132-5140.
- 3 [16] Tuli H, Kumar G, Sandhu S, Sharma A, Kashyap D. Apoptotic effect of cordycepin  
4 on A549 human lung cancer cell line. Turk. J. Biol. 2015;39:306-311.
- 5 [17] Aramwit P, Bang N, Ratanavaraporn J, Nakpheng T, Srichana T. An anti-cancer  
6 cordycepin produced by *Cordyceps militaris* growing on the dead larva of *Bombyx*  
7 *mori* silkworm. J. Agric. Sci. 2014;6:41-53.
- 8 [18] Chiu C, Liu S, Tang C, Chan Y, El-Shazly M, Lee C, Du Y, Wu T, Chang F, Wu Y.  
9 Anti-inflammatory cerebroside from cultivated *Cordyceps militaris*. J. Agric. Food  
10 Chem. 2016;64:1540–1548.
- 11 [19] Choi J, Kim J, Lee M, Park D, Hong Y, Lee C. Metabolomics revealed novel  
12 isoflavones and optimal cultivation time of *Cordyceps militaris* fermentation. J.  
13 Agric. Food Chem. 2010;58:4258–4267.
- 14 [20] Rukachaisirikul V, Pramjit S, Pakawatchai C, Isaka M, Supothina S. 10-Membered  
15 macrolides from the insect pathogenic fungus *Cordyceps militaris* BCC 2816. J.  
16 Nat. Prod. 2004;67:1953-1955.
- 17 [21] Cunningham K, Mason M, Spring F, Hutchinson S. Cordycepin, a metabolic  
18 product isolated from cultures of *Cordyceps militaris*. Nature 1950;166:949.
- 19 [22] Ko B, Lu Y, Yao W, Liu T, Tzean S, Shen T, Liou J. Cordycepin regulates  
20 GSK-3 $\beta$ / $\beta$ -catenin signaling in human leukemia cells. PLOS ONE. 2013;8:issue 9.
- 21 [23] Lee H, Burger P, Vogel M, Friese K, Brüning A. The nucleoside antagonist  
22 cordycepin causes DNA double strand breaks in breast cancer cells. Invest. New  
23 Drugs. 2012;30:1917-1925.
- 24 [24] Jia L, Zhong L, Li H, Jing L. Chemical constituents in water fraction of  
25 *Abelmoschus esculentus*. Chin. Tradit. Herbal Drugs. 2011;42:2186-2188.
- 26 [25] Junko N, Yumiko H, Takashi A. Inhibition of *Trypanosoma cruzi* growth in  
27 mammalian cells by purine and pyrimidine analogs. Antimicrob. Agents Chemother.  
28 1996;40:2455-2458.

- 1 [26] Baer H, Serignese V, Ogbunude P, Dzimir M. Nucleoside transporters in  
2 *Leishmania major*: diversity in adenosine transporter expression or function in  
3 different strains. *Am. J. Trop. Med. Hyg.* 1992;47:87-91.
- 4 [27] Nishida H, Kushida M, Nakajima Y, Ogawa Y, Tatewaki N, Sato S, et al.  
5 Amyloid-beta-induced cytotoxicity of PC-12 cell was attenuated by *Shengmai-san*  
6 through redox regulation and outgrowth induction. *J. Pharmacol. Sci.*  
7 2007;104:73-81.

8  
9  
10  
11 Legend to figure

12  
13 Fig. 1. Structures of compounds **1** and **2**.

14  
15 Fig. 2. Cytotoxicity of compounds **1** and **2** against A549, PANC-1 and MCF-7 cancer  
16 cells tested by MTT assay.

17 Notes: The concentrations of compounds **1** and **2** were adjusted to be 1  $\mu\text{M}$ , 10  $\mu\text{M}$  and  
18 30  $\mu\text{M}$ , respectively. Sulforaphane (SFN; 30  $\mu\text{M}$ ) was used as positive control. All data  
19 are shown as means  $\pm$  SD (n=5-6). Differences among the all groups were evaluated  
20 using a 1-way analysis of variance (ANOVA) followed by the Dunnet test.

21 \* indicates significant difference compared with CT ( $P < 0.05$ ); \*\* indicates significant  
22 difference compared with CT groups ( $P < 0.01$ ).




Fig. 1 Qiu et al

Fig.1

254x190mm (72 x 72 DPI)

review




Fig. 2 Qiu et al

Fig.2

254x190mm (72 x 72 DPI)

review