

Quantitative validation of nicotine production in tea (*Camellia sinensis* L.)

メタデータ	言語: eng 出版者: 公開日: 2019-02-20 キーワード (Ja): キーワード (En): 作成者: Ikka, Takashi, Yamashita, Hiroto, Kurita, Ikuya, Tanaka, Yasuno, Taniguchi, Fumiya, Ogino, Akiko, Takeda, Kazuya, Horie, Nobuhiro, Hojo, Hiroshi, Nanjo, Fumio, Morita, Akio メールアドレス: 所属:
URL	http://hdl.handle.net/10297/00026279

RESEARCH ARTICLE

Quantitative validation of nicotine production in tea (*Camellia sinensis* L.)

Takashi Ikka¹, Hiroto Yamashita¹, Ikuya Kurita¹, Yasuno Tanaka¹, Fumiya Taniguchi², Akiko Ogino², Kazuya Takeda³, Nobuhiro Horie³, Hiroshi Hojo³, Fumio Nanjo³, Akio Morita^{1*}

1 Faculty of Agriculture, Shizuoka University, Shizuoka, Shizuoka, Japan, **2** Tea Research Division of NARO Institute of Fruit and Tea Science (NIFTS), Makurazaki, Kagoshima, Japan, **3** Mitsui Norin Co., Ltd., Fujieda, Shizuoka, Japan

* morita.akio@shizuoka.ac.jp



Abstract

Endogenous nicotine was confirmed to be present in tea plants (*Camellia sinensis* L.) by liquid chromatography-tandem mass spectrometry of tea samples from tea-producing regions in six Asian countries. All samples contained nicotine (0.011–0.694 $\mu\text{g g}^{-1}$ dry weight). Nicotine contents remained constant during manufacturing of green, oolong and black teas, implying that nicotine is stable against heating, drying, enzymatic oxidation and mechanical damage during processing. Flower buds and seeds of cultivar Yabukita also contained nicotine (0.030–0.041 $\mu\text{g g}^{-1}$ dry weight). A comparison of two cultivars revealed that higher nicotine contents were found in the black tea cultivar Benifuki. All plant parts of hydroponic Yabukita contained nicotine (0.003–0.013 $\mu\text{g g}^{-1}$ dry weight). Tea cells cultured in B5 medium as well as roots and stems of tea seedlings contained nicotine levels similar to those of new leaves from field-grown plants. Although the levels of endogenous nicotine in tea plants are extremely low and sample contamination cannot be discounted, these levels exceed the maximum acceptable limit in Japan (0.01 $\mu\text{g g}^{-1}$ dry weight).

OPEN ACCESS

Citation: Ikka T, Yamashita H, Kurita I, Tanaka Y, Taniguchi F, Ogino A, et al. (2018) Quantitative validation of nicotine production in tea (*Camellia sinensis* L.). PLoS ONE 13(4): e0195422. <https://doi.org/10.1371/journal.pone.0195422>

Editor: Wei Wang, Henan Agricultural University, CHINA

Received: December 12, 2017

Accepted: March 22, 2018

Published: April 9, 2018

Copyright: © 2018 Ikka et al. This is an open access article distributed under the terms of the [Creative Commons Attribution License](https://creativecommons.org/licenses/by/4.0/), which permits unrestricted use, distribution, and reproduction in any medium, provided the original author and source are credited.

Data Availability Statement: All relevant data are within the paper and its Supporting Information files.

Funding: This work was supported by the Japanese Society for the Promotion of Science Grant-in-Aid for Scientific Research Number 26292010 (AM) and 17H03756 (AM). We carried out this research entirely through public research funds in Japan (JSPS KAKENHI). We did not receive financial assistance for this study from the relevant private company (Mitsui Norin Co. Ltd); Mitsui Norin Co. Ltd provided salaries for authors

Introduction

Tea, derived from the tea plant (*Camellia sinensis* L.), is one of the world's most popular beverages. Various types, such as green, black and/or oolong, are consumed in different localities. Unlike oolong and black teas, green tea is made without the use of withering and enzymatic oxidation (e.g. polyphenol oxidase and peroxidase) processes. Consequently, catechins and ascorbic acid contents of green tea are generally higher than those of other teas [1, 2]. Green tea, produced mainly in China, Vietnam and Japan, is classified into Chinese and Japanese styles [3]. Japanese-style green tea typically differs from Chinese in the way in which the leaves are heated: Japanese-style tea leaves are heated with a steaming machine—which minimizes the deactivation of oxidation enzymes—whereas Chinese tea production involves a parching machine (S1 Fig). Sencha, the most common Japanese green tea, plays an important role in Japanese culture. The export volume of Japanese green tea has recently increased greatly. Because these compounds are beneficial to human health, their higher levels in green tea have

KT, NH, HH and FN. The specific roles of these authors are articulated in the 'author contributions' section. The funders also did not have any additional role in the study design, data collection, decision to publish, or preparation of the manuscript.

Competing interests: Authors KT, NH, HH, and FN are affiliated with and received salary from a commercial company, Mitsui Norin Co., Ltd. There are no patents, products in development or marketed products to declare. This does not alter our adherence to all the PLOS ONE policies on sharing data and materials.

resulted an increase in green tea production and consumption—up to one-third that of black tea.

Nicotine, which is an alkaloid compound mainly found in the genus *Nicotiana* (e.g. tobacco), has been detected in many food crops [4]. Nicotine acts as an agonist at nicotinic acetylcholine receptors, therefore, it has been used as a pesticide because of its toxicity to organisms. Previous studies have revealed the presence of nicotine not only in solanaceous crops, namely, potato (*Solanum tuberosum*), tomato (*S. lycopersicum*), eggplant (*S. melongena*) and pepper (*Capsicum annum*), but also in wild mushrooms and cauliflower (*Brassica oleracea*) [5–11]. These findings suggest that nicotine is widely distributed in plants and fungi. In a survey reported by the European Food Safety Authority (EFSA), nicotine concentrations were measured in 332 tea samples (87 green, 239 black and 6 Chinese white) originating from numerous tea-producing countries [4]. Nicotine was detected in 274 samples, with contents of 53 samples found to be over the default maximum residue level (MRL) of 0.01 mg kg⁻¹ dry weight (DW). Prior to this EFSA report, other several studies uncovered nicotine in tea at levels ranging from 0 to 28.01 mg kg⁻¹ DW [5–8]. In Japan, where no nicotine-containing pesticides have been listed in the Agricultural Chemicals Regulation Law since 2001, the MRL of nicotine is set at 0.01 mg kg⁻¹ DW. Japanese green tea should thus have no detectable exogenous nicotine. The nicotine content of Japanese green tea was not measured in any of the above-mentioned studies.

Tobacco (*Nicotiana tabacum*) is the only plant in which nicotine biosynthesis has actually been confirmed to occur [6]. Various authors [5, 6] have proposed that the nicotine detected in tea is due to insecticide contamination. Sheen [5] has suggested that the content of nicotine in tomato is lessened during processing due to enzymatic oxidation and other chemical reactions. In contrast, however, a screening for nicotine in various processed tomato products by Siegmund et al. [8] has indicated that nicotine is thermally stable and does not degrade or evaporate during processing. In tobacco leaves, various post-harvest reactions during curing degrade nicotine into its nitrogen oxide as well as into cotinine and other alkaloids [12]. Also in tomato, Siegmund et al. [8] has observed that nicotine contents in tomato decrease with increasing ripeness and differ between varieties. Taken together, these findings suggest that several factors affect nicotine content in plants. Nevertheless, the above-mentioned assumptions as well as the possibility of nicotine biosynthesis have not been confirmed in plants other than tobacco.

To determine whether endogenous nicotine is present in Japanese green tea, in this study we measured nicotine contents of several types of tea from Japan and five other Asian countries. Nicotine was detected in all analyzed Japanese green tea samples, but the contents were lower than those of black teas. To determine the effect of different manufacturing processes used to generate teas such as green, oolong and black, we next measured nicotine contents using two Japanese cultivars: Yabukita, an important Japanese green tea cultivar, and Benifuki, a newly introduced black tea cultivar and Assam hybrid. We also investigated the nicotine contents of various tea plant organs (leaves, stems, roots, flower buds and seeds) and explored seasonal changes in nicotine contents of field-grown tea plants. The nicotine content of aseptically cultured tea cells was also determined. Finally, the origin of nicotine in tea was considered in the light of these findings.

Materials and methods

Plant materials

Tea from various tea-producing districts in Asia. Thirty-two green tea (*C. sinensis* var. *assamica*), 79 black tea (*C. sinensis* var. *sinensis* and/or *C. sinensis* var. *assamica*) and 1 oolong tea (*C. sinensis* var. *sinensis*) sample(s) were subjected to nicotine analysis (S1 Table). The Indian

tea samples were purchased from 16 and 3 tea estates in Darjeeling and Assam, respectively, by Mitsui Norin Co. Ltd., one of the major tea companies in Japan. With the exception of one estate in Assam from which no autumn sample was obtained, tea collections were made during all four crop seasons (first, second, rain/monsoon and autumn; [S4 Fig](#)). All samples were from organically grown tea leaves received Japanese Agricultural Standard (JAS) certification, i.e. grown without the use of chemicals including insecticides. In the case of the Japanese samples, which were collected by the Tea Research Division of the NARO Institute of Fruit and Tea Science (NIFTS) (Makurazaki, Kagoshima, Japan), the leaves came from a cultivated tea garden where no insecticides (including nicotine) have been used for several decades. Other tea samples were purchased from organic cultivation in local markets of each country. All green teas collected in Japan were typical Japanese types (Sencha), produced using steam, while those from Taiwan were Chinese types (pan-fired). Tea samples were dried in a freeze dryer (FDU-2000, EYELA, Tokyo, Japan) and then ground into a fine powder with a vibrating mill (SA300, Yamato, Tokyo, Japan). The samples were stored in desiccators until nicotine analysis.

Tea subjected to different manufacturing processes. New shoots of 30-year-old plants of two tea cultivars (Yabukita and Benifuki) were harvested during the first crop season on 25 April 2015 at the NIFTS tea garden (Makurazaki, Kagoshima, Japan). The shoots were immediately used to manufacture Japanese (Sencha, steaming type) and Chinese (pan-fired tea, pan-firing type) green teas, oolong tea and black teas following the processes shown in [S1 Fig](#). Following processing, samples were collected for nicotine analysis and stored in a freezer at -20°C until tea manufacturing. The manufactured tea samples were then freeze-dried, ground and stored in the same manner as described above.

Tea from each growing season. New leaves, new stems, mature leaves and roots of Yabukita and Benifuki were harvested from the NIFTS tea garden (Makurazaki, Kagoshima, Japan) on 18 April (first crop season), 18 June (second crop season) and 13 August (third crop season) in 2014. Mature leaves and roots were also harvested on 27 February during the off season in 2015. The roots were collected from the soil layer between hedges at a depth of 10 cm. After washing three times with deionized water, the samples were freeze-dried, ground and stored in the same manner as described above.

Tea plant parts and cultured tea cells. One-year-old rooted cuttings of cultivars Yabukita and Benifuki were transplanted on 8 April 2014 into Wagner pots (1/2,000a) containing 10 L of aerated nutrient solution and grown in a greenhouse at Shizuoka University (Shizuoka, Shizuoka, Japan). The nutrient solution, prepared following the method of Konishi et al. [13], was renewed weekly. The pH of the hydroponic solution was adjusted to 4.2 with 5 N H_2SO_4 . After 90 days (on 7 July 2014), the tea plants were harvested and divided into six parts: new leaves, new stems, mature leaves, mature stems, lignifying roots and white roots ([S2 Fig](#)).

Flower buds of 35-year-old Yabukita tea plants were collected from the tea garden of the Shizuoka Prefecture Tea Research Center (Kikugawa, Shizuoka, Japan) on 5 November 2015. The flower buds were immediately divided into petals, filaments and anthers (Panel A in [S3 Fig](#)). On 26 October 2015, fully mature fruits containing one to four seeds were collected from 20-year-old Yabukita tea plants in a private tea garden in Shizuoka, Japan. After peeling away the pericarp, the outer seed coat was broken with a hammer, and the outer and inner seed coats were carefully removed. The collected flower buds and tea seeds were freeze-dried, ground and stored as described earlier. Because collection of adequate samples for nicotine analysis was difficult, only a single replicate was performed.

Suspension-cultured tea cells (Panel B in [S3 Fig](#)) were prepared from calli derived from anthers of Yabukita [14]. The cells were subcultured weekly in B5 medium at pH 5.6 in darkness at 25°C on a rotary shaker (100 rpm). Exponentially growing cells were harvested after 9 days by suction filtration, suspended and rinsed three times with a simple solution containing

0.2 mM CaCl₂ at pH 5.6. The cells were frozen in liquid nitrogen and freeze-dried. B5 medium was also collected before and after 9 days cultivation and stored at -20°C until analysis. The samples were freeze-dried, ground and stored in the same manner as described above.

Aseptic tea seedlings were prepared following the method of Morita et al. [15] from the same tea seeds as mentioned above. Four months after germination, the aseptic tea seedlings were divided into leaves, stems and roots (Panel C in S3 Fig) and freeze-dried, ground and stored as described above.

Measurement of nicotine

Nicotine extraction from tea samples was carried out according to the modified QuEChERS method of Cavalieri et al. [16]. One gram of powdered tea sample was added to a 50-mL polytetrafluoroethylene (PTFE) centrifuge tube along with 50 µL of nicotine d₃ internal standard (Sigma, MO, USA) at a concentration of 5 µg mL⁻¹. To this mixture was added 20 mL of 0.1 M ammonia solution followed by incubation for 10 min to allow for swelling of the tea powder. After adding 10 mL acetonitrile, 6 g MgSO₄ and 1.5 g CH₃COONa, the solution was mixed on a shaker at 3,000 rpm for 5 min (SA300, Yamoto, Tokyo, Japan). Following centrifugation at 5,000 ×g for 4 min, 1.5 mL of the resulting supernatant was transferred to a 2-mL PTFE centrifuge tube containing 50 mg of Primary Secondary Amine (Agilent, CA, USA) and 150 mg of MgSO₄. The sample was mixed for 1 min and then centrifuged at 18,000 ×g for 5 min. The supernatant was analyzed by liquid chromatography-tandem mass spectrometry (LC-MS/MS). Nicotine in B5 medium from cultured tea cells was also extracted using the same procedure, except that the final concentration of added ammonia solution was 0.1 M.

Nicotine was quantified on a liquid chromatograph (1100 series, Agilent, CA, USA) equipped with a triple quadrupole mass spectrometer (3200 QTRAP®, Sciex, MA, USA) with a TurboIon-Spray® (Sciex, MA, USA) electrospray ionization source operated in positive ionization mode. The chromatographic analysis was performed on a 150 × 2.1 mm InertSustain phenylhexyl column (GL Science, Tokyo, Japan). Acetonitrile (B) and 20 mM ammonium hydrogen carbonate (A) were used as the mobile phase at a flow rate of 0.25 mL min⁻¹. Elution was performed with the following gradient: initial concentration of 10% B, followed by a 0.1-min linear gradient from 10% to 35% B, a 0.9-min hold at 35% B, a 0.1-min linear gradient from 35% to 50% B, a 1.9-min hold at 50% B, a 2.0-min linear gradient from 50% to 75% B, a 2.0-min hold at 75% B, a 0.1-min gradient from 75% to 10% B, and a final concentration of 10% B for 4.9 min.

Mass spectrometry parameters were optimized as follows: ion spray voltage (5500 V); ionization source temperature (600°C); curtain and CAD gases (138 kPa); gas 1 (552 kPa); gas 2 (483 kPa). Data analyses were performed using Analyst software (version 1.6, Sciex, MA, USA). Chemicals used for nicotine determination were LC/MS grade or analytical grade. The limit of quantification (LOQ) of our method was set at 0.005 µg g⁻¹ DW, the minimum concentration giving a signal-to-noise ratio of at least 10.

Statistical analysis

Data were analyzed by Tukey's and Dunnett's tests. A *P*-value < 0.05 was considered to represent a significant difference.

Results

Analysis of tea collected from different tea-producing districts in Asia

Nicotine contents of tea samples collected from different tea-producing districts in Asia are shown in Table 1. All samples had nicotine contents above 0.01 µg g⁻¹ DW. The maximum

Table 1. Nicotine contents of tea samples obtained from tea-producing areas in Asia.

Tea Samples				Nicotine content ($\mu\text{g g}^{-1}$ dry weight)						
Type	Subtype	Country or district		Number	Average	\pm	S.D.	Minimum	Median	Maximum
Green tea	Sencha	Japan		31	0.019	\pm	0.006	0.013	0.017	0.040
	Pan-fired	Taiwan		1	0.011					
	Sub total			32	0.019	\pm	0.006	0.011	0.017	0.040
Black tea		India	Assam	11	0.331	\pm	0.069	0.246	0.332	0.449
			Darjeeling	64	0.273	\pm	0.151	0.100	0.219	0.694
			China	1	0.289					
			Taiwan	1	0.121					
			Indonesia	1	0.101					
			Vietnam	1	0.024					
	Sub total			79	0.274	\pm	0.145*	0.024	0.245	0.694
Oolong tea		Taiwan		1	0.020					
Total				112	0.199	\pm	0.169	0.011	0.189	0.694

* Significant difference between green tea and black tea (t-test, $P < 0.01$).

<https://doi.org/10.1371/journal.pone.0195422.t001>

detected nicotine content, $0.694 \mu\text{g g}^{-1}$ DW, was in black tea from a Darjeeling organic tea estate; the minimum, $0.011 \mu\text{g g}^{-1}$ DW, was found in green tea from Taiwan (Table 1). The average nicotine content of Japanese green tea, $0.019 \mu\text{g g}^{-1}$ DW ($0.013\text{--}0.040 \mu\text{g g}^{-1}$ DW), was lower than that of Assam ($0.331 \mu\text{g g}^{-1}$ DW) and Darjeeling ($0.273 \mu\text{g g}^{-1}$ DW). With respect to tea types, the average nicotine content of green tea ($0.019 \mu\text{g g}^{-1}$ DW) was significantly lower than that of black tea ($0.274 \mu\text{g g}^{-1}$ DW).

Seasonal fluctuation in nicotine contents of Indian black tea collected from 19 tea estates is shown in S4 Fig. Because we observed a different pattern of fluctuation on each tea estate, no overall trend was discernable across estates.

Analysis of nicotine content as a function of manufacturing process

Nicotine contents of tea derived from green tea (Sencha and pan-fired), oolong tea and black tea manufacturing processing are shown in Table 2. After plucking, fresh tea leaves containing nicotine contents of $0.021 \mu\text{g g}^{-1}$ DW (in Yabukita) and $0.026 \mu\text{g g}^{-1}$ DW (in Benifuki) were used for the manufacturing treatments. Although heating temperature, treatment period and the application of fermentation and/or withering (oxidation by oxidative enzymes in leaves) varied among processes, nicotine levels in processed tea leaves were unchanged relative to fresh leaves regardless of the manufacturing process used.

Analysis of nicotine content according to growing season

Table 3 displays seasonal changes in nicotine contents of new leaves, new stems, mature leaves and roots of Yabukita and Benifuki. In new leaves and stems of Yabukita, nicotine contents during the first crop season (0.010 and $0.011 \mu\text{g g}^{-1}$ DW, respectively) were significantly lower than those during the second (0.023 and $0.033 \mu\text{g g}^{-1}$ DW) and third (0.022 and $0.036 \mu\text{g g}^{-1}$ DW). Nicotine contents in these two organs were highest in Benifuki during the third season (0.036 and $0.032 \mu\text{g g}^{-1}$ DW). Although nicotine contents in mature leaves of both cultivars were highest in the off season ($0.024 \mu\text{g g}^{-1}$ DW in Yabukita and $0.021 \mu\text{g g}^{-1}$ DW in Benifuki), the ranges of their seasonal changes were smaller than those of new leaves and new stems. In roots of both cultivars, nicotine contents remained constant (ca. $0.028 \mu\text{g g}^{-1}$ DW in Yabukita

Table 2. Nicotine contents of tea leaves following green (Sencha and pan-fired), oolong and black tea manufacturing processes.

Process	Nicotine content ($\mu\text{g g}^{-1}$ dry weight)					
	Yabukita			Benifuki		
Fresh leaves	0.021	±	0.002	0.026	±	0.005
Green tea (Sencha)						
Steaming	0.021	±	0.001	0.023	±	0.001
Primary drying	0.020	±	0.002	0.025	±	0.004
Rolling	0.020	±	0.001	0.022	±	0.001
2nd drying	0.017	±	0.001	0.022	±	0.001
Final rolling	0.018	±	0.001	0.023	±	0.002
Drying	0.020	±	0.003	0.025	±	0.001
Green tea (Pan-fired)						
1st panning	0.021	±	0.005	0.021	±	0.001
1st rolling	0.018	±	0.001	0.023	±	0.001
2nd panning	0.019	±	0.001	0.024	±	0.000
2nd rolling	0.021	±	0.000	0.023	±	0.002
2nd drying	0.018	±	0.002	0.022	±	0.001
Drying	0.021	±	0.002	0.023	±	0.002
Oolong tea						
Sunlight withering	0.020	±	0.001	0.024	±	0.001
Indoor withering	0.018	±	0.001	0.024	±	0.002
Panning	0.021	±	0.002	0.024	±	0.001
Rolling	0.018	±	0.002	0.026	±	0.002
2nd rolling	0.020	±	0.000	0.025	±	0.002
Drying	0.019	±	0.001	0.026	±	0.002
Black tea						
Indoor withering	0.020	±	0.002	0.028	±	0.000
Rolling	0.019	±	0.002	0.028	±	0.001
Fermentation	0.019	±	0.002	0.027	±	0.003
Drying	0.021	±	0.001	0.025	±	0.001

Values are means±S.D. ($n = 3$).

<https://doi.org/10.1371/journal.pone.0195422.t002>

and $0.030 \mu\text{g g}^{-1}$ DW in Benifuki). When averaged over first to third crop seasons, the highest nicotine contents were found in roots.

Analysis of nicotine in various plant organs and cultured tea cells

Nicotine contents in Yabukita and Benifuki cultured hydroponically for 90 days are given in Table 4. In Yabukita, the lowest nicotine contents were in new shoots ($0.003 \mu\text{g g}^{-1}$ DW, below the LOQ), with contents similar among other plant parts (0.009 – $0.013 \mu\text{g g}^{-1}$ DW). In Benifuki, in contrast, white roots and lignifying roots contained nicotine at a significantly higher level than the contents found in leaves and stems.

Nicotine contents in flower buds and seeds of Yabukita are shown in Table 5. In flower buds, nicotine contents of filaments ($0.041 \mu\text{g g}^{-1}$ DW) were higher than those of petals ($0.031 \mu\text{g g}^{-1}$ DW) and anthers ($0.033 \mu\text{g g}^{-1}$ DW). Seeds had nicotine contents ($0.030 \mu\text{g g}^{-1}$ DW) similar to those of petals.

Nicotine contents of suspension-cultured tea cells and B5 medium before and after 9 days incubation are displayed in Table 5. Nicotine was detected in tea cells at a content of $0.031 \mu\text{g g}^{-1}$ DW, but was not found in B5 medium either before or after incubation.

Table 3. Nicotine contents of new leaves, new stems, mature leaves and roots of two tea cultivars (Yabukita and Benifuki) harvested during first, second and third crop seasons and the off season in 2014 and 2015.

Parts	Cultivars	Crop season (nicotine content, $\mu\text{g g}^{-1}$ dry weight)				Statistical analysis*		
		1st season	2nd season	3rd season	Off- season	Cultivar	Season	Cultivar \times Season
		(18th Apr.)	(18th Jun.)	(13th Aug.)	(27th Feb.)			
New leaves	Yabukita	0.010 \pm 0.001 a	0.023 \pm 0.003 b	0.022 \pm 0.004 b	n.t.	$P < 0.05$	$P < 0.01$	$P < 0.05$
	Benifuki	0.015 \pm 0.004 a	0.016 \pm 0.001 a	0.036 \pm 0.007 b	n.t.			
New stems	Yabukita	0.011 \pm 0.000 a	0.033 \pm 0.005 b	0.036 \pm 0.008 b	n.t.	NS	$P < 0.01$	NS
	Benifuki	0.013 \pm 0.000 a	0.023 \pm 0.001 a	0.032 \pm 0.003 b	n.t.			
Mature leaves	Yabukita	0.018 \pm 0.002 ab	0.014 \pm 0.001 a	0.013 \pm 0.000 a	0.023 \pm 0.003 b	NS	$P < 0.01$	NS
	Benifuki	0.019 \pm 0.004 ab	0.014 \pm 0.001 a	0.015 \pm 0.001 ab	0.021 \pm 0.001 b			
Roots	Yabukita	0.028 \pm 0.001	0.027 \pm 0.001	0.027 \pm 0.001	0.028 \pm 0.001	$P < 0.01$	NS	NS
	Benifuki	0.028 \pm 0.002	0.030 \pm 0.003	0.031 \pm 0.001	0.031 \pm 0.001			

Data represent means \pm S.D. ($n = 3$). n.t.: not tested. Different letters within the same row indicate significant differences (Tukey's HSD test, $P < 0.05$).

* Significant differences among organs of the two cultivars were estimated by two-way ANOVA. NS: not significant ($P > 0.05$).

<https://doi.org/10.1371/journal.pone.0195422.t003>

Table 5 shows nicotine contents of leaves, stems and roots of tea seedlings grown under aseptic condition for 90 days. Nicotine was detected at a content of at least $0.033 \mu\text{g g}^{-1}$ DW in all plant organs, with the maximum level found in roots.

Discussion

Among the tea samples collected from tea-producing localities in six Asian countries, we found detectable levels of nicotine in Japanese green tea (Table 1). This result reconfirms the presence of nicotine in Japanese green tea. With respect to maximum values among tea-producing regions, nicotine contents of organically grown black tea from Assam and Darjeeling were higher than those of Japanese green tea. In a study of 41 black, 14 oolong, 26 green and 6 white tea samples, conducted with LC-ESI-MS/MS using a modified QuEChERS method, Thräne et al. [17] observed that the highest nicotine contents ($> 0.6 \mu\text{g g}^{-1}$ DW) were found in

Table 4. Nicotine contents of various plant organs of two hydroponically cultivated tea cultivars (Yabukita and Benifuki).

Parts	Nicotine content ($\mu\text{g g}^{-1}$ dry weight)					
	Yabukita			Benifuki		
New leaves	0.003	\pm	0.000 a	0.008	\pm	0.006 A
New stems	0.01	\pm	0.000 b	0.01	\pm	0.000 A
Mature leaves	0.009	\pm	0.001 b	0.009	\pm	0.000 A
Mature stems	0.012	\pm	0.001 b	0.01	\pm	0.001 A
White roots	0.011	\pm	0.000 b	0.029	\pm	0.003 B
Lignifying roots	0.013	\pm	0.004 b	0.024	\pm	0.003 B

Data are means \pm S.D. ($n = 3$). Different letters indicate significant differences ($P < 0.05$, Tukey's). Lowercase and uppercase characters indicate significant differences among organs of Yabukita and Benifuki, respectively.

<https://doi.org/10.1371/journal.pone.0195422.t004>

Table 5. Nicotine contents of flower buds and seeds of tea plants (Yabukita), cultured tea cells and B5 medium before and after 9 days cultivation, and leaves, stems and roots of tea seedlings cultured under aseptic conditions for 90 days.

Samples		Nicotine content ($\mu\text{g g}^{-1}$ dry weight)
Flower buds ¹⁾	Petals	0.031
	Filaments	0.041
	Anthers	0.033
Seeds ¹⁾		0.030
B5 medium ²⁾	Before use	n.q.
	After cultivation	n.q.
Tea cells ²⁾		0.033 \pm 0.019
Tea seedlings ¹⁾	Leaves	0.033
	Stems	0.044
	Roots	0.059

¹⁾ Values are from a single replicate ($n = 1$).

²⁾ Values are means \pm S.D. ($n = 10$). n.q.: not quantified [below the LOQ ($0.005 \mu\text{g g}^{-1}$ dry weight)].

<https://doi.org/10.1371/journal.pone.0195422.t005>

samples from India and China. They also reported that tea samples from Japan had contents below $0.19 \mu\text{g g}^{-1}$ DW. These results suggest that nicotine contents of tea leaves differ among producing regions and cultivars. In spite of the relatively small number of samples in the present study, we also observed that nicotine contents of black tea ($0.024\text{--}0.694 \mu\text{g g}^{-1}$ DW) were significantly higher than those of green tea ($0.011\text{--}0.40 \mu\text{g g}^{-1}$ DW) (Table 1). This trend is in line with average nicotine contents reported by EFSA 2011 [4] of $0.130 \mu\text{g g}^{-1}$ DW ($0\text{--}0.873 \mu\text{g g}^{-1}$ DW) for black tea vs. $0.075 \mu\text{g g}^{-1}$ DW for green tea ($0\text{--}0.440 \mu\text{g g}^{-1}$ DW)—although it should be noted that these samples did not include Japanese green tea. In tobacco plants, it is reported that nicotine synthesized in roots accumulates in vacuoles [18] and so also the flavonoids such as catechins are found in the vacuoles in tea [19]. Understanding the significance of these observed differences among producing regions and cultivars will require future investigation of cultivation environments, such as soil conditions and climate, and the introduction of additional cultivars. Thräne et al. [17] found that average nicotine contents seemed to increase with increasing degree of fermentation, although black teas with contents below a LOQ of $0.01 \mu\text{g g}^{-1}$ DW were also observed. In contrast, the results of Sheen [5] suggest that nicotine is reduced in tomato via enzymatic oxidation and other chemical reactions during processing. As is well known, tea is classified into three types based on the degree of fermentation: non-fermented (green tea), semi-fermented (oolong tea) and full-fermented (black tea). During the tea manufacturing process, leaves are typically heated by steaming (ca. 100°C), hot air exposure ($65\text{--}115^\circ\text{C}$) or by placement on high-temperature pans (ca. 300°C) to diminish oxidative enzyme activities and for drying (S1 Fig). Nicotine contents in tea leaves would thus be expected to change during manufacturing. In this study, however, nicotine contents remained constant regardless of tea type (Sencha, pan-fired, oolong or black). This result suggests that heating and fermentation have no effect on nicotine contents in tea leaves. Similarly, the application of withering, a leaf-damaging process, during the manufacturing of oolong and black teas had no effect on nicotine contents.

A comparison of two cultivars revealed that nicotine contents in *C. sinensis* var. *sinensis* ‘Yabukita’ were lower than those of *C. sinensis* var. *sinensis* \times var. *assamica* ‘Benifuki’ in new leaves subjected to tea manufacturing processes (Table 2), in new leaves of field-grown tea plants (Table 3) and in new leaves and roots of hydroponically grown tea plants (Table 4).

These results suggest that nicotine contents of *C. sinensis* var. *assamica* are higher than those of var. *sinensis*. Because leaves of varieties *sinensis* and *assamica* are generally used to make green and black teas, respectively, this implies that a difference exists in nicotine accumulation among tea varieties. In samples of var. *assamica* from India, however, we detected nicotine contents as low as $0.100 \mu\text{g g}^{-1}$ DW, even though the highest value was also detected in an Indian tea sample ($0.694 \mu\text{g g}^{-1}$ DW; Table 1). A wide range of nicotine contents in Indian black tea has also been observed in the above studies. Two possible explanations for these results can be proposed. One possibility is genetic variation in nicotine synthesis in tea plants; the other is environmental contamination due to various reasons—for example, from smokers (cigarette smoke or nicotine residues on fingers of harvesters) [20] or seepage into ground water from nearby cultivated tobacco plants [21]. Because nicotine was detected in all types of tea samples, the latter possibility seems unlikely but must still be considered.

New tea shoots (new leaves and stems) were harvested several times every year from spring to autumn. In general, plucking was carried out four times annually in India and two or three times in Japan. No clear seasonal trend, such as highest value, could be discerned in the nicotine content of Indian black teas collected from 14 tea estates (S4 Fig). In Japan, nicotine contents of new leaves and stems of field-grown tea plants were highest during second and third crop seasons (Table 3). In tobacco plants, damage to leaves increases nicotine contents, indicating that nicotine biosynthesis is affected in response to mechanical wounding [22–24]. Wang et al. [25] have also reported that excision of tobacco shoot apices causes dramatic increases in nicotine contents in leaves. Before beginning this study, we therefore assumed that wounding caused by plucking of tea new shoots might increase nicotine contents, thereby leading to higher levels in tea leaves during second and third crop seasons compared with the first season. Although this trend was observed in Japanese field-grown plants, no such pattern was uncovered in Indian black tea. In tobacco plants, nicotine biosynthesis is regulated by multiple biotic and abiotic stresses during the growing season, such as drought, heat and herbivore or insect damage [26, 27]. Because the type and degree of stress experienced under field conditions among tea-growing regions was variable, the individual seasonal fluctuation of nicotine content in new tea shoots differed between tea estates. The seasonal change in nicotine contents of new shoots observed in this study consequently suggests that nicotine biosynthesis and degradation is carried out in tea plants.

Leaves, stems and roots as well as seeds and flower buds also contained nicotine (Table 5). In field-grown tea plants, the annual average nicotine content was highest in roots (Table 3). Roots were also the organs with the highest contents of nicotine in tea plants grown hydroponically in a greenhouse and aseptic seedlings (Tables 4 and 5). In tobacco plants, nicotine is synthesized in roots [28], where it remains most abundant until topping [25]. On the basis of nicotine presence in other plant organs besides fruits, Sheen [5] concluded that tomato also biosynthesizes nicotine. Although no other reliable evidence is currently available, our data strongly suggest that nicotine is synthesized in the roots of tea plants. Similar to aseptic seedlings, suspension-cultured tea cells contained nicotine ($0.033 \mu\text{g g}^{-1}$ DW) (Table 5). These cells were derived from anthers of Yabukita and had been sub-cultured weekly since 1994 in B5 medium under aseptic conditions. Nicotine was not detected in the B5 medium alone. Even when grown under these conditions, which virtually guaranteed the complete exclusion of nicotine contamination from exogenous sources, tea cells contained nicotine. These results strongly support that endogenous nicotine is present in tea plants.

In tobacco plants, nicotine is synthesized from putrescine, a polyamine produced either directly from ornithine in a reaction catalyzed by ornithine decarboxylase (EC 4.1.1.17) or indirectly from arginine in a reaction catalyzed by arginine decarboxylase (EC 4.1.1.17) [29, 30]. The first step in nicotine biosynthesis is the conversion of putrescine to N-

methylputrescine, which is catalyzed by putrescine N-methyltransferase [31–35]. Although the final step of nicotine biosynthesis is still not known clearly, recently it was reported that an A622, a member of PIP family of NADPH-dependent reductases, and berberine bridge enzyme like proteins (BBLs) have these function [36]. Besides, the production of nicotine and the expression levels of genes involved in nicotine biosynthesis showed an increase after wounding and jasmonate treatment [37] mediated by the jasmonate signaling cascade, which is regulated defense responses against environmental stresses [38]. Wang et al. [25] reported that nicotine content increases in tobacco plants harvest at the upper leaves during cultivation. In general, even in the cultivation of tea, the risk of infestation of insects and pathogenic attacks also increases in the summer season compared with the winter season. Indeed, our results represent that nicotine contents in tea differ in each region and season (Tables 1 and 3), especially their values tends to be high in summer season. In other words, environmental factors related to jasmonate signaling may affect nicotine biosynthesis in tea. The activity and gene expression of these enzymes have not been examined in tea plants. To provide further evidence of endogenous nicotine, molecular-level investigation of nicotine synthesis in tea plants is needed.

Compared with other plant species, e.g. solanaceous; $< 100 \mu\text{g kg}^{-1}$ FW (fresh weight) on average [8, 10], it was confirmed that the content of nicotine contained in tea was of same level. Based on the EFSA report 2011 [4], intake of nicotine via a cup of tea (1.5 g leaf/150 ml) calculated from our results is 0.008 to 0.500 μg when calculated at a transition rate of 48% (median), and it is 0.017 to 1.041 μg when it is 100%, these were also reported to be of same level ($\leq 1 \mu\text{g day}^{-1}$) as in solanaceous, and much lower than that via passive smoking (80 $\mu\text{g day}^{-1}$) or smoking (number of cigarettes $\times 1 \text{ mg day}^{-1}$) [10]. According to an EFSA report 2011 [4], tea is the major source of ingested nicotine among herbal infusions and spices, but its maximum contribution amounts to only 0.7% of acceptable daily intake. On this basis, EFSA proposed a MRL of $0.6 \mu\text{g g}^{-1}$ for tea. With the exception of a few samples, nicotine contents obtained in this study were lower than this value, which suggests that tea has no toxic effect when consumed as a beverage. In light of our data, we recommend future revision of this MRL, with careful consideration given to samples exceeding this value to avoid the risk of contamination.

Conclusions

In this study, all purchased tea samples and samples of roots, stems, flower buds and seeds contained nicotine ranging from 0.003 to 0.694 $\mu\text{g g}^{-1}$ DW. In addition, tea cells cultured in B5 medium without nicotine contained nicotine at a content of 0.033 $\mu\text{g g}^{-1}$ DW. These results suggest that tea plants contain nicotine that is not derived from exogenous sources. In addition, nicotine contents in tea leaves changed seasonally and differed among cultivars. Furthermore, a difference in nicotine contents was observed among tea plant-growing regions. All these results strongly suggest that tea plants, similar to members of the Solanaceae, contain endogenous nicotine. Although low, the levels are higher than allowed MRLs. Despite these findings, the possibility of contamination of commercial tea by exogenous nicotine cannot be completely excluded.

Supporting information

S1 Table. Type, subtype, country, sampling year and number of tea samples used to measure nicotine in this study.

(DOCX)

S1 Fig. Manufacturing processes used to produce Sencha, pan-fired, oolong and black teas. (PDF)

S2 Fig. New leaves, new stems, mature leaves, mature stems, lignifying roots and white roots of tea plants cultured hydroponically. The white bar corresponds to 10 cm. (PDF)

S3 Fig. Flower buds and seeds (A), cultured tea cells (B) and aseptic tea seedlings (C) of Yabukita tea plants. (PDF)

S4 Fig. Nicotine contents of tea samples collected during first, second, rain/monsoon and autumn seasons from individual tea estates (designated as A to S) in India. The dotted line indicates the residue standard for nicotine ($0.01 \mu\text{g g}^{-1}$ dry weight). (PDF)

Acknowledgments

The all purchased tea samples were kindly collected by Dr. Masayuki Suzuki of Mitsui Norin Co., Ltd.

Author Contributions

Conceptualization: Takashi Ikka, Akio Morita.

Data curation: Takashi Ikka.

Formal analysis: Takashi Ikka, Fumio Nanjo.

Funding acquisition: Takashi Ikka, Akio Morita.

Investigation: Takashi Ikka, Hiroto Yamashita, Ikuya Kurita, Yasuno Tanaka, Akio Morita.

Methodology: Kazuya Takeda, Nobuhiro Horie, Hiroshi Hojo, Fumio Nanjo.

Project administration: Akio Morita.

Resources: Fumiya Taniguchi, Akiko Ogino.

Writing – original draft: Takashi Ikka, Akio Morita.

References

1. Lee KW, Lee HJ, Lee CY. Antioxidant activity of black tea versus green tea. *J Nutr Edu.* 2002; 132(4): 785.
2. Cabrera C, Artacho R, Giménez R. Beneficial effects of green tea—a review. *J Am Col Nutr.* 2006; 25(2): 79–99.
3. International Tea Committee (ITC). International Tea Committee: Annual Bulletin of Statistics. International Tea Committee, Ltd. 2015.
4. European food Safety Authority (EFSA). Setting of temporary MRLs for nicotine in tea, herbal infusions, spices, rose hips and fresh herbs. *EFSA Journal* 2011; 9(3): 50.
5. Sheen SJ. Detection of nicotine in foods and plant materials. *J Food Sci.* 1988; 53(5): 1572–1573.
6. Davis RA, Stiles MF, Reynolds JH. Dietary nicotine: a source of urinary cotinine. *Food Chem Toxicol.* 1991; 29(12): 821–827. PMID: [1765327](https://pubmed.ncbi.nlm.nih.gov/1765327/)
7. Domino EF, Hornbach E, Demana T. The nicotine content of common vegetables. *New Eng J Med.* 1993; 329(6): 437. <https://doi.org/10.1056/NEJM199308053290619> PMID: [8326992](https://pubmed.ncbi.nlm.nih.gov/8326992/)
8. Siegmund B, Leitner E, Pfannhauser W. Determination of the nicotine content of various edible nightshades (Solanaceae) and their products and estimation of the associated dietary nicotine intake. *J Agr Food Chem.* 1999; 47(8): 3113–3120.

9. Castro A, Monji N. Dietary nicotine and its significance in studies on tobacco smoking. *Biochemical Archives* 1986; 2(2): 91–97.
10. Andersson C, Wennström P, Gry J. Nicotine alkaloids in Solanaceous food plants. *TemaNord*. 2003.
11. European food Safety Authority (EFSA). Potential risks for public health due to the presence of nicotine in wild mushrooms. *EFSA Journal* 2009; RN-286: 1–47.
12. Petterson DS, Harris DJ, Allen DG. *Toxic Substances in Crop Plants*. Elsevier, (Chapter 7) 1991.
13. Konishi S, Miyamoto S, Taki T. Stimulatory effects of aluminum on tea plants grown under low and high phosphorus supply. *Soil Sci Plant Nutr*. 1985; 31(3): 361–368.
14. Kuboi T, Kaji M. Establishment of tea cell lines with high growth rate. *Tea Res J (Japan)* 1994; 80(1): 8.
15. Morita A, Yanagisawa O, Maeda S, Takatsu S, Ikka T. Tea plant (*Camellia sinensis* L.) roots secrete oxalic acid and caffeine into medium containing aluminum. *Soil Sci Plant Nutr*. 2011; 57(6): 796–802.
16. Cavalieri C, Bolzoni L, Bandini M. Nicotine determination in mushrooms by LC–MS/MS with preliminary studies on the impact of drying on nicotine formation. *Food Additives and Contaminants* 2010; 27(4): 473–477. <https://doi.org/10.1080/19440040903479768> PMID: 20234963
17. Thräne C, Isemer C, Engelhardt UH. Determination of nicotine in tea (*Camellia sinensis*) by LC–ESI–MS/MS using a modified QuEChERS method. *Euro Food Res Technol*. 2015; 241(2): 227–232.
18. Morita M, Shitan N, Sawada K, Van Montagu MCE., Inzé D, Rischer H, et al. Vacuolar transport of nicotine is mediated by a multidrug and toxic compound extrusion (MATE) transporter in *Nicotiana tabacum*. *Proc Nat Acad Sci USA* 2009; 106: 2447–2452. <https://doi.org/10.1073/pnas.0812512106> PMID: 19168636
19. Suzuki T, Yamazaki N, Sada Y, Oguni I, Moriyasu Y. Tissue distribution and intracellular localization of catechins in tea leaves. *Biosci Biotechnol Biochem*. 2003; 67: 2683–2686. <https://doi.org/10.1271/bbb.67.2683> PMID: 14730155
20. El-Ramady H, Abdalla N, Alshaal T, Elhenawy A, Amer M, Shehata A, et al. Contamination of Plant Foods with Nicotine: An Overview. *DBG Reports* 2015.
21. Gonzalez Alonso S, Valcarcel Y, Montero JC, Catala M. Nicotine occurrence in bottled mineral water: Analysis of 10 brands of water in Spain. *Sci Total Environ*. 2012; 416: 527–531. <https://doi.org/10.1016/j.scitotenv.2011.11.046> PMID: 22137651
22. Baldwin IT, Schmelz EA, Ohnmeiss TE. Wound-induced changes in root and shoot jasmonic acid pools correlate with induced nicotine synthesis in *Nicotiana sylvestris* spegazzini and comes. *J Chemical Ecology* 1994; 20(8): 2139–2157.
23. Baldwin IT, Zhang ZP, Diab N, Ohnmeiss TE, McCloud ES, Lynds GY, et al. Quantification, correlations and manipulations of wound-induced changes in jasmonic acid and nicotine in *Nicotiana sylvestris*. *Planta* 1997; 201(4): 397–404.
24. Shi Q, Li C, Zhang F. Nicotine synthesis in *Nicotiana tabacum* L. induced by mechanical wounding is regulated by auxin. *J Exp Bot*. 2006; 57(11), 2899–2907. <https://doi.org/10.1093/jxb/erl051> PMID: 16868042
25. Wang SS, Shi QM, Li WQ, Niu JF, Li CJ, Zhang FS. Nicotine concentration in leaves of flue-cured tobacco plants as affected by removal of the shoot apex and lateral buds. *J Integrative Plant Biol*. 2008; 50(8): 958–964.
26. Jin YF, Wang SS, Li JY, Pang T, Zhang JB, Gong M. Effects of different growth temperatures on alkaloid content and nicotine metabolism in roots, stems and leaves of tobacco (*Nicotiana tabacum* L.) plants. CORESTA Meeting, Agronomy/Phytopathology AP41, 2015.
27. Li F, Zhang H, Wang S, Xiao W, Ding C, Liu W, et al. Identification of Topping Responsive Proteins in Tobacco Roots. *Front Plant Sci*. 2016; 7: 1–11. <https://doi.org/10.3389/fpls.2016.00001>
28. Dawson RF. Alkaloid biogenesis. *Advances in Enzymology and Related Areas of Molecular Biology*, 1948; 8: 203–251.
29. Shoji T, Hashimoto T. Biosynthesis and regulation of tobacco alkaloids. *Herbaceous Plants: Cultivation Methods, Grazing and Environmental Impacts*, Nova Biomedical 2013; 37–67.
30. Michael AJ. Polyamines in Eukaryotes, Bacteria and Archaea. *J Biol Chem*. 2016; jbc-R116.
31. Hashimoto T, Yamada Y. Alkaloid biogenesis: molecular aspects. *Annual rev plant biol*. 1994; 45(1): 257–285.
32. Hibi N. Gene Expression in Tobacco Low-Nicotine Mutants. *Plant Cell* 1994; 6(5): 723–735. <https://doi.org/10.1105/tpc.6.5.723> PMID: 8038607
33. Chou WM, Kutchan TM. Enzymatic oxidations in the biosynthesis of complex alkaloids. *Plant J*. 1998; 15(3): 289–300. PMID: 9750342

34. Malmberg RL, Watson MB, Galloway GL, Yu W. Molecular genetic analyses of plant polyamines. *Critical Rev Plant Sci*. 1998; 17(2): 199–224.
35. Xu B, Sheehan MJ, Timko MP. Differential induction of ornithine decarboxylase (ODC) gene family members in transgenic tobacco (*Nicotiana tabacum* L. cv. Bright Yellow 2) cell suspensions by methyl-jasmonate treatment. *Plant Growth Regulation* 2004; 44(2): 101–116.
36. Kajikawa M, Shoji T, Kato A, Hashimoto T. Vacuolar- localized berberine bridge enzyme-like proteins are required for a late step of nicotine biosynthesis in tobacco. *Plant Physiol*. 2011; 155: 2010–2022. <https://doi.org/10.1104/pp.110.170878> PMID: 21343426
37. Shoji T, Nakajima K, Hashimoto T. Ethylene suppresses jasmonate-induced gene expression in nicotine biosynthesis. *Plant Cell Physiol*. 2000; 41: 1072–1076. PMID: 11100780
38. Farmer EE, Almeras E, Krishnamurthy V. Jasmonates and related oxylipins in plant responses to pathogenesis and herbivory. *Current Opinion Plant Biol*. 2003; 6: 372–378.