

Armillariols A to C from the culture broth of *Armillaria* sp.




メタデータ	言語: eng 出版者: 公開日: 2013-10-23 キーワード (Ja): キーワード (En): 作成者: Kobori, Hajime, Sekiya, Atsushi, Yasuda, Nobuhiro, Noguchi, Keiichi, Suzuki, Tomohiro, Choi, Jae-Hoon, Hirai, Hirofumi, Kawagishi, Hirokazu メールアドレス: 所属:
URL	http://hdl.handle.net/10297/7451

Graphical Abstract

Armillariols A to C from the culture broth of *Armillaria* sp.

Leave this area blank for abstract info.

Hajime Kobori, Atsushi Sekiya, Nobuhiro Yasuda, Keiichi Noguchi, Tomohiro Suzuki, Jae-Hoon Choi, Hirofumi Hirai, Hirokazu Kawagishi*





Armiliariols A to C from the culture broth of *Armillaria* sp.

Hajime Kobori^a, Atsushi Sekiya^b, Nobuhiro Yasuda^c, Keiichi Noguchi^d, Tomohiro Suzuki^e, Jae-Hoon Choi^e, Hirofumi Hirai^f, Hirokazu Kawagishi^{a,e,f,*}

^a Graduate School of Science and Technology, Shizuoka University, 836 Ohya, Suruga-ku, Shizuoka 422-8529, Japan

^b Kyushu Research Center, Forestry & Forest Products Research Institute, 4-11-16 Kurokami, Kumamoto, Kumamoto, 860-0862 Japan

^c Japan Synchrotron Radiation Research Institute, SPring-8, 1-1-1 Kouto, Sayo-cho, Sayo-gun, Hyogo 679-5198, Japan

^d Instrumentation Analysis Center, Tokyo University of Agriculture and Technology Koganei, Tokyo 184-8588, Japan

^e Research Institute of Green Science and Technology, Shizuoka University, 836 Ohya, Suruga-ku, Shizuoka 422-8529, Japan

^f Graduate School of Agriculture, Shizuoka University, 836 Ohya, Suruga-ku, Shizuoka 422-8529, Japan

ARTICLE INFO

Article history:

Received

Received in revised form

Accepted

Available online

ABSTRACT

Three novel compounds were isolated from the culture broth of *Armillaria* sp.. Their structures were elucidated mainly by spectroscopic data analyses. All the compounds regulated hypocotyl and root growth of lettuce.

2013 Elsevier Ltd. All rights reserved.

Keywords:

culture broth

fungus

Armillaria sp

lettuce growth inhibition

structure determination

Introduction

The genus *Armillaria* (English name, Honey Fungus; Japanese name, Naratake) belonging to the family Physalacriaceae is a well known group of edible mushroom throughout the world. People have utilized it for its medicinal properties and edible qualities for a long time. On the other hand, the genus has been also known as a serious plant pathogen which causes root rot in various plant species¹ and the phenomenon is called "Armillaria root disease"^{2,3}. The rot is one of the most serious diseases of plants, which occurs in many broadleaf trees and conifers, and several herbaceous plants including *Alchemilla mollis*, *Beta vulgaris* and so on⁴. Furthermore, it is known that penetration of *Armillaria* mycelia to the fungus *Entoloma abortivum* and *Wynnea americana* induces spherical deformity of the fruiting bodies of those mushrooms⁵. These facts mean that *Armillaria* produces allelopathic substance(s). Protoilludane sesquiterpene aryl esters have been isolated from the genus *Armillaria* mushrooms and *Clitocybe illudens*.⁶⁻¹² Those compounds showed antimicrobial activity^{7,13,14} and cytotoxicity against human cancer cell lines^{15,16}. However, there are no evidences that the compounds are toxic principles for "Armillaria root disease". Therefore, we tried to isolate compounds with plant growth regulatory activity from the fungus.

Here, we describe the isolation, structural determination, and biological activity of three novel compounds from the culture broth of a strain of the genus.




Fig. 1. Structures of **1** to **3**.

Results and discussion

Isolation of the active compounds from *Armillaria* sp. was guided by their growth-regulating activity on lettuce. Since culture broth of the fungus inhibited the growth, the broth were extracted with hexane, EtOAc, and 1-BuOH, successively. The active fractions, the hexane soluble- and the EtOAc soluble-fractions, were subjected to repeated chromatography respectively to afford armillariol A to C (**1-3**) (Fig. 1).

Armillariol A (**1**) was purified as colorless oil. The molecular formula was determined as C₁₅H₂₆O₂ by HR-ESI-MS *m/z* 261.1826 [M+Na]⁺ (calcd for C₁₅H₂₆NaO₂, 261.1830), indicating

* Corresponding author. Tel.: +81-54-238-4885; fax: +81-54-238-4885; e-mail: ahkawa@ipc.shizuoka.ac.jp (H. Kawagishi)

the presence of three degrees of unsaturation in the molecule. Structure of **1** was elucidated by interpretation of NMR spectra including DEPT, COSY, HMQC, and HMBC (Fig. 2). The DEPT experiment indicated the presence of four methyls, four methylenes, four methines and three quaternary carbons. The molecular formula, the unsaturation degrees, ^{13}C -NMR data (δ_{C} 72.6, 77.9, 120.8, 120.9, 133.8, 143.5, Table 1), and the DEPT data indicated the presence of a ring, two double bonds, and two hydroxy groups in the molecule. The complete assignment of the protons and carbons of NMR was accomplished as shown in Table 1. The structure of the alkyl chain (C1 to C7) was elucidated by the COSY correlations (H3/H4; H4/H5) and the HMBC correlations (H1'/C2, C2-CH₃, C3; H2-CH₃/C1, C2, C3; H3/C2-CH₃, C4, C5; H4/C5, C6; H5/C4, C7; H7/C5, C6) (Fig. 2). The presence of the cyclohexene moiety (C1' to C6') and its linking positions to a methyl and the chain were confirmed by the COSY correlations (H1'/H2', H6'; H2'/H3'; H5'/H6') and the HMBC correlations (H5'/C1', H7'/C1', H1'/C5'; H4'-CH₃/C3', C4', C5') (Fig. 2). NOE was observed between H4 and H1' in the NOESY and NOE-difference experiments, indicating that the geometry of the double bond was Z. As a result, structure of **1** was supposed to be (Z)-2-methyl-6-(4-methylcyclohex-3-enyl)hept-5-ene-2,3-diol (Fig. 1). 2-Methyl-6-(4-methylcyclohex-3-en-1-yl)hept-6-ene-2,3-diol (10,11-dihydroxy-10,11H- β -bisabolene) that has the same skeleton as **1** has been isolated from the aerial parts of the plant *Acritopappus confertus*¹⁷.




Fig. 2. COSY and HMBC correlations in **1** to **3**.

Table 1. ^1H and ^{13}C NMR data for **1-3** (**1** and **3** in CDCl_3 , **2** in CD_3OD).

position	1		position	2		position	3	
	^1H	^{13}C		^1H	^{13}C		^1H	^{13}C
	δ (mult., J in Hz)	δ		δ (mult., J in Hz)	δ		δ (mult., J in Hz)	δ
1	1.22 (s)	26.6	1		174.8	1		186.5
2		72.6	2	2.22 (d, 7.0)	44.1	2	2.43 (s)	25.9
3	3.36 (dd, 8.9, 4.0)	77.9	3	2.08 (m)	26.8	2'		152.0
4	2.16 (m)	29.8	4	0.95 (d, 6.7)	22.7	3'	7.12 (d, 3.7)	118.6
	2.20 (m)		3-CH ₃	0.95 (d, 6.7)	22.7	4'	6.47 (d, 3.7)	109.8
5	5.18 (dd, 7.3, 7.0)	120.9	1'	4.13 (m)	66.6	5'		159.6
6		143.5		4.16 (m)		1''	4.56 (br. s)	71.1
7	1.66 (br. s)	19.5	2'	3.82 (m)	70.4	2''	3.92 (m)	73.2
1'	2.60 (m)	35.4	3'	3.60 (m)	73.0	3''	1.45 (m)	33.0
2'	1.80 (m)	29.4	4'	3.60 (m)	64.1		1.52 (m)	
	2.02 (m)					4''	1.28 (m)	25.2
3'	5.38 (m)	120.8					1.47 (m)	
4'		133.8				5''	1.27 (m)	31.7
5'	1.92 (m)	30.5				6''	1.27 (m)	22.5
	2.05 (m)					7''	0.86 (t, 6.8)	14.0
6'	1.53 (m)	27.5						
	1.58 (m)							
2-CH ₃	1.16 (s)	23.7						
4'-CH ₃	1.64 (br. s)	23.6						

Armillariol B (**2**) was purified as pale yellow oil. Its molecular formula was determined as $\text{C}_9\text{H}_{18}\text{O}_5$ by HR-ESI-MS m/z 229.1066 $[\text{M}+\text{Na}]^+$ (calcd for $\text{C}_9\text{H}_{18}\text{NaO}_5$, 229.1052), indicating the presence of one degree of unsaturation in the molecule. Structure of **2** was elucidated by interpretation of NMR spectra including DEPT, COSY, HMQC, and HMBC (Fig. 2). The one unsaturation degree was explained by the presence of an ester (δ_{C} 173.6). The acid part, 3-methylbutanoate, was elucidated by the COSY correlations (H2/H3; H3/H4, H3-CH₃) and the HMBC correlations (H2/C1, C3, C4, C3-CH₃; H3/C2, C4, C3-CH₃, H4/C2, C3, C3-CH₃, H3-CH₃/C2, C3, C4) (Fig. 2). The COSY correlations (H1'/H2', H2'/H3', H3'/H4') and the HMBC correlations (H1'/C2', C3', C1; H2'/C1', C3', C4'; H3'/C1', C2'; H4'/C2', C3') indicated the structure of the alcohol part and its linkage position to the acid part. As a consequence, **2** was determined as 2,3,4-trihydroxybutyl 3-methylbutanoate (Fig. 1).

Armillariol C (**3**) was purified as pale yellow oil. Its molecular formula was determined as $\text{C}_{13}\text{H}_{20}\text{O}_4$ by HR-ESI-MS m/z 263.1224 $[\text{M}+\text{Na}]^+$ (calcd for $\text{C}_{13}\text{H}_{20}\text{NaO}_4$, 263.1259), indicating that the unsaturation degrees in the molecule was four. Structure of **1** was elucidated by interpretation of NMR spectra including DEPT, COSY, HMQC, and HMBC (Fig. 2). The DEPT experiment indicated the presence of two methyls, four methylenes, four methines and three quaternary carbons. The single carbon chain from C1'' to C7'' was constructed by the ^1H - and ^{13}C -NMR (Table 1) spectra along with COSY correlations (H3'/H4', H1''/H2'', H2''/H3'', H3''/H4'', H4''/H5'', H6''/H7'') and HMBC ones (H2'/C1, C2'; H3'/C-1, C2', C4', C5'; H4'/C2', C3', C5'; H1''/C4', C5', C2'; H2''/C3'', C4''; H3''/C1'', C2'', C4'', C5''; H4''/C2'', C5'', C6''; H5''/C4'', C6'', C7''; H6''/C4'', C5'', C7''; H7''/H5'', H6''). The molecular formula, the presence of a carbonyl-conjugated diene (C1, δ_{C} 185.5; C2', δ_{C} 152.0; C3', δ_{C} 118.6, δ_{H} 7.12 (d, J = 3.7 Hz); C4', δ_{C} 109.8, δ_{H} 6.47 (d, J = 3.7 Hz); C5', δ_{C} 159.6) including two quaternary sp^2 carbons, and the unsaturation degree indicated that this compound possessed a furan. As a result, structure of **3** was determined to be 1-(5-(1,2-dihydroxyheptyl)furan-2-yl)ethanone (Fig. 1).

Confirmation of the planar structures and determination of stereochemistry of **2** and **3** were performed by X-ray




Fig. 3. ORTEP drawings of *p*-bromobenzoate of **2** and **3** with ellipsoids at the 30% probability level. Hydrogen atoms are shown as small spheres of arbitrary radii.

crystallography analysis of their (*p*-bromo)benzoates (Fig. 3). The Flack parameter ($X = 0.16$ (**3**)) of **2** in the analysis indicated that its absolute configuration might be as shown in Fig. 1. However, the possibility of the opposite configuration could not be excluded. Therefore, the relative configuration of **2** and absolute configuration of **3** were determined to be as shown in Fig. 1. The stereochemistry at the asymmetric carbons in **1** remains undetermined.

In order to examine plant growth regulatory activity of these compounds, lettuce was used as the test plant. All the compounds inhibited hypocotyl and root growth of the plant at $1 \mu\text{mol}/\text{paper}$ with significant differences (rate of growth length compared with control \pm standard deviation: **1**, root $81.2 \pm 11.9\%$, hypocotyl $55.5 \pm 21.9\%$; **2**, root $68.0 \pm 24.6\%$, hypocotyl $66.3 \pm 21.5\%$; **3**, root $68.3 \pm 25.9\%$, hypocotyl $66.5 \pm 21.8\%$). **1** promoted the hypocotyl growth at $10^{-2} \mu\text{mol}/\text{paper}$ ($114 \pm 9.15\%$).

Since *Armillaria* species mainly infect woody plants and then cause "Armillaria root disease" to them^{2,3}. Woody plants should have been used in the bioassay. However, such a assay is very difficult for us because of low quantities of the compounds. Occasionally several herbaceous plants are also damaged by those pathogens⁴. Therefore, the activity of the compounds toward lettuce suggested that armillariols A to C might play some roles in *Armillaria* root disease.

Acknowledgment

We thank V. K. Deo (Shizuoka University) for valuable discussion and the synchrotron radiation experiment was

performed at the BL02B1 of SPring-8 with the approval of the Japan Synchrotron Radiation Research Institute (JASRI) (Proposal No. 2012B1109). This work was partially supported by a Grant-in-Aid for Scientific Research on Innovative Areas "Chemical Biology of Natural Products" from MEXT.

References and notes

1. Roll-Hansen, F. *Eur. J. For. Pathol.* **1985**, *15*, 22-31.
2. Cox, K. D.; Scherm, H. *Biol. Control* **2006**, *37*, 291-300.
3. Thomidis, T.; Exadaktylou, E. *Crop Prot.* **2012**, *36*, 49-51.
4. Robinson-Bax, C.; Fox, R. T. V. *Mycologist* **2002**, *16*, 21-22.
5. Ando, Y. *Chiba Mycol. Club Bull.* **2000**, *16-17*, 19-25 (in Japanese).
6. Ayer, W. A.; Browne, L. M. *Tetrahedron* **1981**, *37*, 2197-2248.
7. Midland, S. L.; Izak, R. R.; Wing, R. M.; Zaki, A. I.; Munnecke, D. E.; Sims, J. J. *Tetrahedron Lett.* **1982**, *23*, 2515-2518.
8. Arnone, A.; Cardillo, R.; Nasini, G. *Phytochemistry* **1986**, *25*, 471-474.
9. Donnelly, D. M. X.; Quigley, P. F.; Coveney, J. D.; Polonsky J. *Phytochemistry* **1987**, *26*, 3075-3077.
10. Donnelly, D. M. X.; Hutchinson, R. M.; Coveney, J. D.; Yonemitsu, M. *Phytochemistry* **1990**, *29*, 2569-2572.
11. Yang, J.-S.; Su, Y.-L.; Yu, D.-Q.; Liang, X.-T. *J. Chinese Pharm. Sci.* **1993**, *2*, 10-17.
12. Donnelly, D. M. X.; Konishi, T.; Dunne, O.; Cremin, P. *Phytochemistry* **1997**, *44*, 1473-1478.
13. Oduro, K. A.; Munnecke, D. E.; Sims, J. J.; Keen, N. T. *Trans. Br. Mycol. Soc.* **1976**, *66*, 195-199.
14. Peipp, H.; Sonnenbichler, J. *Biol. Chem. Hoppe-Seyler* **1992**, *373*, 675-683.
15. Misiek, M.; Williams, J.; Schmich, K.; Hüttel, W.; Merfort, I.; Salomon, C. E.; Aldrich, C. C.; Hoffmeister, D. *J. Nat. Prod.* **2009**, *72*, 1888-1891.
16. Bohnert, M.; Miethbauer, S.; Dahse, H.-M.; Ziemer, J.; Nett, M.; Hoffmeister, D. *Bioorg. Med. Chem. Lett.* **2011**, *21*, 2003-2006.
17. Bohlmann, F.; Zdero, C.; Jakupovic, J.; King, R. M.; Robinson H. *Phytochemistry* **1983**, *22*, 2243-2252.
18. Sugimoto, K.; Ohsumi, H.; Aoyagi, S.; Nishibori, E.; Moriyoshi, C.; Kuroiwa, Y.; Sawa, H. and Takata, M. *AIP Conf. Proc.*, **2010**, *1234*, 887-890.

Supplementary Material

Supplementary data associated with this article can be found, in the online version, at:

[Click here to remove instruction text...](#)