

Improving xylitol production through  
recombinant expression of xylose reductase in  
the white-rot fungus *Phanerochaete sordida*  
YK-624

メタデータ	言語: eng 出版者: 公開日: 2015-01-05 キーワード (Ja): キーワード (En): 作成者: Hirabayashi, Sho, Wang, Jianqiao, Kawagishi, Hirokazu, Hirai, Hirofumi メールアドレス: 所属:
URL	<a href="http://hdl.handle.net/10297/8000">http://hdl.handle.net/10297/8000</a>

1 Note

2 **Improving xylitol production through recombinant expression of xylose reductase**  
3 **in the white-rot fungus *Phanerochaete sordida* YK-624**

4 Sho Hirabayashi <sup>1, †</sup>, Jianqiao Wang <sup>1, †</sup>, Hirokazu Kawagishi <sup>1, 2, 3</sup>, Hirofumi Hirai <sup>1, 2 \*</sup>

5 <sup>1</sup> Graduate School of Agriculture, Shizuoka University, 836 Ohya, Suruga-ku, Shizuoka  
6 422-8529, Japan

7 <sup>2</sup> Research Institute of Green Science and Technology, Shizuoka University, 836 Ohya,  
8 Suruga-ku, Shizuoka 422-8529, Japan

9 <sup>3</sup> Graduate School of Science and Technology, Shizuoka University, 836 Ohya,  
10 Suruga-ku, Shizuoka 422-8529, Japan

11 \* Corresponding author. Department of Applied Biological Chemistry, Graduate School  
12 of Agriculture, Shizuoka University, 836 Ohya, Suruga-ku, Shizuoka 422-8529, Japan.

13 Tel. & fax: +81 54 238 4853

14 *E-mail address*: ahhirai@ipc.shizuoka.ac.jp (H. Hirai).

15

16 † These authors contributed equally to this work.

17 **Keywords**: white-rot fungi, *Phanerochaete sordida* YK-624, xylose reductase,  
18 recombinant expression, xylitol

19 **Running title**: Improving xylitol production by XR gene expression

## Abstract

20        We generated an expression construct consisting of the xylose reductase (XR) gene  
21    (*xr*) from *P. chrysosporium*. Transformant X7 exhibited increased xylitol production and  
22    markedly higher XR activities than the wild-type strain. RT-PCR analysis demonstrated  
23    that the increased XR activity was associated with constant expression of the  
24    recombinant *xr* gene.

25 Lignocellulosic biomass is the most abundant organic compound and has  
26 therefore attracted worldwide interest as a feedstock for the production of bioethanol (1,  
27 2). For the cost-effective production of value-added products from renewable  
28 lignocellulosic resources, microbial bioconversion processes must effectively utilize the  
29 pentose sugar xylose, as it is a major component of lignocellulose hydrolysates (3-5).  
30 However, the yeast *Saccharomyces cerevisiae*, which is typically used in ethanol  
31 production from hexoses, is unable to use pentose sugars such as a xylose (6, 7).  
32 Therefore, the development of a microbial-based system that can directly utilize pentose  
33 is necessary for the economic conversion of lignocellulose in biorefinery processes.

34 In yeasts and fungi, typically convert xylose to xylulose in a two-step reaction  
35 mediated by two kinds of oxidoreductases. An NADPH xylose reductase (XR) first  
36 reduces xylose to xylitol, which is then oxidized to xylulose by an NAD-linked xylitol  
37 dehydrogenase. And, improvement of xylitol production by recombinant  
38 microorganisms was received much attention (4, 8). Thus, the simultaneous production  
39 of ethanol and xylitol would significantly increase the efficiency of bioethanol  
40 production from lignocellulosic biomass.

41 The white-rot fungus *Phanerochaete sordida* YK-624, which was originally  
42 isolated from rotten wood, exhibits greater ligninolytic activity and selectivity than

43 either *P. chrysosporium* or *Trametes versicolor* (9). In our previous studies, we  
44 successfully developed a superior lignin-degrading strain of *P. sordida* YK-624 using a  
45 molecular breeding approach with a homologous expression system (10, 11). Here, we  
46 used a similar approach to generate high XR gene (*xr*)-expressing transformants of *P.*  
47 *sordida* YK-624 and investigated whether xylitol production was improved in these  
48 transformants.

49 Genomic DNA of *P. chrysosporium* ME-446 was extracted from mycelia using  
50 ISOPLANT II (Nippon Gene, Tokyo) and was then used as template for the PCR  
51 amplification of the full-length genomic *xr* gene (1144 bp) using the specific primers  
52 PcXRF1 and PcXRR1. The obtained PCR product was ligated into the cloning vector  
53 pMD20-T (Takara Bio, Shiga, Japan) and introduced into *Escherichia coli* DH5 $\alpha$  for  
54 sequencing. Primers XRF1 and XRR1 were designed to amplify the *xr* gene and  
55 introduce an *Xba* I site for cloning into the expression plasmid p*PsGPD-pro*, which was  
56 generated in our previous study (12). The amplified DNA fragment was digested with  
57 *Xba* I and cloned into *Xba* I-digested p*PsGPD-pro*, yielding plasmid p*PsGPD-XR*.

58 After p*PsGPD-XR* was sequenced to verify the absence of PCR errors, it was  
59 co-transformed with p*PsURA5* using the PEG method into UV-64 protoplasts that were  
60 prepared by a standard technique using cellulases (13). Co-transformed clones (X

61 strains) were identified by PCR with the primers *gpdF1* and *XRR2*, which were  
62 designed to amplify the recombinant *xr* gene. A total of 13 strains (X1-13) that were  
63 co-transformed with *pPsGPD-XR* and *pPsURA5* were obtained.

64 The effect of recombinant *xr* expression on xylitol production by the X strains was  
65 next investigated. *P. sordida* YK-624 (WT) and the X strains were incubated on PDA  
66 plates at 30 °C for 3 days, and 10-mm diameter disks were then punched out from the  
67 growing edge of the mycelia using a sterile cork borer. Two mycelial disks for each  
68 strain were placed into a 100-mL Erlenmeyer flask containing 10 mL nitrogen-limited  
69 Kirk medium (14) supplemented with 1.5% xylose as a carbon source, and the flask was  
70 then statically incubated at 30 °C for 7 days. Xylitol and xylose concentrations were  
71 determined using a high-performance liquid chromatograph equipped with a Shodex  
72 SH1821 column (8.0 mm × 300 mm, Showa Denko K.K., Tokyo, Japan) at 75 °C with  
73 0.5 mM H<sub>2</sub>SO<sub>4</sub> as the mobile phase at a flow rate of 0.6 mL min<sup>-1</sup>. Xylitol and xylose  
74 were detected using an online refractive index detector. Transformant strain X7  
75 exhibited the highest xylitol production among the 13 transformant strains (data not  
76 shown) and was therefore selected for subsequent analyses.

77 We next analyzed the transcription levels of *xr* by RT-PCR and measured the XR  
78 activity of strain X7 during 8 days of culture in nitrogen-limited Kirk medium. Mycelial

79 samples were collected and stored at -80 °C. Total RNA was isolated from the thawed  
80 samples using an RNeasy Plant Mini kit (Qiagen, Valencia, CA). RT-PCR was  
81 performed using 200 ng of total RNA with a PrimeScript RT-PCR Kit (Takara Bio) and  
82 the gene-specific primer sets *gpdF1-XRR2* and *ActinF-ActinR*.

83 For the measurement of XR activity, each mycelial sample was added to 2 ml of  
84 250 mM phosphate buffer (pH 7.0) and was then homogenized using a Polytron  
85 PT1200E (Kinematica, Canada) at 4 °C. The homogenate was centrifuged (4 °C, 10,000  
86 x g, 10 min) and the obtained supernatant was used as a cell-free extract for the  
87 measurement of XR activity, which was determined by monitoring the oxidation of  
88 NADPH to NADP<sup>+</sup> ( $\epsilon_{340} = 6.22 \text{ mM}^{-1} \text{ cm}^{-1}$ ) (15). The reaction mixture (1 ml) consisted  
89 of 10 mM 2-mercaptoethanol, 50 mM xylose, and 0.17 mM NADPH in 250 mM  
90 phosphate buffer (pH 7.0).

91 The time course of xylitol production by strains X7 and WT over 2, 4, 6, and 8  
92 days of incubation in nitrogen-limited Kirk medium was monitored. After 4 days of  
93 incubation, xylitol production by strain X7 reached 1.39 g L<sup>-1</sup>, whereas the WT strain  
94 had only produced 0.61 g L<sup>-1</sup> (Fig. 1). In contrast to the WT strain, the production of  
95 xylitol by strain X7 continued to increase, reaching 3.61 g L<sup>-1</sup> of xylitol after 8 days of  
96 incubation (Fig. 1). We next evaluated the consumption rates by the xylose

97 consumption, and the xylose consumption rate of the strain X7 (31.8%) was higher than  
98 that of WT strain (11.2%).

99 XR activity of strain X7 increased gradually for 8 days and was higher than that of  
100 WT throughout the incubation period (Fig. 2a). The highest XR activity for strain X7  
101 was 9.14 nkat flask<sup>-1</sup> after 8 days of incubation, whereas that for WT was 6.54 nkat  
102 flask<sup>-1</sup> (Fig. 2a). The difference of fungal grown in WT strain and strain X7 was not  
103 observed under glucose- or xylose-containing medium. Consistent with these findings,  
104 RT-PCR indicated that strain X7 had constant *xr* expression between days 2 to 8 (Fig.  
105 2b). Taken together, these results suggest that the increase of XR activity in strain X7  
106 was due to constant expression of the recombinant *xr* gene, and that the recombinant  
107 expression of *xr* in the white-rot fungus *P. sordida* YK-624 improves xylitol production.

108 To conclude, the recombinant expression of *xr* is effective for the improvement of  
109 the xylitol production in white-rot fungi. Recently, direct ethanol production by the  
110 white-rot fungus *Phlebia* sp. MG-60 in pure culture without the addition of exogenous  
111 chemicals or enzymes was reported (16). In future studies, we intend to develop a  
112 molecularly bred strain of white-rot fungus that can simultaneously produce ethanol and  
113 xylitol from woody biomass with high yields.

114



115 This work was partially supported by Grants-in-Aid for Scientific Research (A)  
116 (Nos. 21248023 and 24248030) from the Ministry of Education, Culture, Sports,  
117 Science and Technology of Japan.

## 118 **Reference**

- 119 1. FitzPatrick, M., Champagne, P., Cunningham, M.F., and Whitney, R.A.: A  
120 biorefinery processing perspective: treatment of lignocellulosic materials for the  
121 production of value-added products, *Bioresour. Technol.*, **101**, 8915–8922 (2010).
- 122 2. Lin, Y. and Tanaka, S.: Ethanol fermentation from biomass resources: current state  
123 and prospects, *Appl. Microbiol. Biotechnol.*, **69**, 627–642 (2006).
- 124 3. Sasaki, M., Jojima, T., Inui, M., and Yukawa, H.: Xylitol production by  
125 recombinant *Corynebacterium glutamicum* under oxygen deprivation, *Appl.*  
126 *Microbiol. Biotechnol.*, **86**, 1057–1066 (2010).
- 127 4. Kötter, P., and Ciriacy, M.: Xylose fermentation by *Saccharomyces cerevisiae*, *Appl.*  
128 *Microbiol. Biotechnol.*, **38**, 776–783 (1993).
- 129 5. Eliasson, A., Christensson, C., Wahlbom, C.F., and Hahn-Hägerdal, B.: Anaerobic  
130 xylose fermentation by recombinant *Saccharomyces cerevisiae* carrying *XYL1*,  
131 *XYL2*, and *XKSI* in mineral medium chemostat cultures, *Appl. Environ. Microbiol.*,  
132 **66**, 3381–3386 (2000).

- 133 6. Vallet, C., Said, R., Rabiller, C., and Martin, M.L.: Natural abundance isotopic  
134 fractionation in the fermentation reaction: influence of the nature of the yeast,  
135 Bioorg. Chem., **24**, 319–330 (1996).
- 136 7. Leticia, P., Miguel, C., Humberto, G., and Jaime, A.J.: Fermentation parameters  
137 influencing higher alcohol production in the tequila process, Biotechnol. Lett., **19**,  
138 45–47 (1997).
- 139 8. Harcus, D., Dignard, D., Lépine, G., Askew, C., Raymond, M., Whiteway, M., and  
140 Wu, C.: Comparative xylose metabolism among the Ascomycetes *C. albicans*, *S.*  
141 *stipitis* and *S. cerevisiae*, PLoS One, **8**, DOI: 10.1371/journal.pone.0080733 (2013).
- 142 9. Hirai, H., Kondo, R., and Sakai, K.: Screening of lignin degrading fungi and their  
143 ligninolytic enzyme activities during biological bleaching of kraft pulp. Mokzai  
144 Gakkaishi, **40**, 980–986 (1994).
- 145 10. Sugiura, T., Mori, T., Kamei, I., Hirai, H., Kawagishi, H., and Kondo, R.:  
146 Improvement of ligninolytic properties in the hyper lignin-degrading fungus  
147 *Phanerochaete sordida* YK-624 using a novel gene promoter, FEMS Microbiol.  
148 Lett., **331**, 81–88 (2012).
- 149 11. Sugiura, T., Yamagishi, K., Kimura, T., Nishida, T., Kawagishi, H., and Hirai, H.:  
150 Cloning and homologous expression of novel lignin peroxidase genes in the

- 151 white-rot fungus *Phanerochaete sordida* YK-624, Biosci. Biotechnol. Biochem., **73**,  
152 1793–1798 (2009).
- 153 12. Suzuki, T., Dohra, H., Omae, S., Takeshima, Y., Choi, J.H., Hirai, H., and  
154 Kawagishi, H.: Heterologous expression of a lectin from *Pleurocybella porrigens*  
155 (PPL) in *Phanerochaete sordida* YK-624, J. Microbiol. Methods, **100**, 70–76  
156 (2014).
- 157 13. Yamagishi, K., Kimura, T., Oita, S., Sugiura, T., and Hirai, H.: Transformation by  
158 complementation of a uracil auxotroph of the hyper lignin-degrading basidiomycete  
159 *Phanerochaete sordida* YK-624, Appl. Microbiol. Biotechnol., **76**, 1079–1091  
160 (2007).
- 161 14. Tien, M., and Kirk, T.K.: Lignin peroxidase of *Phanerochaete chrysosporium*,  
162 Methods Enzymol., **161**, 238–249 (1988).
- 163 15. Yokoyama, I., Suzuki, T., Kawai, K., Horitsu, H., and Takamizawa, K.: Purification,  
164 characterization and structure analysis of NADPH-dependent d-xylose reductases  
165 from *Candida tropicalis*, J. Ferm. Bioeng., **79**, 217–223 (1995).
- 166 16. Kamei, I., Hirota, Y., and Meguro, S.: Integrated delignification and simultaneous  
167 saccharification and fermentation of hard wood by a white-rot fungus, *Phlebia* sp.  
168 MG-60, Bioresour. Technol., **126**, 137–141 (2012).

Table 1 Oligonucleotides used as primers in this study.

Primer name	Nucleotide sequence (5'-sequence-3')
PcXRF1	ATGCTTTCTTCGCCAACCCCTC
PcXRR1	TCAGCATTACGGAGAAAGTACG
XRF1	AATCTAGAATGCTTTCTTCGCCAACCCCTC
XRR1	AATCTAGATCAGCATTACGGAGAAAGTACG
gpdF1	AAGCAGCGAGGATTGTACC
XRR2	GATGGAAGGTGTTCCACAG
ActinF	AGCACGGTATCGTCACCAAC
ActinR	AGCGAAACCCTCGTAGATGG




Fig. 1 Time course of xylitol production by the wild-type (WT) strain (closed diamonds) and strain X7 (closed squares). Values are the means  $\pm$  SD of triplicate samples. Asterisks indicate values that were determined by the Student's t-test to be significantly different from WT (\* $P$ <0.05, \*\* $P$ <0.01).





Fig. 2 XR activity and expression of the *xr* gene by strain X7. a) Time course of XR activities in the wild-type strain (closed diamonds) and strain X7 (closed squares). Values are the means  $\pm$  SD of triplicate samples. Asterisks indicate values that were determined by the Student's t-test to be significantly different from WT (\* $P < 0.05$ , \*\* $P < 0.01$ ). b) RT-PCR analysis of *xr* gene transcription in strain X7. The expression of *actin* was used as an internal control.