

Efficient removal of perchlorate ion from water by a water-insoluble M2L4type compound

メタデータ	言語: eng 出版者: 公開日: 2015-11-02 キーワード (Ja): キーワード (En): 作成者: Mochizuki, Masaru, Inoue, Tatsunari, Yamanishi, Katsunori, Koike, Shiori, Kondo, Mitsuru, Zhang, Liang, Aoki, Hiromi メールアドレス: 所属:
URL	http://hdl.handle.net/10297/9211

Cite this: DOI: 10.1039/c0xx00000x

www.rsc.org/dalton

COMMUNICATION

Efficient Removal of Perchlorate Ion from Water by Water-insoluble M₂L₄ Type CompoundMasaru Mochizuki,^a Tatsunari Inoue,^a Katsunori Yamanishi,^b Shiori Koike,^a Mitsuru Kondo,^{*,a,c} Liang Zhang,^d and Hiromi Aoki^d⁵ Received (in XXX, XXX) Xth XXXXXXXXXX 20XX, Accepted Xth XXXXXXXXXX 20XX

DOI: 10.1039/b000000x

Reaction of CuSO₄·5H₂O with 1,3-bis(benzimidazol-1-ylmethyl)-2,4,6-trimethylbenzene (*m*-bbitrb) provides a new positively charged M₂L₄ type molecule, [SO₄ ⊂ Cu₂(*m*-bbitrb)₄]SO₄. This compound includes SO₄²⁻ inside and outside the cationic cage. Despite this compound being insoluble in water, it shows high efficient perchlorate removal activity from water by the exchange with SO₄²⁻ outside the cage.

Materials which show anion-exchange property have attracted intense attentions as useful methods for removal of toxic anions from water. One of the recent tough targets of the anions that should be removed from water is perchlorate ion (ClO₄⁻). The anion interferes with the thyroid's uptake of iodide ion, which is an essential component of growth hormones produced in the thyroid gland. Therefore, ingestion of excess ClO₄⁻ would seriously affect the growth of newborns and children.¹ In 2005, the U.S. Environmental Protection Agency (EPA) announced that the acceptable exposure level of this anion was 24.5 ppb in drinking water.² However, reduction of the concentration of the anion to the low concentration level in a short period is difficult because of its high solubility.³

Numerous coordination compounds with M₂L₄ cationic cages have been reported to date.⁴ Their cages include various guest molecules such as anions,⁵⁻⁷ neutral molecules,⁸ and metal complexes.⁹ Among them, the self-assembled M₂L₄ cages which are constructed by metal ions and bis(imidazole)- or bis(benzimidazole)-type neutral ligands have attracted attention as molecular capsules for anions.^{6,7} Many of them have been isolated with hydrophobic anions such as ClO₄⁻, BF₄⁻, and CF₃SO₃⁻ within the cationic cages. This is because the hydrophobic anions assist the self-assembled construction of the M₂L₄ frameworks.^{6,7,10} In contrast, M₂L₄-type molecular capsules including sulfate ion, which is a harmless anion, were not known for the series of coordination compounds constructed by bis(imidazole)- or bis(benzimidazole)-type ligands.

We have recently shown that ClO₄⁻ in aqueous solutions can be removed by encapsulating it in a water-insoluble M₂L₄ molecular capsule.¹¹ This method, however, could not remove ClO₄⁻ from water to below 1 ppm concentration level. On continuing works, we have succeeded in isolation of a new water-insoluble M₂L₄ type compound that includes SO₄²⁻ inside and outside the cage. This compound removed ClO₄⁻ from water efficiently. Here, we report the synthesis, structure, and the high

perchlorate removal activities from water of the compound.

50




Figure 1. Molecular structure of **1-SO₄**. Disorders of SO₄²⁻ inside the cage, and hydrogen atoms are omitted for clarity. Color code: 55 blue, copper; red, oxygen; yellow, sulfur; sky-blue, nitrogen; black, carbon.

Compound **1-SO₄** was obtained as blue microcrystals by evaporation of an EtOH/H₂O mixed solution of CuSO₄·5H₂O and *m*-bbitrb on heating. Single crystals for X-ray diffraction analysis were grown by slow evaporation of an EtOH/H₂O solution of the product in a test tube for a few months. Figure 1 shows the M₂L₄ type capsule structure of **1-SO₄** bearing SO₄²⁻ inside and outside the cage. The SO₄²⁻ inside the cage associates with the two Cu²⁺ with Cu—O distances of 2.143 (8) Å. These distances are shorter than the Cu—O distance (2.388 (4) Å) between ClO₄⁻ and Cu²⁺ in [ClO₄ ⊂ Cu₂(*m*-bbitrb)₄](ClO₄)₂(ClO₄) (**2**) that was reported by Su and co-workers.⁶ The coordination of SO₄²⁻ pulls the Cu²⁺ toward the center of the capsule. The Cu²⁺ of **1-SO₄** is withdrawn about 0.202 (2) Å to the center of the capsule from the N₄ plane that is defined by four nitrogen atoms at the coordination condition, while Cu²⁺ of **2** remains in the N₄ plane. As a result the Cu····Cu distance of the M₂L₄ cage of **1-SO₄** (about 6.6 Å) is shorter than that of **2** (7.36 Å). This is the first M₂L₄-type molecule capturing divalent anions in M₂L₄ cationic cages constructed by bis(imidazole)- or bis(benzimidazole)-type ligands.

It was found that **1-SO₄** has high activity for removal of ClO₄⁻ ion from water despite the compound being insoluble in water. Figure 2a shows changes the concentrations of anions in the

aqueous solution as a function of time after addition of 0.20 g (0.10 mmol) of **1-SO₄** to an aqueous solution (100 mL) containing NaF, NaCl, NaBr, NaNO₃, NaClO₄, and Na₂SO₄ at the same concentration (1.0 mM). The change in the concentration of each anion was monitored using ion chromatography. As shown in the chart, 1.0 mM ClO₄⁻ was reduced to about 0.070 mM at 5 min, and then 0.013 mM within 30 min. Other anions, NO₃⁻ and Br⁻, were also reduced from 1.0 mM to about 0.55 mM and 0.80 mM, respectively, at 30 min. Concentrations of Cl⁻ and F⁻ were not changed essentially. The concentration of SO₄²⁻ was increased from 1.0 mM to 1.73 mM at 30 min. This increase corresponds to the decreases of concentrations of other anions, indicating that SO₄²⁻ of **1-SO₄**, which is insoluble in water, was exchanged with the anions during the reaction.




Figure 2. Changes of concentration of anions in an aqueous solution (100 mL), containing NaF, NaCl, NaBr, NaNO₃, NaClO₄, and Na₂SO₄ (all 0.1 mmol) after addition of **1-SO₄** (0.20 g, 0.1 mmol) (a) and anion-exchange resin Purolite A530E (0.20 g) (b).

To estimate the ClO₄⁻ removal ability of **1-SO₄**, a similar experiment was carried out for comparison using an anion-exchange resin (Purolite A530E),¹² which has been widely used for ClO₄⁻ removal.¹³ This study showed that 1.0 mM ClO₄⁻ was reduced to about 0.42 mM at 30 min and 0.15 mM at 180 min (Figure 2b). Three other anions, NO₃⁻, Br⁻, and SO₄²⁻, were reduced from 1.0 mM to 0.71 mM, 0.79 mM, and 0.91 mM, respectively, at 30 min. The concentration of Cl⁻, which is originally included in the resin, increased corresponding to the decreases of concentrations of other anions. This result shows that **1-SO₄** efficiently removes ClO₄⁻ from water. This ability could arise from the hydrophobic space created among the [SO₄ ⊂ Cu₂(*m*-bbitr_b)₄]²⁺ units, which encapsulates the hydrophobic anion ClO₄⁻ in preference to SO₄²⁻. This study showed that **1-SO₄** can remove ClO₄⁻ effectively from aqueous solutions even in the presence of other anions.

To study the anion exchange reaction shown by Figure 2a, structure of the reaction product was determined by single crystal X-ray analysis technique at 173 K (Figure 3) and elemental analysis (Supporting Information). Immersion of **1-SO₄** in an aqueous solution that contained excess NaClO₄ for a few days provided a blue powder. The solution remained colorless throughout the reaction. After careful washing of the reaction product with water, the obtained product was recrystallized from MeCN. The single-crystal X-ray diffraction study confirmed that the product was [SO₄ ⊂ Cu₂(*m*-bbitr_b)₄](ClO₄)₂·4MeCN (**1-ClO₄**). The overall structure is shown in Figure 3. Molecular structure of the cationic unit of **1-ClO₄** is essentially same to that of **1-SO₄**. This result clearly demonstrates that only SO₄²⁻ outside the cage was exchanged with ClO₄⁻ by the anion-exchange reactions. Compound **1-ClO₄** was also insoluble in water, meaning that ClO₄⁻ was separated from the aqueous solution by treatment with **1-SO₄**.




Figure 3. Molecular structure of **1-ClO₄**. Disorders of SO₄²⁻ inside the cage, ClO₄⁻ outside the cage, and hydrogen atoms are omitted for clarity. Color code: blue, copper; red, oxygen; yellow, sulfur; sky-blue, nitrogen; black, carbon; green, chlorine.

Although the solution in Figure 2a remained colorless throughout the experiment, the concentration of Cu²⁺ dissolved in the solution was determined using inductively coupled plasma–optical emission spectroscopy (ICP–OES). This measurement detected only 0.13 mg L⁻¹ of Cu²⁺ in the aqueous solutions after the ClO₄⁻ removal experiment. This value is lower than the concentration in drinking water mandated by the World Health Organization (2 mg L⁻¹).¹⁴ These results show that **1-SO₄** has high potential as a material for removal of ClO₄⁻ from water for many purposes.

Since the ClO₄⁻ removal speed by **1-SO₄** was very fast, we have attempted to prepare a **1-SO₄**-fixed filter. The filter was made from a commercially available syringe filter (ϕ 40 mm, 0.2 μm pores). The fixation of **1-SO₄** on the membrane was achieved by filtration of a powder sample of **1-SO₄** suspended in water using the syringe filter. After the aqueous solution of NaClO₄ (ca. 1,000 ppb for ClO₄⁻) was passed through the filter under hand-pressure, it was found that the concentration of ClO₄⁻ was reduced to ca. 12 ppb for ClO₄⁻. The ion chromatography charts are shown in Figure 4. This result means that the **1-SO₄** is useful for convenient removal of ClO₄⁻ from water in a short time period.




Figure 4. Ion chromatograph charts of the aqueous solution of NaClO_4 (ca. 1000 ppb) (6.0 mL) before (a) and after (b) passing through the syringe filter fixed with **1-SO₄**.

5

Although removal of ClO_4^- by anion-exchange resins is currently a promising method, the resins are generally difficult to regenerate for reuse.¹⁵ We confirmed that **1-SO₄** is conveniently regenerated for reuse after the ClO_4^- removal experiments. As summarized in the Supporting Information (p. S18), analytically pure *m*-bbitr_b was recovered by a general procedure from the reaction products obtained after ClO_4^- removal experiments in about 53% yield. The ligand could be used for re-preparation of **1-SO₄**.

15

In conclusion, we have prepared a new water-insoluble M_2L_4 type compound that includes SO_4^{2-} inside and outside the cage. The compound removed ClO_4^- from water by exchange with SO_4^{2-} outside the cage of the compound. The activity was useful for reduction of ClO_4^- in water to lower concentration level. The removal reactions were not interfered by other general anions. The compound was conveniently regenerated by general chemical procedure from the obtained product after the ClO_4^- removal experiments. While many anion-exchange resins have been developed, utilizations of positively charged nano-scale molecules as anion-exchange materials were not explored well. This work reveals that coordination compounds have high potentials as new anion-exchange materials.

20

We thank Y. Yamamoto of Research Institute of Green Science Technology in Shizuoka University for support in obtaining the elemental analysis data. This work was supported by a grant from the New Energy and Industrial Technology Development Organization (NEDO) in Japan (No. 07A20008a).

30

Notes and references

^a Department of Chemistry, Faculty of Science, Shizuoka University, 836 Ohya, Suruga-ku, Shizuoka, 422-8529 Japan

^b Graduate School of Science and Technology, Shizuoka University, 836 Ohya, Suruga-ku, Shizuoka 422-8529 Japan.

^c Green Chemistry Research Division, Research Institute of Green Science Technology, Shizuoka University, 836 Ohya, Suruga-ku, Shizuoka 422-8529, Japan. scmkond@ipc.shizuoka.ac.jp

^d Research and Development Department, Central Laboratory, Maezawa Industries, Inc., 5-11, Naka-cho, Kawaguchi, Saitama 332-8556, Japan

[†] Electronic supplementary information (ESI) available: Materials and methods, crystallographic data, XRD charts, UV-vis. reflectance spectra, IR spectra. 854022 (1-SO₄), CCDC 854021 (1-ClO₄). For ESI and crystallographic data in CIF or other electronic format see DOI: [xx.xxxx/xxxxxxxxxx](https://doi.org/10.1039/C4XX00000X).

1. (a) J. Wolff, *Pharmacol. Rev.*, 1998, **50**, 89-105. (b) ITRC, ITRC (Interstate Technology & Regulatory Council) Perchlorate: Overview of Issues, Status, and Remedial Options. PERCHLORATE-1. Interstate Technology & Regulatory Council, Perchlorate Team, Washington, DC Available on the Internet at <http://www.itrcweb.org/>. 2005.
2. D. R. Parker, A. L. Seyffert and B. K. Reese, *Environ. Sci. Technol.*, 2008, **42**, 1465-1471.
3. P. B. Hatzinger, *Environ. Sci. Technol.*, 2005, **39**, 239A-247A.
4. M. Han, D. M. Engelhard and G. H. Clever, *Chem. Soc. Rev.*, 2014, **43**, 1848-1860
5. (a) D. A. McMorran, P. J. Steel, *Angew. Chem. Int. Ed.* 1998, **37**, 3295-3297. (b) L. J. Barbour, G. W. Orr and J. L. Atwood, *Nature*, 1998, **393**, 671-673. (c) D. K. Chand, K. Biradha and M. Fujita, *Chem. Commun.*, 2001, 1652-1653. (d) N. Yue, Z. Qin, M. C. Jennings, D. J. Eisler and R. J. Puddephatt, *Inorg. Chem. Commun.*, 2003, **6**, 1269-1271.
6. C.-Y. Su, Y.-P. Cai, C.-L. Chen, H.-X. Zhang and B.-S. Kang, *J. Chem. Soc., Dalton Trans.*, 2001, 359-361.
7. (a) H.-K. Liu, J. Hu, T.-W. Wang, X.-L. Yu, J. Liu and B. Kang, *J. Chem. Soc., Dalton Trans.*, 2001, 3534-3540. (b) C.-Y. Su, Y.-P. Cai, C.-L. Chen, M. D. Smith, W. Kaim and H.-C. zur Loye, *J. Am. Chem. Soc.*, 2003, **125**, 8595-8613. (d) H. Amouri, C. Desmaret, A. Bettoschi, M. N. Rager, K. Boubekeur, P. Rabu and M. Drillon, *Chem. Eur. J.*, 2007, **13**, 5401-5407.
8. (a) P. Liao, B. W. Langloss, A. M. Johnson, E. R. Knudsen, F. S. Tham, R. Julian and R. J. Hooley, *Chem. Commun.*, 2010, **46**, 4932-4934. (b) N. Kishi, Z. Li, K. Yoza, M. Akita and M. Yoshizawa, *J. Am. Chem. Soc.*, 2011, **133**, 11438-11441.
9. (a) Y.-B. Dong, P. Wang, J.-P. Ma, X.-X. Zhao, H.-Y. Wang, B. Tang and R.-Q. Huang, *J. Am. Chem. Soc.*, 2007, **129**, 4872-4873. (b) G. H. Clever, S. Tashiro and M. Shionoya, *Angew. Chem., Int. Ed.*, 2009, **48**, 7010-7012. (c) G. H. Clever, W. Kawamura, S. Tashiro, M. Shiro and M. Shionoya, *Angew. Chem., Int. Ed.*, 2012, **51**, 2606-2609. (d) J. E. M. Lewis, E. L. Gavey, S. A. Cameron and J. D. Crowley, *Chem. Sci.*, 2012, **3**, 778-784.
10. (a) H.-K. Liu, Y. Cai, W. Luo, F. Tong, C. You, H. Lü, X. Huang, H.-YY e, F. Su, X. Wang, *Inorg. Chem. Commun.*, 2009, **12**, 457-460. (b) W.-Y. Sun, J. Fan, T. Okamura, J. Xie, K.-B. Yu, N. Ueyama, *Chem. Eur. J.*, 2001, **7**, 2557-2562.
11. (a) T. Hirakawa, M. Yamaguchi, N. Ito, M. Miyazawa, N. Nishina, M. Kondo, R. Ikeya, S. Yasue, K. Maeda and F. Uchida, *Chem. Lett.*, 2009, **38**, 290-291. (b) M. Yamaguchi, T. Hirakawa, N. Nishina, M. Kondo, H. Aoki, E. Okuda, L. Zhang, *Chem. Lett.* 2010, **39**, 1192-1193.
12. Purolite A530E Microporous strong base anion exchange resin. <http://www.purolite.com/default.aspx?RelID=606288&ProductID=333>. The ClO_4^- removal activity of the resin was drastically increased by grinding it. The anion removal result by using anion-exchange resin without grinding is shown in Supplementary Information (Supporting Information).
13. B. Gu, Y.-K. Ku, G. M. Brown, *Environ. Sci. Technol.* 2005, **39**, 901-907.
14. World Health Organization 2011, Guidelines for drinking-water quality, fourth edition. http://www.who.int/water_sanitation_health/publications/2011/dwg_guidelines/en/index.html
15. D. P. Chen, C. Yu, C.-Y. Chang, Y. Wan, J. M. J. Frechet, W. A. Goddard III, M. S. Diallo, *Environ. Sci. Technol.* 2012, **46**, 10718-10726.

105