

# High-Quality Draft Genome Sequence of *Burkholderia contaminans* CH-1, a Gram-Negative Bacterium That Metabolizes 2-Azahypoxanthine, a Plant Growth-Regulating Compound

メタデータ	言語: eng 出版者: 公開日: 2018-01-11 キーワード (Ja): キーワード (En): 作成者: Choi, Jae-Hoon, Sugiura, Hikaru, Moriuchi, Ryota, Kawagishi, Hirokazu, Dohra, Hideo メールアドレス: 所属:
URL	<a href="http://hdl.handle.net/10297/00024384">http://hdl.handle.net/10297/00024384</a>



# High-Quality Draft Genome Sequence of *Burkholderia contaminans* CH-1, a Gram-Negative Bacterium That Metabolizes 2-Azahypoxanthine, a Plant Growth-Regulating Compound

Jae-Hoon Choi,<sup>a,b</sup> Hikaru Sugiura,<sup>b</sup> Ryota Moriuchi,<sup>a</sup> Hirokazu Kawagishi,<sup>a,b,c</sup> Hideo Dohra<sup>a,d</sup>

Research Institute of Green Science and Technology, Shizuoka University, Shizuoka, Japan<sup>a</sup>; College of Agriculture, Academic Institute, Shizuoka University, Shizuoka, Japan<sup>b</sup>; Graduate School of Science and Technology, Shizuoka University, Shizuoka, Japan<sup>c</sup>; Department of Biological Science, Graduate School of Integrated Science and Technology, Shizuoka University, Shizuoka, Japan<sup>d</sup>

**ABSTRACT** *Burkholderia contaminans* strain CH-1 converts 2-azahypoxanthine to 2-aza-8-oxohypoxanthine, plant growth-regulating compounds, by oxidation. We report here the high-quality draft genome sequence of *B. contaminans* CH-1. The genome contains 8,065 protein-coding sequences, including several genes possibly involved in metabolizing 2-azahypoxanthine.

**B***urkholderia contaminans* belongs to the *Burkholderia cepacia* complex, which comprises approximately 20 closely related species (1). We recently reported that *B. contaminans* strain CH-1, isolated from an airborne-contaminated nutrient medium, showed a high level of activity in converting 2-azahypoxanthine (AHX) to 2-aza-8-oxohypoxanthine (AOH). Both compounds were novel plant growth-regulating compounds related to fairy ring formation (2).

Genomic DNA of *B. contaminans* strain CH-1 was extracted using the DNeasy blood and tissue kit and fragmented using the Covaris acoustic solubilizer. A paired-end library constructed by the TruSeq DNA PCR-free library preparation kit was sequenced using the Illumina MiSeq platform (301-bp paired-end) at the Instrumental Research Support Office, Research Institute of Green Science and Technology, Shizuoka University. The raw read sequences were cleaned up using Trimmomatic (3) by trimming adapter sequences, low-quality ends (quality score, <15), the last 301 bases, and reads less than 150 bp. The resultant 2,052,682 high-quality read pairs, totaling 1,012 Mb and corresponding to an approximately 114-fold coverage of the genome, were assembled using SPAdes version 3.10.0 (4) with a default set of *k*-mer sizes and options (–careful, –only-assembler, and –cov-cutoff auto); contigs less than 200 bp were eliminated. Average nucleotide identity (ANI) analysis (5, 6) of the resultant 94 contigs showed an extremely high ANI value (99.98%) with the improved high-quality draft genome of *B. contaminans* LMG 23361 (GenBank accession number MCAU02000000) (7).

To construct scaffolds of the CH-1 genome, the contigs were aligned to the LMG 23361 genome sequence using CONTIGuator version 2.7 (8). The pseudocontigs were manually curated by alignments with contigs and the LMG 23361 genome, using BLAST and Geneious version 10.2.2 (9), and then circularized using Circlator version 1.1.2 (10). The high-quality draft genome of *B. contaminans* CH-1 contained three circular chromosomes, in which only five gaps remained, as well as a putative linear plasmid (designated pBC453), totaling 8,901,643 bp with a G+C content of 66.0%. The CH-1 genome was annotated using the DFAST-core stand-alone program version 0.9.4 (11)

**Received** 13 September 2017 **Accepted** 18 September 2017 **Published** 12 October 2017

**Citation** Choi J-H, Sugiura H, Moriuchi R, Kawagishi H, Dohra H. 2017. High-quality draft genome sequence of *Burkholderia contaminans* CH-1, a Gram-negative bacterium that metabolizes 2-azahypoxanthine, a plant growth-regulating compound. *Genome Announc* 5:e01148-17. <https://doi.org/10.1128/genomeA.01148-17>.

**Copyright** © 2017 Choi et al. This is an open-access article distributed under the terms of the [Creative Commons Attribution 4.0 International license](https://creativecommons.org/licenses/by/4.0/).

Address correspondence to Hirokazu Kawagishi, [kawagishi.hirokazu@shizuoka.ac.jp](mailto:kawagishi.hirokazu@shizuoka.ac.jp), or Hideo Dohra, [dora.hideo@shizuoka.ac.jp](mailto:dora.hideo@shizuoka.ac.jp).

and manually curated. The CH-1 genome contains 8,065 protein-coding sequences, including 85 possible pseudogenes (1.05%), 18 rRNAs (6 clusters of 5S, 16S, and 23S rRNAs), and 82 tRNAs. The proteins encoded by the CH-1 genome were additionally annotated using BlastKOALA (12) to characterize protein functions and reconstruct the purine metabolism pathway in the KEGG database (13). Among 8,065 proteins, 82 were assigned to the purine metabolism pathway (ko00230), including the xanthine dehydrogenase small subunit (XdhA, K13481) and large subunit (XdhB, K13482), which are possibly involved in the conversion of AHX to AOH, as predicted by the purine metabolism pathway in plants (14). In addition, three proteins were assigned to the xanthine dehydrogenase accessory factor (XdhC, K07402), which is required for xanthine dehydrogenase activity in *Rhodobacter capsulatus* (15). These genome data will help elucidate the mechanism for converting AHX to AOH and improve methods for the large-scale production of AOH for its practical use in agriculture.

**Accession number(s).** The high-quality draft genome sequence of *B. contaminans* CH-1 has been deposited in DDBJ/ENA/GenBank under the accession numbers AP018357 to AP018360 (four entries).

## ACKNOWLEDGMENT

This work was partially supported by a Grant-in-Aid for Young Scientists A (16H06192) from MEXT to Jae-Hoon Choi.

## REFERENCES

1. Vanlaere E, Baldwin A, Gevers D, Henry D, De Brandt E, LiPuma JJ, Mahenthalingam E, Speert DP, Dowson C, Vandamme P. 2009. Taxon K, a complex within the *Burkholderia cepacia* complex, comprises at least two novel species, *Burkholderia contaminans* sp. nov. and *Burkholderia lata* sp. nov. *Int J Syst Evol Microbiol* 59:102–111. <https://doi.org/10.1099/ijs.0.001123-0>.
2. Choi JH, Kikuchi A, Pumkiao P, Hirai H, Tokuyama S, Kawagishi H. 2016. Bioconversion of AHX to AOH by resting cells of *Burkholderia contaminans* CH-1. *Biosci Biotechnol Biochem* 80:2045–2050. <https://doi.org/10.1080/09168451.2016.1189314>.
3. Bolger AM, Lohse M, Usadel B. 2014. Trimmomatic: a flexible trimmer for Illumina sequence data. *Bioinformatics* 30:2114–2120. <https://doi.org/10.1093/bioinformatics/btu170>.
4. Bankevich A, Nurk S, Antipov D, Gurevich AA, Dvorkin M, Kulikov AS, Lesin VM, Nikolenko SI, Pham S, Prjibelski AD, Pyshkin AV, Sirotkin AV, Vyahhi N, Tesler G, Alekseyev MA, Pevzner PA. 2012. SPAdes: a new genome assembly algorithm and its applications to single-cell sequencing. *J Comput Biol* 19:455–477. <https://doi.org/10.1089/cmb.2012.0021>.
5. Goris J, Konstantinidis KT, Klappenbach JA, Coenye T, Vandamme P, Tiedje JM. 2007. DNA-DNA hybridization values and their relationship to whole-genome sequence similarities. *Int J Syst Evol Microbiol* 57:81–91. <https://doi.org/10.1099/ijs.0.64483-0>.
6. Rodriguez-R LM, Konstantinidis KT. 2016. The enveomics collection: a toolbox for specialized analyses of microbial genomes and metagenomes. *PeerJ* 4:e1900v1.
7. Jung JY, Ahn Y, Kweon O, LiPuma JJ, Hussong D, Marasa BS, Cerniglia CE. 2017. Improved high-quality draft genome sequence and annotation of *Burkholderia contaminans* LMG 23361T. *Genome Announc* 5(16):e00245-17. <https://doi.org/10.1128/genomeA.00245-17>.
8. Galardini M, Biondi EG, Bazzicalupo M, Mengoni A. 2011. CONTIGuator: a bacterial genomes finishing tool for structural insights on draft genomes. *Source Code Biol Med* 6:11. <https://doi.org/10.1186/1751-0473-6-11>.
9. Kearse M, Moir R, Wilson A, Stones-Havas S, Cheung M, Sturrock S, Buxton S, Cooper A, Markowitz S, Duran C, Thierer T, Ashton B, Meintjes P, Drummond A. 2012. Geneious Basic: an integrated and extendable desktop software platform for the organization and analysis of sequence data. *Bioinformatics* 28:1647–1649. <https://doi.org/10.1093/bioinformatics/bts199>.
10. Hunt M, Silva ND, Otto TD, Parkhill J, Keane JA, Harris SR. 2015. Circlator: automated circularization of genome assemblies using long sequencing reads. *Genome Biol* 16:294. <https://doi.org/10.1186/s13059-015-0849-0>.
11. Tanizawa Y, Fujisawa T, Kaminuma E, Nakamura Y, Arita M. 2016. DFAST and DAGA: web-based integrated genome annotation tools and resources. *Food Health* 35:173–184. <https://doi.org/10.12938/bmfh.16-003>.
12. Kanehisa M, Sato Y, Morishima K. 2016. BlastKOALA and GhostKOALA: KEGG tools for functional characterization of genome and metagenome sequences. *J Mol Biol* 428:726–731. <https://doi.org/10.1016/j.jmb.2015.11.006>.
13. Kanehisa M, Goto S, Sato Y, Kawashima M, Furumichi M, Tanabe M. 2014. Data, information, knowledge and principle: back to metabolism in KEGG. *Nucleic Acids Res* 42:D199–D205. <https://doi.org/10.1093/nar/gkt1076>.
14. Choi JH, Ohnishi T, Yamakawa Y, Takeda S, Sekiguchi S, Maruyama W, Yamashita K, Suzuki T, Morita A, Ikka T, Motohashi R, Kiriwa Y, Tobina H, Asai T, Tokuyama S, Hirai H, Yasuda N, Noguchi K, Asakawa T, Sugiyama S, Kan T, Kawagishi H. 2014. The source of “fairy rings”: 2-azahypoxanthine and its metabolite found in a novel purine metabolic pathway in plants. *Angew Chem Int Ed Engl* 53:1552–1555. <https://doi.org/10.1002/anie.201308109>.
15. Leimkühler S, Klipp W. 1999. Role of XDHC in molybdenum cofactor insertion into xanthine dehydrogenase of *Rhodobacter capsulatus*. *J Bacteriol* 181:2745–2751.